

Lyapunov Exponents for the Parabolic Anderson Model

M. Cranston , T. S. Mountford , T. Shiga *

March 22, 2002

Abstract

We consider the asymptotic almost sure behavior of the solution of the equation

$$\begin{aligned} u(t, x) &= u_0(x) + \kappa \int_0^t \Delta u(s, x) ds + \int_0^t u(s, x) \partial B_x(s) \\ u(0, x) &= u_0(x) \end{aligned}$$

where $\{B_x : x \in \mathbf{Z}^d\}$ is a field of independent Brownian motions.

Keywords and phrases: parabolic Anderson model, Feynman-Kac formula, Lyapunov exponents, Borell's inequality, percolation.

1 Introduction

We make an asymptotic study of the Lyapunov exponent of the parabolic Anderson model with white noise potential. Start with a collection of independent, one-dimensional, Brownian motions $\{B_x(t), x \in \mathbf{Z}^d, t \geq 0\}$. This provides a random environment defined on a probability space (Ω, \mathcal{F}, Q) . Fixing a $\kappa > 0$ denote by $(X(t), t \geq 0, \mathcal{F}_t, P_x)$ the symmetric random walk on \mathbf{Z}^d with jump rate κ . That is, under P_x , $\{X(t), t \geq 0\}$ is the pure jump Markov process on \mathbf{Z}^d started at x with generator $\kappa\Delta$ where Δ is the discrete Laplacian defined by $\Delta f(x) = \frac{1}{2d} \sum_{|y-x|=1} (f(y) - f(x))$. Then we consider the solution of the stochastic

*Research of the first two authors supported in part by a grant from NSF, of the third author by a grant from Grant-in Aid for Scientific Research.

equation

$$u(t, x) = u_0(x) + \kappa \int_0^t \Delta u(s, x) ds + \int_0^t u(s, x) \partial B_x(s) \quad (1.1)$$

$$u(0, x) = u_0(x) \quad (1.2)$$

where ∂ denotes the Stratonovich differential. A result of Carmona and Molchanov [CM94] is that provided u_0 is bounded, the solution is given by means of the Feynman-Kac formula

$$u(t, x) = E_x[u_0(X(t))e^{\int_0^t dB_{X(s)}(t-s)}].$$

For important background material on the significance of the model we refer the reader to the memoir [CM94] and the references therein. The equation (1.1) and its variants arise in a variety of circumstances. Equation (1.1) can be written in differential form as

$$\frac{\partial u}{\partial t} = \kappa \Delta u + V u$$

with a potential $V = \partial B$. The elliptic version of this equation originated in the work of Anderson on entrapment of electrons in crystals with impurities. The parabolic equation can be viewed as a scalar version of magnetic fields in the presence of turbulence as described in Molcahnov and Ruzmaikin [MR94] and also has an interpretation as a population model.

When $\kappa = 0$, the solution is simply $u(t, x) = u_0(x)e^{-B_x(t)}$. In this case, $\lim_{t \rightarrow \infty} \frac{1}{t} \log u(t, x) = 0$. However, when $\kappa > 0$ the situation is quite different. A result of [CM94] is that when u_0 has compact support, then

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log u(t, x) = \gamma(\kappa), \quad Q \text{ a.s.}$$

The positive constant $\gamma(\kappa)$ is called the Lyapunov exponent. This was proven by a subadditivity argument which didn't extend to the noncompact support case. Moreover, $\gamma(\kappa)$ is independent of u_0 and a principal result of [CM94] was the existence of constants c_1, c_2 such that for all small $\kappa > 0$,

$$c_1 / \log \frac{1}{\kappa} \leq \gamma(\kappa) \leq c_2 \log \log \frac{1}{\kappa} / \log \frac{1}{\kappa}.$$

This was improved in their later work [CMV94] to

$$c_1 / \log \frac{1}{\kappa} \leq \gamma(\kappa) \leq c_2 / \log \frac{1}{\kappa},$$

again for small κ . That is, γ is continuous at 0 but increases dramatically with κ . In the present paper we remove the restriction that u_0 have compact support by showing

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log u(t, x) = \gamma(\kappa)$$

when u_0 is a bounded nonnegative, not identically 0 function and that again $\gamma(\kappa)$ is independent of u_0 . This will follow from the case where u_0 has compact support and a recurrence property of oriented percolation. We also show

$$\lim_{\kappa \downarrow 0} \gamma(\kappa) \log \frac{1}{\kappa} = c$$

where we identify c by means of a subadditive ergodic theory argument. In fact, this is where our results differ in approach from earlier work on the subject. Our approach was developed in [Mou01] in dealing with survival probabilities of the random walk $X(t)$ in a disastrous environment. In the present context this relies on introducing the space of deterministic paths on which the measure P_x is concentrated when the paths of X are restricted to have a specific number of jumps. For an interval $[a, b] = I \subset [0, \infty)$ and $\gamma : I \rightarrow \mathbf{Z}$ let $N(\gamma, I)$ denote the number of jumps of γ in the interval I . We shall only deal with paths γ which are right continuous with left limits. Then for $x \in \mathbf{Z}^d$ define

$$\Gamma_{I,j}^x = \{\gamma : I \rightarrow \mathbf{Z}^d, \gamma(a) = x, |\gamma(s) - \gamma(s-)| \leq 1, \text{ for all } s \in I, N(\gamma, I) = j\}.$$

We also define $\Gamma_{I,j}^{x,\epsilon}$ to be the intersection of $\Gamma_{I,j}^x$ with those paths with jump times separated from one another by ϵ and such that there are no jump times within $\frac{\epsilon}{2}$ of the boundary of I . Now we consider the functional

$$A_{I,j}^x = \sup_{\gamma \in \Gamma_{I,j}^x} \int_a^b dB_{\gamma(s)}(b-s).$$

The corresponding functional when restricting jumps to be ϵ apart will be

$$A_{I,j}^{x,\epsilon} = \sup_{\gamma \in \Gamma_{I,j}^{x,\epsilon}} \int_a^b dB_{\gamma(s)}(b-s).$$

Obviously, $A_{I,j}^x \geq A_{I,j}^{x,\epsilon}$. The functional $A_{I,j}^{x,\epsilon}$ will be important in establishing the lower bound in **Theorem 2.14**. We simplify the notation a little by first setting $A_{0,n} = A_{[0,n],n}^0$. The fact that time is running in opposite directions in the functional A makes it a little clumsy to work with. Instead we define a functional with the same distribution. This will suffice to give all the conclusions we need about A . Thus, we set

$$\vec{A}_{I,j}^x = \sup_{\gamma \in \Gamma_{I,j}^x} \int_a^b dB_{\gamma(s)}(s).$$

Make the simplified notation $\vec{A}_{[0,n],n}^0 = \vec{A}_{0,n}$. Similarly, set

$$\vec{A}_{I,j}^{x,\epsilon} = \sup_{\gamma \in \Gamma_{I,j}^{x,\epsilon}} \int_a^b dB_{\gamma(s)}(s)$$

and use the simplified notation $\vec{A}_{0,n}^\epsilon = \vec{A}_{[0,n],n}^{0,\epsilon}$. One thing to notice here is that time in the dB and γ terms are running in the same direction for the functionals denoted by \vec{A} . A very important fact is a scaling relation for the functionals $A_{[0,n],j}$ and $\vec{A}_{[0,n],j}$ inherited from the scaling properties of the Brownian field $\{B_x(t) : t \geq 0, x \in \mathbf{Z}^d\}$. Namely, for $j, n \in \mathbf{Z}$, (using the notation $\stackrel{\mathcal{L}}{=}$ for equality in law)

$$\begin{aligned}
A_{[0,n],jn}^0 &= \sup_{\gamma \in \Gamma_{[0,n],jn}^0} \int_0^n dB_{\gamma(s)}(n-s) \\
&= \sup_{\gamma \in \Gamma_{[0,n],jn}^0} \int_0^{jn} dB_{\gamma(\frac{s}{j})}(n-\frac{s}{j}) \\
&= \sup_{\gamma \in \Gamma_{[0,jn],jn}^0} \int_0^{jn} dB_{\gamma(s)}(\frac{1}{j}(jn-s)) \\
&\stackrel{\mathcal{L}}{=} \frac{1}{\sqrt{j}} \sup_{\gamma \in \Gamma_{[0,jn],jn}^0} \int_0^{jn} dB_{\gamma(s)}(jn-s) \\
&= \frac{1}{\sqrt{j}} A_{[0,jn],jn}^0.
\end{aligned}$$

A similar computation gives the same scaling relation for $\vec{A}_{[0,n],jn}$.

Another observation is that $\vec{A}_{0,n}$ is the supremum of the centered Gaussian field $\{\int_a^b dB_{\gamma(s)}(s) : \gamma \in \Gamma_{[0,n],n}^0\}$. Thus, results such as Borell's inequality are available for our analysis. For the reader's convenience, we state the results from the theory of Gaussian processes which we shall use. Both results may be found in Adler [Ad90]

Theorem 1.1. (*C. Borell*)

Let T be a Polish space and $\{X_t\}_{t \in T}$ be a centered separable Gaussian field with $\sup_{t \in T} X_t < \infty$ a.s.. Then $E(\sup_{t \in T} X_t) < \infty$ and for all $\lambda > 0$

$$P\left(\left|\sup_{t \in T} X_t - E\left(\sup_{t \in T} X_t\right)\right| > \lambda\right) \leq 2e^{-\lambda^2/2\sigma_T^2},$$

where $\sigma_T^2 = \sup_{t \in T} E(X_t^2)$.

Let $(X_t)_{t \in T}$ be a centered separable Gaussian field with the pseudo-metric ρ on T ;

$$\rho(t, s) = \sqrt{E(X_t - X_s)^2}.$$

Theorem 1.2. (*Fernique-Talagrand*) There exists a universal constant $K > 0$ such that

$$E(\sup_{t \in T} X_t) \leq K \int_0^\infty \sqrt{\log N(\epsilon)} d\epsilon$$

where $N(\epsilon)$ is the least number of ρ -balls of radius ϵ required to cover T .

Next, given $m, n \in \mathbf{Z}^+$ determine $x \in \mathbf{Z}^d$ as follows, let x be the smallest value of $\gamma(m)$ under any well ordering of \mathbf{Z}^d taken among paths γ which achieve the maximum for $\int_0^m dB_{\gamma(s)}(s)$. Then set $\vec{A}_{m,m+n} = \vec{A}_{[m,m+n],n}^x$. Since $\vec{A}_{0,m+n}$ is a supremum over paths which have $m+n$ jumps on the interval $[0, m+n]$ which doesn't impose any restriction on how many jumps occur in $[0, m]$ or $[m, m+n]$ it follows that

$$\vec{A}_{0,m+n} \geq \vec{A}_{0,m} + \vec{A}_{m,m+n}.$$

Then by the subadditive ergodic theorem (applied to $-\vec{A}_{0,n}$) we have the following

Theorem 1.3. *There is a positive α such that*

$$\lim_{n \rightarrow \infty} \frac{\vec{A}_{0,n}}{n} = \alpha.$$

For $\epsilon > 0$ there is an $\alpha(\epsilon)$ such that

$$\lim_{n \rightarrow \infty} \frac{\vec{A}_{0,n}^\epsilon}{n} = \alpha(\epsilon).$$

Moreover, $\lim_{\epsilon \downarrow 0} \alpha(\epsilon) = \alpha$.

Proof. The conditions of Liggett's subadditive ergodic theorem [Lig85] apply. The positivity of α is already apparent if one considers the supremum of $\int_0^t dB_{\gamma(s)}(s)$ over paths which are constrained to visit only two adjacent sites in \mathbf{Z}^d . The finiteness of α follows from the entropy bound in **Theorem 1.2**. Namely, for some universal constant K ,

$$E_Q[\vec{A}_{0,n}] \leq K \int_0^\infty \sqrt{\log N(\epsilon)} d\epsilon,$$

where $N(\epsilon)$ is the smallest number of ϵ balls required to cover $\Gamma_{[0,n],n}^0$ using balls with respect to the canonical metric ρ . By scaling, $E_Q[\vec{A}_{0,n}] = \sqrt{n} E_Q[\vec{A}_{[0,1],n}]$. Thus we will apply the entropy bound to $E_Q[\vec{A}_{[0,1],n}]$. Let W_n be the set of n -step random walk paths on \mathbb{Z}^d and let

$$T_n = \{t = (t_1, \dots, t_n) : 0 < t_1 \leq \dots \leq t_n \leq 1\}.$$

Identify $\Gamma_{[0,1],n}$ with $W_n \times T_n$, then it holds that

$$\begin{aligned} \rho((w, t), (\tilde{w}, s))^2 &= E_Q \left[\sum_{k=0}^n (B_{w_k}(t_{k+1}) - B_{w_k}(t_k)) - \sum_{l=0}^n (B_{\tilde{w}_l}(s_{l+1}) - B_{\tilde{w}_l}(s_l)) \right]^2 \\ &= 2 \sum_{k=1}^n \sum_{l=1}^n \delta_{w_k \tilde{w}_l} |(t_k, t_{k+1}) \cap (s_l, s_{l+1})| \\ &\leq 2, \end{aligned}$$

with $t_{n+1} = s_{n+1} = 1$. As is easily seen, the ρ balls of radius $\sqrt{2\epsilon n}$ centered at the points

$$\{(w, (\epsilon k_1, \dots, \epsilon k_n)) \in W_n \times \epsilon \mathbb{Z}_+^n : 1 \leq k_1 \leq k_2 \leq \dots \leq k_n \leq \frac{1}{\epsilon}\}$$

cover $W_n \times T_n$. And

$$N(\sqrt{2\epsilon n}) \leq (2d)^n \binom{[1/\epsilon] + n}{n},$$

so that there is a constant $M > 0$ such that for any $\epsilon > 0$

$$N(\epsilon) \leq (2d)^n \binom{[2n/\epsilon^2] + n}{n} \leq M^n \epsilon^{-2n}.$$

Hence

$$\int_0^\infty \sqrt{\log N(\epsilon)} d\epsilon \leq \sqrt{n} \int_0^2 \sqrt{\log \frac{M}{\epsilon^2}} d\epsilon \leq C\sqrt{n}.$$

Thus we obtain

$$E_Q[\vec{A}_{[0,n],n}] \leq Cn,$$

which implies

$$\alpha = \lim_{n \rightarrow \infty} \frac{1}{n} E_Q[\vec{A}_{[0,n],n}] = \sup_n \frac{1}{n} E_Q[\vec{A}_{[0,n],n}] \leq C.$$

The proof for the second claim is entirely analogous. The proof that $\lim_{\epsilon \downarrow 0} \alpha(\epsilon) = \alpha$ is straightforward and is omitted. \square

From the scaling property we have immediately

Corollary 1.4. *For any $j \in \mathbf{Z}$*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \vec{A}_{[0,n],jn}^0 = \sqrt{j}\alpha, \text{ } Q \text{ a. s. .}$$

The Corollary quantifies the improvement in the functional $\vec{A}_{[0,n],jn}^0$ as a function of j : with more jumps, a greater value is achieved. Now when considering which paths of the Markov process X contribute the principal term in $u(t, x)$, paths with jt jumps give a potentially greater value for $\int_0^t dB_{X(s)}(t-s)$ but the probability of making jt jumps in $[0, t]$ decreases with j for j large. Our work shows that j^*t jumps, with

$$j^* = \frac{\alpha^2}{4 \log^2 \frac{1}{\kappa}}$$

is the optimal number of jumps during $[0, t]$ in terms of balancing payoff $\int_0^t dB_{X(s)}(t-s)$ versus probability. Thus, the principal contribution in the Feynman-Kac representation for $u(t, x)$ arises from paths with j^*t jumps.

2 Existence of a limit for $\frac{\log u(t,x)}{t}$

We aim to establish the existence, for arbitrary bounded, nonnegative u_0 , of the

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log u(t, x), \quad Q \text{ a. s.}$$

In fact, we shall assume without loss of generality that $u_0(0) > 0$. Recall, this was established by Carmona and Molchanov in the case of compactly supported u_0 . We first establish an existence result for a functional with the same distribution as $u(t, x)$ when u_0 is taken to be identically 1. In this section we treat explicitly the case where dimension $d = 1$. This is simply to minimize notation. It will be clear that the arguments easily extend to higher dimensions. Also we make use of some large deviations results, **Theorem 1.1** and **Theorem 2.11**.

Theorem 2.1. *There is a $\gamma_1(\kappa)$ such that for any $x \in \mathbf{Z}^d$*

$$\gamma_1(\kappa) = \lim_{t \rightarrow \infty} \frac{1}{t} \log E_x[e^{\int_0^t dB_{X(s)}(s)}], \quad Q \text{ a. s.}$$

Proof. By translation invariance we may take x equal to 0. Set

$$\begin{aligned} v(t, y) &= E_0[e^{\int_0^t dB_{X(s)}(s)} 1_y(X(t))] \\ Z(t) &= \max_{y \in \mathbf{Z}} v(t, y) \end{aligned}$$

The functional $\log Z(t)$ is superadditive and Liggett's subadditive ergodic Theorem (applied to $-\log Z(t)$) gives the *Q a. s.* existence of

$$\lim_{t \rightarrow \infty} \frac{\log Z(t)}{t} = \gamma_1(\kappa)$$

for some constant $\gamma_1(\kappa)$. From its definition we have that $Z(t) \geq E_0[e^{\int_0^t dB_{X(s)}(s)} 1_0(X(t))]$ and so (recalling the results of Carmona and Molchanov [?] mentioned in the **Introduction**) we have that $\gamma_1(\kappa) \geq \gamma(\kappa) > 0$.

However,

$$\log E_0[e^{\int_0^t dB_{X(s)}(s)}] \geq \log Z(t)$$

so

$$\liminf_{t \rightarrow \infty} \frac{1}{t} \log E_0[e^{\int_0^t dB_{X(s)}(s)}] \geq \gamma_1(\kappa), \quad Q \text{ a. s.}$$

On the other hand,

$$\begin{aligned}
\frac{1}{t} \log E_0[e^{\int_0^t dB_{X(s)}(s)}] &= \frac{1}{t} \log (E_0[\Sigma_{|x| \leq t^2} e^{\int_0^t dB_{X(s)}(s)} \mathbf{1}_x(X(t))]) \\
&+ E_0[e^{\int_0^t dB_{X(s)}(s)} \mathbf{1}_{(t^2, \infty)}(|X(t)|)] \\
&\leq \frac{1}{t} (\log(ct^2 e^{Z(t)}) \\
&+ E_0[e^{\int_0^t dB_{X(s)}(s)} \mathbf{1}_{(t^2, \infty)}(|X(t)|)]) .
\end{aligned}$$

But,

$$\begin{aligned}
E_Q[E_0[e^{\int_0^t dB_{X(s)}(s)} \mathbf{1}_{(t^2/2, \infty)}(|X(t)|)]^2] &\leq P_0(|X(t)| \geq t^2/2) E_Q[E_0[e^{2 \int_0^t dB_{X(s)}(s)}]] \\
&\leq e^{-ct^2} E_Q[E_0[e^{2 \int_0^t dB_{X(s)}(s)}]] \\
&\leq e^{-ct^2} ,
\end{aligned}$$

for another value of c . Chebychev's inequality then implies

$$Q(E_0[e^{\int_0^t dB_{X(s)}(s)} \mathbf{1}_{(t^2/2, \infty)}(|X(t)|)] > \frac{1}{K}) \leq e^{-ct^2}$$

for K very large but fixed. So by Borel-Cantelli,

$$Q(E_0[e^{\int_0^n dB_{X(s)}(s)} \mathbf{1}_{(n^2/2, \infty)}(|X(n)|)] > \frac{1}{K}, i.o.) = 0.$$

Now define a stopping time T with respect to $\mathcal{F}_t = \sigma\{B_x(s) : x \in \mathbf{Z}, 0 \leq s \leq t\}$ by $T = \inf\{t \geq N_0 : E_0[e^{\int_0^t dB_{X(s)}(s)} \mathbf{1}_{(t^2, \infty)}(|X(t)|)] \geq 1\}$.

A simple two moment argument then gives that on the event $\{T < \infty\}$, for $T \in (n-1, n]$, one has $Q(E_0[e^{\int_0^n dB_{X(s)}(s)} \mathbf{1}_{(n^2/2, \infty)}(|X(n)|)] < \frac{1}{K} | \mathcal{F}_T)$ tends to 0 as N_0 tends to ∞ . Letting N_0 tend to infinity one obtains that Q a.s. there exists t_0 so that

$$E_0[e^{\int_0^t dB_{X(s)}(s)} \mathbf{1}_{(t^2, \infty)}(|X(t)|)] \leq 1, \text{ for } t \geq t_0$$

Thus, for t large,

$$\begin{aligned}
\frac{1}{t} \log E_0[e^{\int_0^t dB_{X(s)}(s)}] &\leq \frac{1}{t} \log(ct^2 e^{Z(t)} + 1) \\
&\leq \frac{Z(t)}{t} + \frac{O(\log t)}{t}.
\end{aligned}$$

Consequently, Q a.s.,

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log E_0[e^{\int_0^t dB_{X(s)}(s)}] \leq \gamma_1(\kappa).$$

and therefore,

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log E_0[e^{\int_0^t dB_{X(s)}(s)}] = \lim_{t \rightarrow \infty} \frac{1}{t} \log Z(t).$$

and the Theorem is proved. \square

Our goal of proving the existence of $\lim_{t \rightarrow \infty} \frac{1}{t} \log u(t, x)$, Q a. s., is enabled by the introduction of a few auxiliary functionals.

Set

$$Z(t, t^2) = \max_{x \in \mathbf{Z}^d, |x| \leq t^2} v(t, x).$$

Notice that $v(t, x) \stackrel{\mathcal{L}}{=} E_0[e^{\int_0^t dB_{X(s)}(t-s)} 1_x(X(t))]$ and we'll use information about v to get the result on u . We also remark that by translation invariance, $v(t, x-y) \stackrel{\mathcal{L}}{=} E_y[e^{\int_0^t dB_{X(s)}(s)} 1_x(X(t))]$

Theorem 2.2. *For any $x \in \mathbf{Z}^d$*

$$\gamma(\kappa) = \lim_{t \rightarrow \infty} \frac{1}{t} \log E_x[e^{\int_0^t dB_{X(s)}(s)}], \quad Q \text{ a. s.}$$

The proof will be broken up into a sequence of lemmas.

Lemma 2.3. *For M sufficiently large, if $a < 2$, then*

$$Q(E_0[e^{\int_0^t dB_{X(s)}(s)} 1_{[Mt, \infty)}(N(X, t)) \leq e^{at}] \geq 1 - e^{-(2+a)t}.$$

Proof. By Fubini and Cauchy-Schwarz,

$$\begin{aligned} E_Q[E_0[e^{\int_0^t dB_{X(s)}(s)} 1_{[Mt, \infty)}(N(X, t))]] &= E_0[E_Q[e^{\int_0^t dB_{X(s)}(s)} 1_{[Mt, \infty)}(N(X, t))]] \\ &\leq E_0[E_Q[e^{2 \int_0^t dB_{X(s)}(s)}]^{1/2} P_0(N(X, t) \in [Mt, \infty))^{1/2}] \\ &\leq e^t e^{-\frac{1}{2}(\epsilon\kappa - \kappa - M)t}. \end{aligned}$$

Then, provided M is large enough, Chebychev's inequality completes the proof. \square

Now fix an $\epsilon > 0$. In all that follows we shall assume, without loss of generality, that $Mt \in \mathbf{Z}$.

Lemma 2.4. *For M and t sufficiently large,*

$$Q(E_0[e^{\int_0^t dB_{X(s)}(s)} 1_{[0, Mt]}(N(X, t)) \geq e^{(\gamma_1(\kappa) - \frac{\epsilon}{100})t}] \geq 1 - \epsilon^{100}.$$

Proof. Using **Lemma 2.3**, with $a = -1$, and **Theorem 2.1**, we have for large t , with Q probability at least $1 - \epsilon^{100}$, that

$$\begin{aligned} E_0[e^{\int_0^t dB_{X(s)}(s)} 1_{[0, Mt]}(N(X, t))] &= E_0[e^{\int_0^t dB_{X(s)}(s)}] \\ &\quad - E_0[e^{\int_0^t dB_{X(s)}(s)} 1_{[Mt, \infty)}(N(X, t))] \\ &\geq e^{(\gamma_1(\kappa) - \frac{\epsilon}{50})t} - e^{-t}. \end{aligned}$$

and the lemma follows by a little algebra. \square

An immediate consequence is the following,

Corollary 2.5. *For t and M sufficiently large, given any $x \in \mathbf{Z}$, there exists an $x^*(t, x) \in \sigma(B_y(s) : y \in \mathbf{Z}, |y - x| \leq Mt, s \leq t)$ such that*

$$Q(E_x[e^{\int_0^t dB_{X(s)}(s)} 1_{[0, Mt]}(N(X, t)) 1_{x^*(t, x)}(X(t))] \geq e^{(\gamma_1(\kappa) - \frac{\epsilon}{99})t}) \geq 1 - \epsilon^{100}.$$

A more refined consequence is

Corollary 2.6. *Given any $x \in \mathbf{Z}$ there exist $x_+^* = x_+^*(t, x) > x$ and $x_-^* = x_-^*(t, x) < x$, with $x_{\pm}^*(t, x) \in \sigma(B_y(s) : y \in \mathbf{Z}, |y - x| \leq Mt, s \leq t)$ such that*

$$Q(E_x[e^{\int_0^t dB_{X(s)}(s)} 1_{[0, Mt]}(N(X, t)) 1_{x_+^*}(X(t))] \geq e^{(\gamma_1(\kappa) - \frac{\epsilon}{99})t}) \geq 1 - \epsilon^{50}$$

and

$$Q(E_x[e^{\int_0^t dB_{X(s)}(s)} 1_{[0, Mt]}(N(X, t)) 1_{x_-^*}(X(t))] \geq e^{(\gamma_1(\kappa) - \frac{\epsilon}{99})t}) \geq 1 - \epsilon^{50}$$

Proof. This is an easy use of the FKG inequalities and the observation that the events

$$\{E_x[e^{\int_0^t dB_{X(s)}(s)} 1_{[0, Mt]}(N(X, t)) 1_{x^*}(X(t))] \geq e^{(\gamma_1(\kappa) - \frac{\epsilon}{99})t}, \text{ for some } x^* > x\}$$

and

$$\{E_x[e^{\int_0^t dB_{X(s)}(s)} 1_{[0, Mt]}(N(X, t)) 1_{x^*}(X(t))] \geq e^{(\gamma_1(\kappa) - \frac{\epsilon}{99})t}, \text{ for some } x^* < x\}$$

are positively correlated, equally likely and the probability of their union is at least $1 - \epsilon^{100}$ by **Corollary 2.5**. \square

We can move the points x_{\pm}^* a little further away from x ,

Lemma 2.7. *For M and t sufficiently large there is a $c_1 = c_1(\kappa, M)$ such that for all $x \in \mathbf{Z}$ there exist $x_+^* = x_+^*(t, x) \in [x + \frac{\epsilon}{2}Mt, x + (1 - \frac{\epsilon}{2})Mt]$ and $x_-^* = x_-^*(t, x) \in [x - (1 - \frac{\epsilon}{2})Mt, x - \frac{\epsilon}{2}Mt]$, $x_{\pm}^*(t, x) \in \sigma(B_y(s) : y \in \mathbf{Z}, |y - x| \leq Mt, s \leq t)$ such that*

$$Q(E_x[e^{\int_0^t dB_{X(s)}(s)} 1_{[0, Mt]}(N(X, t)) 1_{x_+^*}(X(t))] \geq e^{(\gamma_1(\kappa) - c_1\epsilon)t}) \geq 1 - \epsilon^{49}$$

and

$$Q(E_x[e^{\int_0^t dB_{X(s)}(s)} 1_{[0, Mt]}(N(X, t)) 1_{x_-^*}(X(t))] \geq e^{(\gamma_1(\kappa) - c_1\epsilon)t}) \geq 1 - \epsilon^{49}.$$

Proof. By symmetry, we only need argue the first case. By **Corollary 2.6**, with probability at least $1 - \epsilon^{50}$, there is an $x_+^*(t(1 - \epsilon)) \in [x, x + (1 - \epsilon)Mt]$ such that

$$E_x[e^{\int_0^{t(1-\epsilon)} dB_{X(s)}(s)} 1_{[0, Mt(1-\epsilon)]}(N(X, t(1-\epsilon))) 1_{x_+^*(t(1-\epsilon))}(X(t(1-\epsilon)))] \geq e^{(\gamma_1(\kappa) - \frac{\epsilon}{99})(t-\epsilon)}.$$

If $x_+^*(t(1-\epsilon)) \in [x + \frac{\epsilon}{2}Mt, x + (1 - \frac{\epsilon}{2})Mt]$, we consider the constant path identically equal to $x_+^*(t(1-\epsilon))$ over the interval $[t(1-\epsilon), t]$. Then $P_{x_+^*(t(1-\epsilon))}(N(X, \epsilon t) = 0) = e^{-\kappa \epsilon t}$ and $Q(e^{\int_{t(1-\epsilon)}^t dB_{x_+^*(t(1-\epsilon))}(s)} \geq e^{-\epsilon \frac{1}{2}t}) \geq 1 - e^{-\epsilon ct}$. Thus, in the case, we can paste the piece of path from 0 to $x_+^*(t(1-\epsilon)) > x + \frac{\epsilon Mt}{2}$ on $[0, t(1-\epsilon)]$ to a constant path on $[t(1-\epsilon), t]$ and by setting $x_+^*(t) = x_+^*(t(1-\epsilon)) \in [x + \frac{\epsilon}{2}Mt, x + (1 - \frac{\epsilon}{2})Mt]$, we get

$$Q(E_x[e^{\int_0^t dB_{X(s)}(s)} 1_{[0, Mt]}(N(X, t)) 1_{x_+^*(t)}(X(t))] \geq e^{(\gamma_1(\kappa) - \frac{\epsilon}{99} - \epsilon \frac{1}{2}t)}) \geq 1 - \epsilon^{50} - e^{-\epsilon ct}$$

This gives the result if $x_+^*(t(1-\epsilon)) \in [x + \frac{\epsilon}{2}Mt, x + (1 - \frac{\epsilon}{2})Mt]$. On the other hand, if first $x < x_+^*(t(1-\epsilon)) < x + \frac{\epsilon}{2}Mt$, we consider paths in $\Gamma_{[t(1-\epsilon), t], \epsilon Mt}^{x_+^*(t(1-\epsilon))}$ for which all jumps are to the right. Call this set Γ_{right} . Then, for some $c > 0$ it follows that for all $\gamma \in \Gamma_{right}$

$$Q(e^{\int_{t(1-\epsilon)}^t dB_{\gamma(s)}(s)} \geq e^{-\epsilon \frac{1}{2}t}) \geq 1 - e^{-\epsilon ct}.$$

Also, for $K_1 = 2M \log \frac{2M}{\epsilon \kappa}$, $P_{x_+^*(t(1-\epsilon))}(X(\cdot - t(1-\epsilon)) \in \Gamma_{right}) \geq e^{-\epsilon K_1 t}$. Thus, using the Markov property and the above estimates, we have

$$Q(E_x[e^{\int_0^t dB_{X(s)}(s)} 1_{[0, Mt]}(N(X, t)) 1_{x_+^*(t(1-\epsilon)) + \epsilon Mt}(X(t))] \geq e^{(\gamma_1(\kappa) - \frac{\epsilon}{99} - \epsilon(\frac{1}{2} + K_1)t)}) \geq 1 - \epsilon^{50} - e^{-\epsilon ct}$$

which implies the existence of $x_+^* \in [x + \frac{\epsilon}{2}Mt, x + (1 - \frac{\epsilon}{2})Mt]$ and c_1 satisfying the lemma. When $x_+^*(t(1-\epsilon)) > x + (1 - \frac{\epsilon}{2})Mt$, we use an analogous argument with Γ_{left} . The measurability claim is an easy consequence of **Corollary 2.6** and the use of paths in Γ_{right} . \square

We now establish a block argument, a la percolation theory, starting with

Lemma 2.8. *Let $x_{\pm}^*(\epsilon, t, x)$ be the points given by **Lemma 2.7** when using the Brownian fields $\{B_y(s), y \in \mathbf{Z}, \epsilon^2 t \leq s \leq t\}$. Then,*

$$x_+^*(\epsilon, t, x) \in [x + \frac{\epsilon}{2}Mt(1 - \epsilon^2), x + (1 - \frac{\epsilon}{2})Mt(1 - \epsilon^2)] \subset [x + \frac{\epsilon}{4}Mt, x + (1 - \frac{\epsilon}{4})Mt]$$

and

$$x_-^*(\epsilon, t, x) \in [x - (1 - \frac{\epsilon}{2})Mt(1 - \epsilon^2), x - \frac{\epsilon}{2}Mt(1 - \epsilon^2)] \subset [x - (1 - \frac{\epsilon}{4})Mt, x - \frac{\epsilon}{4}Mt].$$

and

$$\begin{aligned} & Q(\forall x \in \{-Mt, (-M + \epsilon^2)t, (-M + 2\epsilon^2)t, \dots, (M - \epsilon^2)t, Mt\}, \\ & E_x[e^{\int_{\epsilon^2 t}^t dB_{X(s-\epsilon^2 t)}(s)} 1_{[0, Mt(1-\epsilon^2)]}(N(X, (1-\epsilon^2)t)) 1_{x_+^*(\epsilon, t, x)}(X(t))] \geq e^{(\gamma_1(\kappa) - c_1 \epsilon)t(1-\epsilon^2)}) \\ & \geq 1 - \epsilon^{46} \end{aligned}$$

and

$$\begin{aligned}
& Q(\forall x \in \{-Mt, (-M + \epsilon^2)t, (-M + 2\epsilon^2)t, \dots, (M - \epsilon^2)t, Mt\}, \\
& E_x[e^{\int_{\epsilon^2 t}^t dB_{X(s-\epsilon^2 t)}(s)} 1_{[0, Mt(1-\epsilon^2)]}(N(X, (1-\epsilon^2)t) 1_{x_-^*(\epsilon, t, x)}(X(t)))] \geq e^{(\gamma_1(\kappa) - c_1 \epsilon)t(1-\epsilon^2)} \\
& \geq 1 - \epsilon^{46}
\end{aligned}$$

provided ϵ is sufficiently small.

Proof. Since there are $\frac{2M}{\epsilon^2}$ points x in question, this follows easily from **Lemma 2.7** if ϵ is sufficiently small. \square

As the next step we establish the previous result for all $x \in \{-Mt, -Mt+1, \dots, Mt-1, Mt\}$.

Lemma 2.9. *For each $x \in [-Mt, Mt] \cap \mathbf{Z}$, there are points $x_{\pm}^*(t, x)$ lying in $[x + \frac{\epsilon}{4}Mt, x + (1 - \frac{\epsilon}{4})Mt]$, $[x - (1 - \frac{\epsilon}{4})Mt, x - \frac{\epsilon}{4}Mt]$, respectively, with $x_{\pm}^*(t, x)$ measurable with respect to $\sigma(B_y(s) : y \in \mathbf{Z}, |y - x| \leq Mt, s \leq t)$ such that for some $c = c(\kappa, M)$*

$$\begin{aligned}
& Q(\forall x \in \{-Mt, -Mt+1, \dots, Mt-1, Mt\}, \\
& E_x[e^{\int_0^t dB_{X(s)}(s)} 1_{[0, Mt]}(N(X, t)) 1_{x_{\pm}^*(t, x)}(X(t))] \geq e^{(\gamma_1(\kappa) - c\epsilon)t} \geq 1 - \epsilon^{46},
\end{aligned}$$

with the analogous statement holding for $x_{\pm}^*(t, x)$. Thus the probability of the intersection of the events for $x_{\pm}^*(t, x)$ and $x_{\mp}^*(t, x)$ is at least $1 - \epsilon^{45}$.

Proof. Start by observing that to each $x \in \{-Mt, -Mt+1, \dots, Mt-1, Mt\}$ we can associate a $y \in \{-Mt, (-M + \epsilon^2)t, (-M + 2\epsilon^2)t, \dots, (M - \epsilon^2)t, Mt\}$ such that $|x - y| \leq \frac{\epsilon^2}{2}t$ and $P_x(N(X, \epsilon^2 t) = |x - y|, X(\epsilon^2 t) = y) \geq e^{-K_3 \epsilon^2 t}$, with K_3 independent of ϵ and t . Also, for some K_4, K_5 independent of ϵ and t we have by **Theorem 1.1**

$$Q(\inf_{\gamma \in \Gamma_{[0, \epsilon^2 t], |x-y|: \gamma(\epsilon^2 t) = x}^x} \int_0^{\epsilon^2 t} dB_{\gamma(s)}(s) \geq -K_4 \epsilon^2 t) \geq 1 - e^{-K_5 \epsilon^2 t}.$$

Put $x_{\pm}^*(t, x) = x_{\pm}^*(\epsilon, t, y)$ and $x_{\mp}^*(t, x) = x_{\mp}^*(\epsilon, t, y)$ where the latter are given by **Lemma 2.8**. Then, piecing together paths on $[0, \epsilon^2 t]$ which connect x to y and on $[\epsilon^2 t, Mt]$ linking y to $x_{\pm}^*(\epsilon, t, y)$ and using the fact that there are $2Mt+1$ points in $\{-Mt, -Mt+1, \dots, Mt-1, Mt\}$, we have the result. \square

We next iterate so we can set up a percolation scheme.

Lemma 2.10. *For $i \in \{0, 1, \dots, (\frac{8}{\epsilon} - 1)\}$, and $x \in [-Mt + \frac{i\epsilon}{4}Mt, Mt + \frac{i\epsilon}{4}Mt]$, $x' \in [-Mt - \frac{i\epsilon}{4}Mt, Mt - \frac{i\epsilon}{4}Mt]$ there exist $x_{\pm}^*(i, x) \in [-Mt + \frac{(i+1)\epsilon}{4}Mt, Mt + \frac{(i+1)\epsilon}{4}Mt]$, $x_{\pm}^*(i, x') \in [-Mt -$*

$\frac{(i+1)\epsilon}{4}Mt, Mt - \frac{(i+1)\epsilon}{4}Mt]$, measurable with respect to the Brownian fields $\{B_y(s+it) : 0 \leq s \leq t\}$, such that the events

$$A(i) = \{\forall x \in [-Mt + \frac{i\epsilon t}{4}Mt, Mt + \frac{i\epsilon t}{4}Mt], \\ E_x[e^{\int_0^t dB_{X(s)}(it+s)} \mathbf{1}_{x_+^*(i,x)}(X(t)) \mathbf{1}_{[0,Mt]}(N(X,t))] \geq e^{(\gamma_1(\kappa) - \epsilon)t}\}$$

and

$$B(i) = \{\forall x \in [-Mt - \frac{i\epsilon t}{4}Mt, Mt - \frac{i\epsilon t}{4}Mt], \\ E_x[e^{\int_0^t dB_{X(s)}(it+s)} \mathbf{1}_{x_-^*(i,x)}(X(t)) \mathbf{1}_{[0,Mt]}(N(X,t))] \geq e^{(\gamma_1(\kappa) - \epsilon)t}\}$$

satisfy, for ϵ sufficiently small (depending only on M)

$$Q(\cap_{i=0}^{\frac{8}{\epsilon}-1} A(i)) \geq 1 - \epsilon^{44}$$

and

$$Q(\cap_{i=0}^{\frac{8}{\epsilon}-1} B(i)) \geq 1 - \epsilon^{44}.$$

Proof. We can apply **Lemma 2.9**. If $x \in [-Mt + \frac{i\epsilon}{4}Mt, Mt + \frac{i\epsilon}{4}Mt]$, we consider the two cases $x \in [-Mt + \frac{i\epsilon}{4}Mt, \frac{\epsilon}{4}Mt + \frac{i\epsilon}{4}Mt]$ and $x \in (\frac{\epsilon}{4}Mt + \frac{i\epsilon}{4}Mt, Mt + \frac{i\epsilon}{4}Mt]$. In the first case, take $x_+^*(i, x) = x_+^*(t, x)$ from **Lemma 2.9**, using the Brownian fields $\{B_y(s+ti) : 0 \leq s \leq t, |y-x| \leq Mt\}$. Then we'll have

$$x_+^*(i, x) \in [-Mt + \frac{(i+1)\epsilon}{4}Mt, \frac{\epsilon}{4}Mt + \frac{i\epsilon}{4}Mt + (1 - \frac{\epsilon}{4})Mt] \subset [-Mt + \frac{(i+1)\epsilon}{4}Mt, Mt + \frac{(i+1)\epsilon}{4}Mt].$$

In the second case, take $x_+^*(i, x) = x_-^*(t, x)$ from **Lemma 2.9**, using the Brownian fields $\{B_y(s+ti) : 0 \leq s \leq t, |y-x| \leq Mt\}$. Then we'll have

$$x_+^*(i, x) \in [x - (1 - \frac{\epsilon}{4})Mt, x - \frac{\epsilon}{4}Mt] \subset [-Mt + \frac{(i+1)\epsilon}{4}Mt, Mt + \frac{(i+1)\epsilon}{4}Mt].$$

An analogous argument handles the selection of the points $x_-^*(i, x')$. Notice that $1 - \frac{8M}{\epsilon}\epsilon^{46} \geq 1 - \epsilon^{44}$ for ϵ sufficiently small (depending only on M) to get the probability estimates. \square

Proof. We now prove **Theorem 2.2**, namely that $\gamma(\kappa) = \lim_{t \rightarrow \infty} \frac{1}{t} \log E_x[e^{\int_0^t dB_{X(s)}(s)}]$. It is clear that we need only do this for $x = 0$. Since

$$\gamma(\kappa) = \lim_{t \rightarrow \infty} \frac{1}{t} \log v(t, 0) \leq \lim_{t \rightarrow \infty} \frac{1}{t} \log E_0[e^{\int_0^t dB_{X(s)}(s)}]$$

and both limits exist Q a.s. we only need show that

$$Q(\lim_{t \rightarrow \infty} \frac{1}{t} \log v(t, 0) \geq \lim_{t \rightarrow \infty} \frac{1}{t} \log E_0[e^{\int_0^t dB_{X(s)}(s)}]) \geq \frac{1}{4}.$$

Using **Lemma 2.10**, we can conclude that with Q -probability at least $1 - \epsilon^{43}$, corresponding to the starting point 0 there exist sequences, $\{x_+^*(i, 0)\}_{i=0}^{\frac{8}{\epsilon}t}$, $\{x_-^*(i, 0)\}_{i=0}^{\frac{8}{\epsilon}t}$, such that $x_{\pm}(0, 0) = 0$ with $x_+^*(i, 0) \in [-Mt + \frac{i\epsilon}{4}Mt, Mt + \frac{i\epsilon}{4}Mt]$, $x_-^*(i, 0) \in [-Mt - \frac{i\epsilon}{4}Mt, Mt - \frac{i\epsilon}{4}Mt]$, and for $0 \leq i \leq \frac{8}{\epsilon}t - 1$,

$$E_{x_+^*(i,0)}[e^{\int_0^t dB_{X(s)}(s+it)} 1_{x_+^*(i+1,0)}(X(t)) 1_{[0,Mt]}(N(X, t))] \geq e^{(\gamma_1(\kappa) - c\epsilon)t}$$

and

$$E_{x_-^*(i,0)}[e^{\int_0^t dB_{X(s)}(s+it)} 1_{x_-^*(i+1,0)}(X(t)) 1_{[0,Mt]}(N(X, t))] \geq e^{(\gamma_1(\kappa) - c\epsilon)t}$$

Thus, by the Markov property

$$E_0[e^{\int_0^{\frac{8}{\epsilon}t} dB_{X(s)}(s)} 1_{x_{\pm}(\frac{8}{\epsilon}t)}(X(\frac{8}{\epsilon}t)) 1_{[0,Mt]}(N(X, t))] \geq e^{(\gamma_1(\kappa) - c\epsilon)\frac{8t}{\epsilon}}.$$

This induces the following percolation system: $\{\psi(l, n), n \geq 0, l \in \mathbf{Z}, l + n \equiv 0 \pmod{2}\}$. We say that $(l, n) \rightarrow (l + 1, n + 1)$ if $\forall x \in [(2l - 1)Mt, (2l + 1)Mt]$, $\exists x_0, \dots, x_{\frac{8}{\epsilon}t - 1}$, with $x_0 = x$, such that $x_i \in [(2l - 1)Mt + \frac{i\epsilon}{4}Mt, (2l + 1)Mt + \frac{i\epsilon}{4}Mt]$, and

$$E_{x_i}[e^{\int_0^t dB_{X(s)}(\frac{8nt}{\epsilon} + it + s)} 1_{[0,Mt]}(N(X(t))) 1_{x_{i+1}}(X(t))] \geq e^{(\gamma_1(\kappa) - c\epsilon)t}.$$

And we make an analogous definition for $(l, n) \rightarrow (l - 1, n + 1)$. By the measurability claim in **Lemma 2.10**, this is a 2-dependent percolation scheme with Q -probability of an open bond at least $1 - \epsilon^{43}$. Thus, by Durrett [D93] we have for any n , with Q -probability at least $\frac{1}{2}$ that there is a path $(0, 0) \rightarrow (l_1, 1) \rightarrow \dots \rightarrow (0, 2n)$. Thus, for some $x^* \in [-Mt, Mt]$,

$$E_0[e^{\int_0^{\frac{8}{\epsilon}t2n} dB_{X(s)}(s)} 1_{x^*}(X(\frac{8}{\epsilon}t2n))] \geq e^{(\gamma_1(\kappa) - c\epsilon)\frac{8}{\epsilon}t2n}.$$

But, for some $h > 0$, independent of t, n, ϵ, x^* with Q -probability at least $\frac{1}{2}$,

$$E_{x^*}[e^{\int_0^{\frac{16t}{\epsilon}} dB_{X(s)}(s + \frac{8}{\epsilon}t2n)} 1_0(X(\frac{16t}{\epsilon}))] \geq e^{-h\frac{16t}{\epsilon}}.$$

Thus, with Q -probability at least $\frac{1}{4}$ we have

$$E_0[e^{\int_0^{\frac{8}{\epsilon}t2(n+1)} dB_{X(s)}(s)} 1_0(X(\frac{8}{\epsilon}t2(n+1)))] \geq e^{(\gamma_1(\kappa) - 2c\epsilon)\frac{8}{\epsilon}t2(n+1)}.$$

This proves the theorem. □

The next result will lead to the existence of $\lim_{t \rightarrow \infty} u(t, x)$.

Theorem 2.11. *Given $\epsilon > 0$ there is a positive constant c such that*

$$Q(E_0[e^{\int_0^t dB_{X(s)}(s)}] \geq e^{(\gamma_1(\kappa) + \epsilon)t}) \leq e^{-ct}.$$

Before proving this Theorem we derive the following consequence,

Corollary 2.12.

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log E_0[e^{\int_0^t dB_{X(s)}(t-s)}] \leq \gamma_1(\kappa).$$

Proof. Since the distribution of $E_0[e^{\int_0^t dB_{X(s)}(t-s)}]$ and $E_0[e^{\int_0^t dB_{X(s)}(s)}]$ are identical, we can apply **Theorem 2.11** and Borel-Cantelli to get the result. \square

Corollary 2.13. *For u_0 positive, strictly positive somewhere and bounded,*

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log u(t, x) = \gamma(\kappa).$$

Proof. Without loss of generality we take u_0 bounded by 1 and strictly positive at the origin. By Carmona and Molchanov,[?] $\liminf_{t \rightarrow \infty} \frac{1}{t} \log u(t, x) \geq \gamma(\kappa)$. By **Corollary 2.12** however $\limsup_{t \rightarrow \infty} \frac{1}{t} \log u(t, x) \leq \limsup_{t \rightarrow \infty} \frac{1}{t} \log E[e^{\int_0^t dB_{X(t-s)}(s)}] = \gamma_1(\kappa) = \gamma(\kappa)$. \square

We can now state our main result.

Theorem 2.14.

$$\lim_{\kappa \downarrow 0} \gamma(\kappa) \log \frac{1}{\kappa} = \frac{\alpha^2}{4}.$$

The proof of **Theorem 2.14** will be given in the next section. Some of the results needed for the proof of **Theorem 2.14** will be used in the proof of **Theorem 2.11**, which is given in the final section.

3 Asymptotics of the Lyapunov exponent

The proof of **Theorem 2.14** is based on Borell's inequality and Fernique-Talagrand's theorem for Gaussian random fields together with a classical large deviation result for Poisson processes.

Recall from **Theorem 1.1**, that $\alpha > 0$ satisfies $\alpha = \sup_n \frac{1}{n} E_Q[\vec{A}_{[0,n],n}] \leq C$.

Lemma 3.1. *Let any $a > 0$ be fixed. For every $\epsilon > 0$*

$$Q\left(\exists n_0(\omega) : (\alpha - \epsilon)\sqrt{mn} \leq \vec{A}_{[0,n],m} \leq (\alpha + \epsilon)\sqrt{nm} \quad (\forall m \geq an), \quad \forall n \geq n_0\right) = 1.$$

Proof. Applying **Theorem 3.1**, we have

$$Q(|\vec{A}_{[0,n],n}^0 - E_Q[\vec{A}_{[0,n],n}^0]| > \frac{\epsilon n}{2}) \leq 2e^{-\epsilon^2 n/8}.$$

Since $E_Q[\vec{A}_{[0,n],n}^0] \sim \alpha n$ by Liggett [Lig85], it holds that for large n

$$Q(|\vec{A}_{[0,n],n}^0 - \alpha n| > \epsilon n) \leq 2e^{-\epsilon^2 n/8},$$

which becomes

$$Q(|\vec{A}_{[0,n],m}^0 - \alpha\sqrt{nm}| > \epsilon\sqrt{nm}) \leq 2e^{-\epsilon^2 m/8},$$

by the Brownian scaling invariance;

$$\vec{A}_{[0,n],m} \stackrel{(d)}{=} \sqrt{\frac{n}{m}} \vec{A}_{[0,m],m}.$$

Finally use the Borel-Cantelli lemma to complete the proof of **Lemma 3.1**. □

Lemma 3.2. (i) Let $F : [0, \infty) \mapsto \mathbb{R}$ be a continuous function satisfying

$$\limsup_{x \rightarrow \infty} \frac{F(x)}{x \log x} < 1.$$

Then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log E_0[e^{nF(\frac{N(X,n)}{n})}] = \sup_{x \geq 0} \{F(x) - x \log \frac{x}{\kappa} + x - \kappa\}.$$

(ii) For $0 \leq a < b \leq \infty$

$$\begin{aligned} & \lim_{n \rightarrow \infty} \frac{1}{n} \log E_0[e^{(\alpha\sqrt{nN(X,n)} - \beta N(X,n))} : an \leq N(X,n) \leq bn] \\ &= \sup_{a \leq x \leq b} \{\alpha\sqrt{x} - \beta x - x \log \frac{x}{\kappa} + x - \kappa\} \end{aligned}$$

Proof. Since $N(X,n)$ has a Poisson distribution with mean κn , it follows from a large deviation theorem by Cramer that for any bounded continuous function $F : [0, \infty) \mapsto \mathbb{R}$

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log E_0[e^{nF(\frac{N(X,n)}{n})}] = \sup_{x \geq 0} \{F(x) - x \log \frac{x}{\kappa} + x - \kappa\}.$$

Furthermore it is easy to extend it for any F satisfying the assumption. (ii) follows from (i) immediately. □

Lemma 3.3. (i) For $\theta > 0$

$$\sup_{x \geq 0} \left\{ \theta \sqrt{x} - x \log \frac{x}{\kappa} + x - \kappa \right\} \sim \frac{\theta^2}{4 \log \frac{1}{\kappa}} \quad (\kappa \downarrow 0).$$

(ii) For $c > 0$, $\theta > 0$, $0 < q < 1$ and $\kappa > 0$,

$$\sup_{\kappa^q \leq x \leq b} \left(\theta \sqrt{x} + (\log c \kappa^q) x - x \log \frac{x}{\kappa} + x - \kappa \right) \sim \frac{\theta^2}{4(1+q)} \frac{1}{\log \frac{1}{\kappa}} \quad (\kappa \downarrow 0).$$

Proof. For (ii) let

$$f(x) = \theta \sqrt{x} + (\log c \kappa^q) x - x \log \frac{x}{\kappa} + x - \kappa.$$

$$f'(x) = \frac{\theta}{2\sqrt{x}} + (\log c \kappa^q) - \log \frac{x}{\kappa}$$

is a decreasing function and $f'(\kappa^q) > 0$ and $f'(b) < 0$ for a sufficiently small $\kappa > 0$, so that f has a unique maximum at $x(\kappa) \in (\kappa^q, b)$ such that

$$\frac{\theta}{2\sqrt{x(\kappa)}} + (\log c \kappa^q) - \log \frac{x(\kappa)}{\kappa} = 0,$$

which implies

$$\frac{\theta}{2\sqrt{x(\kappa)}} \sim (1+q) \log \frac{1}{\kappa} \quad (\kappa \downarrow 0).$$

Hence

$$\sup_{\kappa^q \leq x \leq b} f(x) = f(x(\kappa)) \sim \frac{\theta^2}{4(1+q)} \frac{1}{\log \frac{1}{\kappa}}.$$

□

(Proof of the upper bound)

By **Lemma 3.1**, we can suppose that for $a > 0$ and $\epsilon > 0$

$$\vec{A}_{[0,n],m} \leq (\alpha + \epsilon) \sqrt{mn} \quad (\forall m \geq an) \quad Q - a.s..$$

Then

$$\begin{aligned} E_0[e^{\int_0^n dB_{X(s)}(s)} : N(X, n) \geq an] &\leq \sum_{j=an}^{\infty} e^{\vec{A}_{[0,n],j}} P_0(N(X, n) = j) \\ &\leq \sum_{j=an}^{\infty} e^{(\alpha+\epsilon)\sqrt{jn}} \mathbb{P}(N(X, n) = j) \\ &\leq E_0[e^{(\alpha+\epsilon)\sqrt{nN(X,n)}}], \end{aligned} \tag{3.1}$$

Noting that $\vec{A}_{[0,n],m}$ is increasing in m we have

$$E_0[e^{\int_0^n dB_{X(s)}(s)} : N(X, n) \leq an] \leq e^{\vec{A}_{[0,n],an}} \leq e^{(\alpha+\epsilon)\sqrt{an}}.$$

Combining ([Sh97]), ([ShF98]) together with **Lemma 3.2**, we have

$$\gamma(\kappa) \leq \sup\{(\alpha + \epsilon)\sqrt{x} - x \log \frac{x}{\kappa} - x + \kappa\} + (\alpha + \epsilon)\sqrt{a}.$$

Choosing a suitable $a = a(\kappa)$ using by **Lemma 3.3(i)** we obtain

$$\limsup_{\kappa \searrow 0} \frac{\gamma(\kappa)}{\log \frac{1}{\kappa}} \leq \frac{(\alpha + \epsilon)^2}{4},$$

which gives the upper bound of **Theorem 2.14**.

(Proof of the lower bound.)

For $\gamma \in \Gamma_{[0,n],m}^\epsilon$ let $t_1 < t_2 < \dots < t_m < n$ the jump times of γ during $[0, n]$, and let I_i^γ be the interval $[\frac{j_i \epsilon}{3}, \frac{(j_i+1)\epsilon}{3}]$ with an integer j_i , including t_i . If $\eta \in \Gamma_{[0,n],m}^\epsilon$ has $s_1 < s_2 < \dots < s_m < n$ as its jump times during $[0, n]$, $I_i^\eta = I_i^\gamma$ and $\gamma(t_i) = \eta(s_i)$ for every $1 \leq i \leq m$, we denote $\eta \sim \gamma$. Then it is easy to see that if $\eta, \gamma \in \Gamma_{[0,n],m}^\epsilon$ satisfy $\eta \sim \gamma$,

$$|\int_0^n dB_{\gamma(s)}(s) - \int_0^n dB_{\eta(s)}(s)| < \omega(\gamma),$$

where

$$\omega(\gamma) = \sum_{i=1}^m (\omega(B_{\gamma(t_i)} : I_i) + \omega(B_{\gamma(t_{i-1})} : I_i)),$$

$$\omega(B_x : I) = \sup_{t,s \in I} |B_x(t) - B_x(s)|.$$

Lemma 3.4. For any $\delta > 0$, $a > 0$ and $b > a$ there exists $\epsilon_0 > 0$ such that for all $0 < \epsilon < \epsilon_0$ and for all $an \leq m \leq bn$, (i) there is a $K > 0$ such that

$$Q(\omega(\gamma) > cn) \leq K^m e^{-\frac{cn}{\sqrt{\epsilon}}} \text{ for all } \gamma \in \Gamma_{[0,n],m}^\epsilon.$$

(ii)

$$Q\left(\sup_{\gamma \in \Gamma_{[0,n],m}^\epsilon} \omega(\gamma) \geq \delta \sqrt{nm}\right) \leq ce^{-n}.$$

(iii) Moreover,

$$Q \left(\exists n_0(\omega) : \sup_{\gamma \in \Gamma_{[0,n],m}^\epsilon} \omega(\gamma) \leq \delta \sqrt{nm} \quad , \forall m \in [an, bn], \quad \forall n \geq n_0 \right) = 1$$

Proof. We first count the number of increasing sequences $j_1 < j_2 < \dots < j_m$ of integers which determine the intervals I_i^ϵ . We must have $j_m < \frac{3n}{2\epsilon}$. Thus, there are fewer than

$$\begin{aligned} \binom{\frac{3n}{2\epsilon}}{m} &\leq c_1 \frac{\left(\frac{3n}{2\epsilon}\right)^{\frac{3n}{2\epsilon} + \frac{1}{2}}}{m^{m+\frac{1}{2}} \left(n\left(\frac{3}{2\epsilon} - \frac{m}{n}\right)\right)^{n\left(\frac{3}{\epsilon}-1\right) + \frac{1}{2}}} \\ &\leq c_1 \left(\frac{c_2}{\epsilon}\right)^m, \end{aligned}$$

where c_1, c_2 are positive constants. For each such sequence, $j_1 < j_2 < \dots < j_m$, there are $(2)^n$ possible sequences of sites visited. This means there are $C_1 \left(\frac{2c_2}{\epsilon}\right)^m$ possible variables $2(\omega(B_{\gamma(t_i)}, I_i^\epsilon) + \omega(B_{\gamma(t_{i-1})}, I_{i-1}^\epsilon))$ to consider. For each such sequence,

$$\omega(\gamma) \stackrel{L}{=} \sqrt{\epsilon} \sum_{i=1}^m Y_i$$

where Y_i are *i.i.d.* random variables satisfying

$$E_Q[e^{Y_i}] = K < \infty.$$

Then for fixed $\gamma \in \Gamma_{[0,n],m}^{0,\epsilon}$,

$$Q(\omega(\gamma) > cn) \leq K^m e^{-\frac{cn}{\sqrt{\epsilon}}},$$

which proves (i). Therefore,

$$\begin{aligned} Q \left(\sup_{\gamma \in \Gamma_{[0,n],n}^{0,\epsilon}} \omega(\gamma) > c\sqrt{nm} \right) &\leq c_1 \left(\frac{c_2}{\epsilon}\right)^m K^m e^{-\frac{c\sqrt{nm}}{\sqrt{\epsilon}}} \\ &\leq c_1 e^{-cn}, \end{aligned}$$

provided ϵ is sufficiently small. □

Lemma 3.5. For any delta > 0 , and all $a > 0$ and $b > a$ there exists $\epsilon_0 > 0$ such that for $0 < \epsilon < \epsilon_0$

$$Q \left(\exists n_0(\omega) : |A_{[0,n],m}^\epsilon - \alpha \sqrt{nm}| \leq \delta \sqrt{nm} \quad , \forall m \in [an, bn], \quad \forall n \geq n_0 \right) = 1.$$

Proof. By Borell's inequality,

$$Q \left(|\vec{A}_{[0,n],m}^\epsilon - E_Q[\vec{A}_{[0,n],m}^\epsilon]| \geq \frac{\delta\sqrt{nm}}{2} \right) \leq 2e^{-\delta^2 m/8}.$$

Since

$$E_Q[\vec{A}_{[0,n],m}^\epsilon] = \sqrt{\frac{n}{m}} E_Q[\vec{A}_{[0,m],m}^\epsilon] \sim \sqrt{nm}\alpha$$

uniformly in $m \in [an, bn]$ as $n \rightarrow \infty$, we have

$$Q \left(|\vec{A}_{[0,n],m}^\epsilon - \alpha\sqrt{nm}| \geq \delta\sqrt{nm} \right) \leq 2e^{-\delta^2 m/8}.$$

□

By **Lemmas 3.6** and **3.7** we can assume that the Brownian motions $\{B_x(t)\}$ satisfy the following two conditions:

For every $\delta > 0$, $a > 0$ and $b > a$, there exists $\epsilon_0 > 0$ such that if $0 < \epsilon < \epsilon_0$,

$$\exists n_0 \text{ such that } \sup_{\gamma \in \Gamma_{[0,n],m}^\epsilon} \omega(\gamma) \leq \delta\sqrt{nm} \text{ for all } m \in [an, bn], \text{ for all } n \geq n_0,$$

and

$$\exists n_0 \text{ such that } |\vec{A}_{[0,n],m}^\epsilon - \alpha\sqrt{nm}| \leq \delta\sqrt{nm} \text{ for all } m \in [an, bn], \text{ for all } n \geq n_0.$$

Then,

$$\begin{aligned} & E_0[e^{\int_0^n dB_{X(s)}(s)}] \\ & \geq \sum_{m=an}^{bn} E_0[e^{\int_0^n dB_{X(s)}(s)} : N(X, n) = m] \\ & = \sum_{m=an}^{bn} e^{A_{[0,n],m}^\epsilon} E_0[e^{(\int_0^n dB_{X(s)}(s) - \int_0^n dB_{\gamma^{n,m}(s)}(s))} : N(X, n) = m] \\ & \equiv (*) \end{aligned}$$

where $\gamma^{n,m} \in \Gamma_{[0,n],m}^\epsilon$ is defined by

$$\vec{A}_{[0,n],m}^\epsilon = \int_0^n dB_{\gamma^{n,m}(s)}(s).$$

Thus,

$$\begin{aligned}
(*) &\geq \sum_{m=an}^{bn} e^{(\tilde{A}_{[0,n],m}^\epsilon - \omega(\gamma^{n,m}))} P_0(X \sim \gamma^{n,m}, N(X, n) = m) \\
&\geq \sum_{m=an}^{bn} e^{(\alpha - 2\delta)\sqrt{nm}} P_0(X \sim \gamma^{n,m}, N(X, n) = m)
\end{aligned}$$

Since

$$P_0(X \sim \gamma^{n,m} | N(X, n) = m) = \frac{m!}{n^m} \left(\frac{\epsilon}{6}\right)^m,$$

supposing $a = \kappa^q$ with $0 < q < 1$, we have some $c > 0$ satisfying

$$P_0(X \sim \gamma^{n,m} | N(X, n) = m) \geq (c\epsilon\kappa^q)^m,$$

so that

$$\begin{aligned}
&E_0[e^{\int_0^n dB_{X(s)}(s)}] \\
&\geq E_0[e^{((\alpha - 2\delta)\sqrt{nN(X,n)} + \log(c\epsilon\kappa^q)N(X,n))}] \\
&\quad \kappa^q n \leq N(X, n) \leq bn].
\end{aligned}$$

Hence by **Lemma 3.3(ii)**, we obtain that for every $\delta > 0$ and $0 < q < 1$

$$\liminf_{\kappa \downarrow 0} \gamma(\kappa) \log \frac{1}{\kappa} \geq \frac{(\alpha - 2\delta)^2}{4(1+q)},$$

which gives the lower bound of the **Theorem 2.14**, letting $q \searrow 0$ and $\kappa \searrow 0$.

4 Proof of Theorem 2.11

Before making the next definition we remark that by **Lemma 3.4** with $a = M, b = \infty, \alpha$ replaced by $\alpha + \epsilon, \beta = 0$ then if M is sufficiently large,

$$\begin{aligned}
E_x[e^{\int_0^{N_0} dB_{X(s)}(s)} 1_{[MN_0, \infty)}] &\leq \sum_{m=MN_0}^{\infty} e^{\tilde{A}_{[0, N_0], m}^\epsilon} P(N(X, N_0) = m) \\
&\leq \sum_{m=MN_0}^{\infty} e^{(\alpha + \epsilon)\sqrt{mN_0}} P(N(X, N_0) = m) \\
&= E_x[E^{(\alpha + \epsilon)\sqrt{mN_0}} : MN_0 \leq N(X, N_0)] \\
&\leq e^{(1+\delta)N_0((\alpha + \epsilon)\sqrt{M} - M \log \frac{M}{\kappa} + M - \kappa)} \\
&\leq e^{-cMN_0}, \text{ for some positive } c.
\end{aligned}$$

For the remainder of the section, we assume that the value of M is large enough to make the above estimate valid. This means we no longer need to worry about the random walk paths that make too many jumps (i. e. more than MN_0 in time N_0 .) their contribution to the expectation is negligible. We introduce a notation for the time-shifted Brownian motion field by setting

$$B_x^j(t) = B_x(t + j).$$

Definition 4.1. Break up space-time into disjoint blocks of the form $((i - 1)MN_0, (i + 1)MN_0] \times [(j - 1)N_0, jN_0]$ where $i \in \mathbf{Z}, j \in \mathbf{Z}^+$. satisfy $i + j \equiv 0 \pmod{2}$. Label the block $((i - 1)MN_0, (i + 1)MN_0] \times [(j - 1)N_0, jN_0]$ by (i, j) . We say (i, j) is a good block (denoted $(i, j) \in G$) if $\forall x \in ((i - 1)MN_0, (i + 1)MN_0]$,

$$E_x[e^{\int_0^{N_0} dB_{X(s)}^j(s)} I_{[0, MN_0]}(N(X, N_0))] \leq e^{(\gamma(\kappa) + \frac{\epsilon}{3})N_0}.$$

Next we introduce a discretization scheme for paths in $\Gamma_{[0, t], Mt}^0$ which will allow us via simple counting arguments to derive some estimates on our basic functionals of $\{B_x\}$. For a path $\gamma \in \Gamma_{[0, t], Mt}^0$ we say that its N_0 -skeleton (or skeleton since the value N_0 will be 'understood') is the sequence $\gamma(0), \gamma(N_0), \gamma(2N_0), \dots, \gamma(t)$, where we assume without loss of generality that t is a multiple of N_0 . We claim

Lemma 4.1. *Given $\epsilon > 0$ there is an N so that for $N_0 > N$, the number of distinct skeletons corresponding to paths in $\Gamma_{[0, t], Mt}^0$ is bounded by $e^{\frac{\epsilon t}{10^6}}$.*

Proof. Each skeleton determines a sequence $\{i_j : j = 0, \dots, \frac{t}{N_0}\}$ with $i_j + j \equiv 0 \pmod{2}$ such that $\gamma(jN_0) \in [(i_j - 1)MN_0, (i_j + 1)MN_0]$ for each j . We shall call the sequence $\{i_j : j = 0, \dots, \frac{t}{N_0}\}$ the trace of γ . Letting $A(i_1, i_2, \dots, i_{\frac{t}{N_0}})$ denote the set of skeletons of elements of $\Gamma_{[0, t], Mt}^0$ with trace $\{i_j : j = 0, \dots, \frac{t}{N_0}\}$, it's easy to see that if N_0 is sufficiently large,

$$|A(i_1, i_2, \dots, i_{\frac{t}{N_0}})| \leq (2MN_0)^{\frac{t}{N_0}}$$

Furthermore the number of traces of elements of $\Gamma_{[0, t], Mt}^0$ is constrained by the requirement that the corresponding path have no more than Mt jumps. In terms of the trace, this translates into the bound

$$\sum_j (|i_j - i_{(j-1)}| - 1)_+ \leq \frac{t}{N_0}.$$

There are at most $2^{3\frac{t}{N_0}}$ such sequences. Thus, the total number of skeletons of elements of $\Gamma_{[0, t], Mt}^0$ is bounded by

$$(2MN_0)^{\frac{t}{N_0}} 2^{3\frac{t}{10^6}} \leq e^{\frac{\epsilon t}{N_0}},$$

provided N_0 is chosen sufficiently large. □

Lemma 4.2. For any $\delta > 0$, and $M < \infty$ as above, there is an N_1 so that for $N_0 \geq N_1$, we have

$$Q((i, j) \in G) \geq 1 - \delta.$$

Proof. Without loss of generality we may treat the case $j = 1$. Fix an η with $0 < \eta \ll \frac{\epsilon}{M}$. Then, $\forall x \in ((i-1)MN_0, (i+1)MN_0] \cap \mathbf{Z}$,

$$E_x[e^{\int_0^{N_0} dB_{X(s)}(s)} \mathbf{1}_{[0, MN_0]}(N(X, N_0))] \leq \sum_{l=(i-2)MN_0}^{(i+2)MN_0} E_x[e^{\int_0^{\eta N_0} dB_{X(s)}(s)}] E_l[e^{\int_0^{(1-\eta)N_0} dB_{X(s)}^{\eta N_0}(s)}]$$

An easy argument using Chebychev's inequality and the fact that

$$E_Q E_x[e^{\int_0^{\eta N_0} dB_{X(s)}(s)}] = E_x E_Q[e^{\int_0^{\eta N_0} dB_{X(s)}(s)}] = e^{\frac{\eta N_0}{2}},$$

shows that the event

$$C(N_0) = \cap_{x=(i-1)MN_0}^{(i+1)MN_0} \{E_x[e^{\int_0^{\eta N_0} dB_{X(s)}(s)}] \leq \frac{e^{\frac{\epsilon N_0}{12}}}{4MN_0}\}$$

has $\lim_{N_0 \rightarrow \infty} Q(C(N_0)) = 1$. Moreover, on the event $C(N_0)$, we have

$$\begin{aligned} & \sup_{x \in [(i-1)MN_0, (i+1)MN_0]} E_x[e^{\int_0^{N_0} dB_{X(s)}(s)} \mathbf{1}_{[0, MN_0]}(N(X, N_0))] \\ & \leq e^{\frac{\epsilon N_0}{12}} \sup_{l \in [(i-2)MN_0, (i+2)MN_0]} E_l[e^{\int_0^{(1-\eta)N_0} dB_{X(s)}^{\eta N_0}(s)}]. \end{aligned}$$

The right hand side in this inequality can be bounded by a supremum over a finite number of terms, where that finite number is independent of N_0 . To do this, take points x_1, x_2, \dots, x_R in $[(i-2)MN_0, (i+2)MN_0]$, separated by $\frac{\eta \epsilon N_0}{2}$ so that $R \cong \frac{8M}{\eta \epsilon}$. For a fixed x_i and any l with $|x_i - l| \leq \frac{\eta \epsilon N_0}{2}$,

$$E_{x_i}[e^{\int_0^{N_0} dB_{X(s)}(s)}] \geq E_{x_i}[e^{\int_0^{\eta N_0} dB_{X(s)}(s)} \mathbf{1}_l(X(\eta N_0))] E_l[e^{\int_0^{(1-\eta)N_0} dB_{X(s)}^{\eta N_0}(s)}].$$

Now we claim

$$\lim_{N_0 \rightarrow \infty} Q\left(\inf_{|l-x_i| \leq \frac{\eta \epsilon N_0}{2}} E_{x_i}[e^{\int_0^{\eta N_0} dB_{X(s)}(s)} \mathbf{1}_l(X(\eta N_0))] \geq e^{-\frac{\epsilon N_0}{12}}\right) = 1.$$

Before proving this claim we proceed to show how it proves the lemma. Assuming the claim,

$$\lim_{N_0 \rightarrow \infty} Q\left(\sup_{1 \leq i \leq R} E_{x_i}[e^{\int_0^{N_0} dB_{X(s)}(s)}] \geq e^{-\frac{\epsilon N_0}{12}} \sup_{l \in [(i-2)MN_0, (i+2)MN_0]} E_l[e^{\int_0^{(1-\eta)N_0} dB_{X(s)}^{\eta N_0}(s)}]\right) = 1.$$

Thus, with Q probability approaching 1 as N_0 tends to ∞ we have for all $x \in [(i-1)MN_0, (i+1)MN_0]$,

$$\begin{aligned}
E_x[e^{\int_0^{N_0} dB_{X(s)}(s)} \mathbf{1}_{[0, MN_0]}(N(X, N_0))] &\leq e^{\frac{\epsilon N_0}{12}} \sup_{l \in [(i-2)MN_0, (i+2)MN_0]} E_l[e^{\int_0^{(1-\eta)N_0} dB_{X(s)}^{\eta N_0}(s)}] \\
&\leq e^{\frac{\epsilon N_0}{6}} \sup_{1 \leq i \leq R} E_{x_i}[e^{\int_0^{N_0} dB_{X(s)}(s)}] \\
&\leq e^{\frac{\epsilon N_0}{6}} e^{(\gamma(\kappa) + \frac{\epsilon}{6})N_0}, \text{ R independent of } N_0, \\
&= e^{(\gamma(\kappa) + \frac{\epsilon}{3})N_0}, \text{ as desired.}
\end{aligned}$$

Now back to the proof of the claim. First, given l with $|x_i - l| \leq \frac{\eta \epsilon N_0}{2}$, take an $x_i - to - l$ path γ taking exactly $|x_i - l|$ steps and evenly spaced jump times. Using the notation of the previous section (see material preceding **Lemma 3.6**.) then making sure ϵ is taken so that $\kappa \epsilon < 1$, we have

$$\begin{aligned}
P_{x_i}(X \sim \gamma) &= 2^{-|x_i - l|} \frac{(\lambda \eta N_0)^{|x_i - l|}}{|x_i - l|!} e^{-\lambda \eta N_0 |x_i - l|} (\eta N_0)^{-|x_i - l|} (\epsilon)^{|x_i - l|} \\
&\geq 2^{-\frac{\eta \epsilon N_0}{2}} (\lambda \epsilon)^{\frac{\eta \epsilon N_0}{2}} e^{-\lambda \eta N_0} \\
&\geq e^{-c \eta N_0}, \text{ for some positive } c.
\end{aligned}$$

Then using **Lemma 3.6**, we have for some $c > 0$,

$$\begin{aligned}
Q(E_{x_i}[e^{\int_0^{\eta N_0} dB_{X(s)}(s)} \mathbf{1}_l(X(s))]) \leq e^{-\frac{\epsilon N_0}{12}} &\leq Q(E_{x_i}[e^{\int_0^{\eta N_0} dB_{X(s)}(s)}; X \sim \gamma]) \leq e^{-\frac{\epsilon N_0}{12}} \\
&\leq Q(e^{\int_0^{\eta N_0} dB_{\gamma(s)}(s) + \omega(\gamma)} P_{x_i}(X \sim \gamma)) \leq e^{-\frac{\epsilon N_0}{12}} \\
&\leq Q(e^{\int_0^{\eta N_0} dB_{\gamma(s)}(s) + c \eta N_0} P_{x_i}(X \sim \gamma)) \leq e^{-\frac{\epsilon N_0}{12}} + e^{-c \eta N_0} \\
&\leq Q(e^{\int_0^{\eta N_0} dB_{\gamma(s)}(s)} \leq e^{-\frac{\epsilon N_0}{12} - c \eta N_0}) + e^{-c \eta N_0} \\
&= Q(B_0(\eta N_0) \leq -\frac{\epsilon N_0}{12} - c \eta N_0) + e^{-c \eta N_0},
\end{aligned}$$

which clearly tends to 0 as N_0 tends to ∞ . This is enough to establish the claim and that finishes the proof of the lemma. \square

With this lemma we can now conclude that

Lemma 4.3. *For any $\eta > 0$, there is an N_1 such that $\forall N_0 > N_1$,*

$$Q(\sup \sum_{j=0}^{\frac{t}{N_0} - 1} \mathbf{1}_{G^c}(i_j, j) \geq \eta t / N_0) \leq e^{-\frac{\delta t}{N_0}}.$$

where the sup is taken over all traces compatible with paths starting at zero and having less than Mt jumps in time interval $[0, t]$.

Proof. This follows from elementary large deviations for the binomial random variables and **Lemma 4.2**. \square

The next lemma says that the contribution of any collection of a restricted number of blocks can not be very big with high probability.

Lemma 4.4. *Given $\epsilon > 0$ and $\eta > 0$ as in **Lemma 4.2**, for all N_0 sufficiently large,*

$$Q(\exists(x_{i_1}, x_{i_2}, \dots, x_{i_{\frac{t}{N_0}}}) \text{ compatible with } \leq Mt \text{ jumps, } J \subset \{1, 2, \dots, \frac{t}{N_0}\})$$

$$\text{with } |J| \leq \frac{\eta t}{N_0}, :$$

$$\prod_J E_{x_{i_j}} [e^{\int_0^{N_0} dB_{X(s)}^{jN_0}(s)} \geq e^{\frac{\epsilon t}{10}}] \leq e^{-c\epsilon t}.$$

Proof. First fix a skeleton $(x_{i_1}, x_{i_2}, \dots, x_{i_{\frac{t}{N_0}}})$ For a given subset $J \subset \{1, 2, \dots, \frac{t}{N_0}\}$ with $|J| \leq \frac{\eta t}{N_0}$ and skeleton $(x_{i_1}, x_{i_2}, \dots, x_{i_{\frac{t}{N_0}}})$, we have by Chebychev's inequality,

$$Q[\prod_J E_{x_{i_j}} [e^{\int_0^{N_0} dB_{X(s)}^{jN_0}(s)} \geq e^{\frac{\epsilon t}{10}}] \leq e^{|J|/2} e^{-\epsilon t/10}.$$

We now complete the proof by first summing over the set of possible J and then the set of possible skeletons $(x_{i_1}, x_{i_2}, \dots, x_{i_{\frac{t}{N_0}}})$ (using **Lemma 4.1**) \square

In the same way we have

Lemma 4.5. *Given $\epsilon > 0$ as in **Lemma 4.2**, for all N_0 sufficiently large,*

$$Q(\exists(x_{i_1}, x_{i_2}, \dots, x_{i_{\frac{t}{N_0}}}) \text{ compatible with } \leq Mt \text{ jumps, any } J \subset \{1, 2, \dots, \frac{t}{N_0}\})$$

$$\prod_J E_{x_j} [e^{\int_0^{N_0} dB_{X(s)}^{jN_0}(s)} I_{[MN_0, \infty)}(N(X, N_0))] \geq e^{\frac{\epsilon t}{10}}] \leq e^{-c\epsilon t}.$$

We can now give the proof of **Theorem 2.11**.

Proof. (of **Theorem 2.11**) For a given skeleton $(x_0, x_1, \dots, x_{\frac{t}{N_0}-1})$, compatible with a path starting at zero with no more than Mt jumps, we have that

$$E_0[e^{\int_0^t dB_{X(s)}(s)} \prod_{j=0}^{\frac{t}{N_0}-1} 1_{x_{i_j}}(X(jN_0))] \leq \prod_{j=0}^{\frac{t}{N_0}-1} E_{x_j} [e^{\int_0^t dB_{X(s)}^{jN_0}(s)}]$$

$$= \prod_B E_{x_j} [e^{\int_0^t dB_{X(s)}^{jN_0}(s)}] \prod_G E_{x_j} [e^{\int_0^t dB_{X(s)}^{jN_0}(s)}]$$

where G is the set of j so that block (i_j, j) is a good block, B is the set of j where this condition fails. By **Lemma 4.3** we have that outside of probability e^{-8t/N_0} that for every possible skeleton $|B| < \eta t/N_0$. By **Lemma 4.4** (outside of probability $e^{-c\epsilon t}$) we have for all skeletons and all choices of B , with $|B| < \frac{\eta}{N_0}t$,

$$\prod_B E_{x_j} [e^{\int_0^t dB_{X(s)}^{jN_0}(s)}] < e^{\epsilon t/10}.$$

Moreover we can split up $\prod_G E_{x_j} [e^{\int_0^t dB_{X(s)}^{jN_0}(s)}]$ as

$$\begin{aligned} & \prod_G (E_{x_j} [e^{\int_0^t dB_{X(s)}^{jN_0}(s)}] 1_{(MN_0, \infty)}(N(X, N_0))) + E_{x_j} [e^{\int_0^t dB_{X(s)}^{jN_0}(s)}] 1_{[0, MN_0]}(N(X, N_0)) \\ &= \sum_{I \subset G} \prod_{j \in I} E_{x_j} [e^{\int_0^t dB_{X(s)}^{jN_0}(s)}] 1_{[0, MN_0]}(N(X, N_0)) \times \prod_{j \notin I} E_{x_j} [e^{\int_0^t dB_{X(s)}^{jN_0}(s)}] 1_{(MN_0, \infty)}(N(X, N_0)) \end{aligned}$$

By **Lemma 4.5**, (outside of probability $e^{-c\epsilon t}$) for all skeletons and subsets I , simultaneously, the term $\prod_{j \notin I} E_{x_j} [e^{\int_0^t dB_{X(s)}^{jN_0}(s)}] 1_{(MN_0, \infty)}(N(X, N_0)) < e^{\epsilon t/10}$ so with large probability,

$$\begin{aligned} \prod_G E_{x_j} [e^{\int_0^t dB_{X(s)}^{jN_0}(s)}] &\leq e^{\epsilon t/10} \sum_{I \subset G} \prod_{j \in I} E_{x_j} [e^{\int_0^t dB_{X(s)}^{jN_0}(s)}] 1_{[0, MN_0]}(N(X, N_0)) \\ &\leq e^{\epsilon t/10} 2^{t/N_0} e^{t(\gamma(\kappa) + \epsilon/3)} \end{aligned}$$

by definition of a good block. That is we have shown that provided N_0 was fixed sufficiently large, then outside probability $e^{-c\epsilon t} + e^{-t/N_0}$ for every compatible skeleton

$$E_0 [e^{\int_0^t dB_{X(s)}(s)} \prod_{j=0}^{\frac{t}{N_0}-1} 1_{x_{i_j}}(X(jN_0))] \leq e^{\epsilon t/10} e^{\epsilon t/10} 2^{t/N_0} e^{t(\gamma(\kappa) + \epsilon/3)}.$$

By **Lemma 4.1**, the number of skeletons compatible with no more than Mt jumps is bounded by $e^{\epsilon t/10^6}$. Thus we have outside the above probability

$$E[e^{\int_0^t dB_{X(s)}(s)} I_{[0, Mt]}(N(X, t))] \leq e^{t(\gamma(\kappa) + 2\epsilon/3)}$$

The Theorem now follows from the estimate

$$E[e^{\int_0^t dB_{X(s)}(s)} I_{(Mt, \infty)}(N(X, t))] \leq e^{-cMt}.$$

□

References

- [Ad90] R. Adler. An Introduction to Continuity, Extreme and Related Topics for General Gaussian Processes. *Inst. Math. Stat. Lecture Notes*, 1990.
- [CM94] René A. Carmona and S. A. Molchanov. Parabolic Anderson problem and intermittency. *Mem. Amer. Math. Soc.*, 108(518):viii+125, 1994.
- [CMV94] René A. Carmona, S. A. Molchanov and F. Viens. Sharp upper bound on the almost-sure exponential behavior of a stochastic partial equation. *Random Oper. Stochastic Equations*, 4,no. 1,43-49 1996.
- [D93] R. Durrett. Ten Lectures on Particle Systems, Ecole d't de Probabilits de Saint Flour, XXIII Springer, New York, Berlin.1993
- [Lig85] Thomas M. Liggett. An improved subadditive ergodic theorem. *Ann. Probab.*, 13(4):1279–1285, 1985.
- [Mou01] Thomas S. Mountford. A note on limiting behaviour of disastrous environment exponents. *Electron. J. Probab.*, 6:no. 1, 9 pp. (electronic), 2001.
- [MR94] Stanislav Molchanov and Alexander Ruzmaikin. Lyapunov exponents and distributions of magnetic fields in dynamo models. In *The Dynkin Festschrift*, pages 287–306. Birkhäuser Boston, Boston, MA, 1994.
- [Sh97] Tokuzo Shiga. Exponential decay rate of teh survival probability in a disasterous random environment. *Prob Related Fields* 108, 417'439, 1997.
- [ShF98] T. Furuoya and T. Shiga. Sample Lyapunov Exponenet for a classs of linear Markov systems over Z^d . *Osaka J. Math*, 35, 35-72 ,1998.

M. Cranston	T. S. Mountford	T. Shiga
Department of Mathematics	DMA	Mathematics
University of Rochester	EPFL	Tokyo Inst. of Tech.
Rochester, New York 14627	CH-1015	Tokyo
USA	Switzerland	Japan
cran@math.rochester.edu	thomas.mountford@epfl.ch	shiga@neptune. ap. titech. ac. jp