

## References

- Banerjee, S., Gelfand, A. E., Finley, A. O., and Sang, H. (2008). Gaussian predictive process models for large spatial data sets. *Journal of the Royal Statistical Society, Series B: Statistical Methodology*, **70**(4), 825–848.
- Calder, K. A. (2007). Dynamic factor process convolution models for multivariate spacetime data with application to air quality assessment. *Environmental and Ecological Statistics*, **14**, 229–247.
- Cressie, N. and Johannesson, G. (2008). Fixed rank kriging for very large data sets. *Journal of the Royal Statistical Society, B*, **70**, 209–226.
- Du, J., Zhang, H., and Mandrekar, V. (2008). Fixed-domain asymptotic properties of tapered maximum likelihood estimators. *Annals of Statistics (under revision)*.
- Fuentes, M. (2007). Appropriate likelihood for large irregularly spaced spatial data. *Journal of the American Statistical Association*, **102**, 321–331.
- Furrer, R., Genton, M. G., and Nychika, D. (2006). Covariance tapering for interpolation of large spatial datasets. *Journal of Computational and Graphical Statistics*, **15**(3), 502–523.
- Gelfand, A., Banerjee, S., and Gamerman, D. (2005). Spatial process modeling for univariate and multivariate dynamic spatial data. *Environmetrics*, **16**, 465–497.
- Gelfand, A. E., Schmidt, A. M., Banerjee, S., and Sirmans, C. F. (2004). Non-stationary multivariate process modeling through spatially varying coregionalization. *Test (Madrid)*, **13**(2), 263–294.
- Higdon, D. (1998). A process-convolution approach to modelling temperatures in the North Atlantic Ocean (Disc: P191-192). *Environmental and Ecological Statistics*, **5**, 173–190.
- Paciorek, C. and Schervish, M. J. (2006). Spatial modeling using a new class of non-stationary covariance functions. *Environmetrics*, **17**, 483–506.
- Rue, H. and Tjelmeland, H. (2002). Fitting Gaussian Markov random fields to Gaussian fields. *Scandinavian Journal of Statistics*, **29**(1), 31–49.
- Stein, M. L. (2008). A modeling approach for large spatial datasets. *Journal of the Korean Statistical Society*, **37**, 3–10.
- Stein, M. L., Chi, Z., and Welty, L. J. (2004). Approximating likelihoods for large spatial data sets. *Journal of the Royal Statistical Society, Series B: Statistical Methodology*, **66**(2), 275–296.
- Vecchia, A. V. (1998). Estimation and model identification for continuous spatial processes. *Journal of the Royal Statistical Society, B*, **50**, 297–312.
- Ver Hoef, J. M., Cressie, N. A. C., and Barry, R. P. (2004). Flexible spatial models based on the fast fourier transform (FFT) for cokriging. *Journal of Computational and Graphical Statistics*, **13**, 265–282.