

ON SOME MULTIPLE DECISION PROBLEMS

by

Ming-Wei Lu
Purdue University

Department of Statistics
Division of Mathematical Sciences
Mimeograph Series #464

*This research was supported by the Office of Naval Research contract N00014-67-A-0226-0014 at Purdue University. Reproduction in whole or in part is permitted for any purpose of the United States Government.



INTRODUCTION

In practice, there often do arise situations in which the researcher wishes to choose the best (or the t best) from a group of k populations, where the "bestness" of a population is based on ranking them according to the value of some characteristic of interest. In such situations, the classical tests of homogeneity are inadequate in the sense that they have not been designed to answer several possible questions in which the researcher may really be interested. Bahadur [1], Mosteller [54] and Paulson [59] were among the first research workers to recognize the inadequacy of the usual tests of homogeneity and to consider more meaningful formulations in order to answer these questions. These developments set the stage for the early investigations of multiple decision problem which have now come to be known as selection and ranking problems.

In the theory of selection and ranking procedures, there are two basic formulations to the problem. The first one is called the indifference zone formulation due to Bechhofer [13] and the other is the subset formulation due to Gupta [29]. The goal of the basic problem in the formulation of Bechhofer is to choose one of the populations as the best. The researcher is required to specify an "indifference zone" in the parameter space and the

procedure determines the smallest sample size so that a certain probability condition is satisfied whenever the unknown parameters lie in the "preference zone". For example, if we are interested in selecting the population with the largest mean in $N(\theta_i, 1)$, $i=1, \dots, k$, the indifference zone is the set of all θ 's such that the largest and the second largest differ by an amount $\leq \delta^*$, whereas the "preference zone" is the set of all θ 's such that the above difference $> \delta^*$. Other contributions to this indifference zone formulation have been made by Bechhofer and Sobel [16], Sobel and Huyett [70], Barr and Rizvi [12], Desu and Sobel [22], Bechhofer, Kiefer and Sobel [15], Santner [64] and others. The goal of the basic problem in the subset selection formulation of Gupta is to select a subset of the given populations which depends on the outcome of the experiments and is not fixed in advance such that it includes the best population with a specified minimum probability regardless of the unknown configuration of parameters, i.e., over the whole parameter space. Some recent results in the area of subset selection formulation are Gnanadesikan and Gupta [28], Gupta and Studden [43], Gupta and Panchapakesan [38], Gupta and Santner [41], Huang [46] and Wong [72].

Many problems in reliability can be considered in the context of selection and ranking problems. For example, one may wish to choose one or more of the several systems or components which has the largest mean life or the largest median life. In general, reliability problems also deal where the distributions are unknown but assumed to belong to a class of distributions such as that

having an increasing failure rate (IFR). Such distributions form a special cases of what are now commonly known as restricted families of probability distributions. The investigations of Barlow and Gupta [5] form the initial efforts on ranking problems for such families. Other contributions to selection problem for the restricted families of probability distributions are Gupta and Panchapakesan [39,40] and Patel [58]. This area of research still remains largely unexplored.

The main investigation of this thesis is to propose and study selection procedures for some problems.

Chapter I deals with some selection and ranking procedures for restricted families of probability distributions. In Section 1.1, definitions of various partial orderings on the space of distributions are given. In Section 1.2, we propose and study a selection rule for distributions which are convex-ordered with respect to specified distribution G . Some properties of this selection rule are discussed. The asymptotic relative efficiencies of this rule with respect to some other selection rules are derived. Section 1.3 deals with the selecting the best population using the indifference zone approach. In Section 1.5, we propose and study a selection rule for distributions which are s -ordered with respect to G where we are interested in the scale parameter case. The estimation of ordered parameters from the k unknown distributions is discussed in Section 1.8.

Chapter II discusses some interval estimation problems from k populations. We are interested in finding the smallest sample

size N to be chosen from each population such that the probability that a given confidence interval I which is based on $T_{[k]} = \max_{1 \leq i \leq k} T_i$ (T_i is an appropriate statistic from i -th population) contains at least one good population, is at least P^* , where P^* is a specified number, $0 < P^* < 1$. Also we are interested in finding the smallest sample size N such that the probability that I contains all good populations (or excludes all bad populations) is at least P^* . Section 2.2 deals with the above problems for the location parameter case. The infima of coverage probabilities are obtained. For scale parameter case, the above problems are investigated in Section 2.3. In Section 2.4, we illustrate the above results by means of two examples.

In some situations, one has to deal with a vector-valued parameter $\underline{\lambda}_i$ associated $\underline{\lambda}_i$. In such cases one may consider comparing populations or $\underline{\lambda}_i$'s in terms of majorization and weak majorization of these vectors. Chapter III deals with such problems. The parameter space is partially ordered by means of majorization or weak majorization. Selection procedures are proposed and studied. In Section 3.2, a class of procedures R_h for selecting the best population is defined. A sufficient condition is obtained for the infimum of the probability of a correct selection to be Schur-convex in $\underline{\lambda}$. Also another sufficient condition for the same infimum of the probability of a correct selection to be nondecreasing and Schur-convex in $\underline{\lambda}$ is obtained in Section 3.3. Section 3.5 and 3.6 deal with selection procedures

for multivariate normal distributions in terms of majorization and weak majorization. Various cases corresponding to the known or unknown common covariance matrix Σ are studied. Properties of these selection procedures are also established.

CHAPTER I
SELECTION PROCEDURES FOR RESTRICTED FAMILIES
OF PROBABILITY DISTRIBUTIONS

1.1 Introduction

In many problems, especially those in reliability theory, one is interested in using a model for life length distribution which belongs, for example, to a family of distributions having increasing failure rate (IFR), or increasing failure rate on the average (IFRA). Such distributions form special cases of what are now commonly known as restricted families of probability distributions. These are defined more precisely later in this section. The idea of using such families stems from the fact that in many cases the experimenter cannot specify the model (distribution) exactly but is able to say whether it comes from a family of distributions such as IFR, IFRA. Families of probability distributions of these types have been studied by several authors, see, for example, Barlow, Marshall and Proschan [7]. These authors have mainly concerned themselves with probabilistic aspects of these distributions. To some extent, there have been some investigations dealing with statistical inference for some of these families; see for example, Barlow and Proschan [8], [9], Barlow and Doksum [3].

In this chapter we are interested in studying multiple decision procedures for k ($k \geq 2$) populations which are themselves unknown but which are assumed to belong to a restricted family. First we give some notations and definitions. A binary ordering relation (\prec) is called a partial ordering in the space of probability distributions if

- (a) $F \prec F$ for all distributions F , and
- (b) $F \prec G, G \prec H$ imply $F \prec H$.

Note that $F \prec G$ and $G \prec F$ do not necessarily imply $F \equiv G$.

We now define some of the special order relations of interest to us (see Barlow and Gupta [5]).

- (i) F is said to be convex with respect to G (written $F \prec_C G$) if and only if $G^{-1}F(x)$ is convex on the support of F .
- (ii) F is said to be star-shaped with respect to G (written $F \prec_* G$) if and only if $F(0) = G(0) = 0$ and $\frac{G^{-1}F(x)}{x}$ is increasing in $x \geq 0$ on the support of F .
- (iii) F is said to be r -ordered with respect to G ($F \prec_r G$) if and only if $F(0) = G(0) = \frac{1}{2}$ and $\frac{G^{-1}F(x)}{x}$ is increasing (decreasing) for x positive (negative) on the support of F .
- (iv) F is said to be s -ordered with respect to G ($F \prec_s G$) if and only if $F(0) = G(0) = \frac{1}{2}$ and $G^{-1}F$ is concave-convex about the origin, on the support of F .

If $G(x) = 1 - e^{-\lambda x}$, $x \geq 0$, then $F \prec_C G$ is equivalent to saying that F has increasing failure rate (IFR). The class of IFR distributions has been studied by Barlow, Marshall and Proschan [7].

Again if $G(x) = 1 - e^{-\lambda x}$, $x \geq 0$, $F \prec_* G$ is equivalent to saying that F has increasing failure rate on average (IFRA). The class of IFRA distributions has been studied by Barlow, Esary and Marshall [4]. The r -ordering has been defined and investigated by Lawrence [48]. The s -ordering and c -ordering have been studied by Van Zwet [71] and Lawrence [48].

In the statistical literature, selection problems for restricted families were first investigated by Barlow and Gupta [5]. Some further results in this direction and a review of some important results concerning inequalities for restricted families and problems of inference for such families have been given by Gupta and Panchapakesan [39]. The selection of the population with largest α -quantile from distributions which are star-shaped with respect to the folded normal distribution has been considered by Gupta and Panchapakesan [40]. In a recent paper, Patel [58] has studied the selection of IFR populations which differ only in the scale parameters. His procedure is based on the total life statistic until r -th failure.

In Section 1.2, we propose and study a selection rule for distributions which are \prec_c ordered with respect to a specified distribution G assuming there exists a best one. Some properties of this rule are discussed. The infimum of the probability of a correct selection is obtained and an asymptotic expression is also given. We also study the asymptotic relative efficiencies of this rule with respect to some selection procedures. Section 1.3

deals with the selecting the best population in the frame work of Section 1.2 using the indifference zone approach. In Section 1.4, we propose a selection procedure for distributions that are \prec_* ordered with respect to G. Section 1.5 deals with selection procedure with respect to the means for distributions that are s-ordered with respect to G where we are interested in the scale parameter case. The distribution of V_j (see Section 1.5) is also investigated. Section 1.8 deals with estimation of ordered parameters from k unknown distributions where we are interested in the scale parameter case.

1.2 Selection rules for distributions \prec_c ordered with respect to a specified distribution G.

Let \mathfrak{F} be the class of absolutely continuous distribution functions F on R with positive and right-(or left-) continuous density f on the interval where $0 < F < 1$. It follows that the inverse function F^{-1} is uniquely determined on (0,1). We take $F^{-1}(0)$ and $F^{-1}(1)$ to be equal to the left hand and right hand endpoints of the support of F. For $F, G \in \mathfrak{F}$, consider the following transformation (see Barlow and Doksum [3])

$$(1.2.1) \quad H_F^{-1}(t) = \int_{F^{-1}(0)}^{F^{-1}(t)} g[G^{-1}F(u)]du, \quad 0 \leq t \leq 1.$$

We assume that G is always fixed. Since H_F^{-1} (the inverse of H) is strictly increasing on [0,1], H_F is a distribution. We know that $F \prec_c G$ if and only if H_F is convex on the interval where $0 < H_F < 1$.

Since G is assumed known we can estimate H_F^{-1} by substituting the empirical distribution F_n of F ; that is

$$(1.2.2) \quad H_n^{-1}(t) = H_{F_n}^{-1}(t) = \int_{F^{-1}(0)}^{F_n^{-1}(t)} g[G^{-1}F_n(u)] du$$

and

$$(1.2.3) \quad H_n^{-1}\left(\frac{r}{n}\right) = \int_{F^{-1}(0)}^{X_{r,n}} g[G^{-1}F_n(u)] du = \sum_{i=1}^r g\left[G^{-1}\left(\frac{i-1}{n}\right)\right] (X_{i,n} - X_{i-1,n})$$

where $X_{i,n}$ is the i -th order statistic in a sample of size n from F and $X_{0,n} \equiv 0$.

If $G(x) = 1 - e^{-x}$ for $x \geq 0$, then (1.2.3) can be written as

$$(1.2.4) \quad H_n^{-1}\left(\frac{r}{n}\right) = \frac{1}{n} [X_{1,n} + \dots + X_{r-1,n} + (n-r+1)X_{r,n}].$$

We say that $X_{1,n} + \dots + X_{r-1,n} + (n-r+1)X_{r,n}$ is the total life statistic until r -th failure from F . Thus, $H_n^{-1}\left(\frac{r}{n}\right) = \frac{1}{n}$ {total life statistic until r -th failure}.

(A) Selection procedure and its properties

Let π_1, \dots, π_k be k populations. The random variable X_i associated with π_i has distribution function F_i , $i = 1, 2, \dots, k$, where $F_i \in \mathcal{F}$ ($i = 1, \dots, k$). Let $F_{[k]}$ denote the cumulative distribution function (c.d.f.) of the "best" population. For example, if we are interested in the quantile selection problem, the "best" population may be defined as that $F_{[k]}$ (unknown) which satisfies (a) below.

We assume that

$$(a) \quad F_{[i]}(x) \geq F_{[k]}(x) \text{ for all } x, i = 1, \dots, k-1;$$

(b) there exists a distribution G such that

$$F_{[i]} \leq G \quad i = 1, \dots, k,$$

where \leq denotes a partial ordering relation in the space of probability distributions. It should be pointed out that the condition (a) above may also imply that $F_{[k]}$ is the distribution with the largest (smallest) parameter. For example, if

$F_i(x) = F(\frac{x}{\theta_i})$ for $x \geq 0$, $\theta_i > 0$ ($i = 1, \dots, k$), then $F_{[k]}$ is the distribution with the largest θ_i . We are given a sample of size n from each π_i ($i = 1, \dots, k$). Our goal is to select a subset from the k populations so as to include the population with $F_{[k]}$. Let $\Omega = \{F = (F_1, \dots, F_k) : \exists a_j \text{ such that } F_i(x) \geq F_j(x) \forall i \neq j\}$. Let

$$(1.2.5) \quad T_i = \sum_{j=1}^r a_j X_{i;j,n} \quad \text{for } i = 1, \dots, k.$$

$$(1.2.6) \quad T = \sum_{j=1}^r a_j Y_{j,n}$$

where $X_{i;j,n}$ is the j -th order statistic from F_i , $Y_{j,n}$ is the j -th order statistic from G , r is a fixed positive integer ($1 \leq r \leq n$),

$$a_j = g G^{-1}(\frac{j-1}{n}) - g G^{-1}(\frac{j}{n}) \quad \text{for } j = 1, \dots, r-1$$

and $a_r = g G^{-1}(\frac{r-1}{n})$.

For selecting a subset containing $F_{[k]}$, we propose the selection rule R_1 as follows:

R_1 : Select population π_i if and only if

$$(1.2.7) \quad T_i \geq c_1 \max_{1 \leq j \leq k} T_j$$

where $c_1 = c_1(k, P^*, n, r)$ is some number between 0 and 1 which is determined as to satisfy the probability requirement

$$(1.2.8) \quad \inf_{\Omega} P\{CS|R_1\} \geq P^*$$

where CS stands for a correct selection, i.e., the selection of any subset which contains the population with distribution $F_{[k]}$. Let $T_{(i)}$ be associated with $F_{[i]}$ and $W_i(x)$ be the c.d.f. of $T_{(i)}$ from $F_{[i]}$. We now state a few preliminary lemmas.

Lemma 1.2.1. (Lehman [49] p. 112) Let $F(x)$ be a distribution function on the real line. If $\psi(x)$ is any nondecreasing function of x , then $E\psi(X)$ is a nonincreasing function of F , i.e., if $F_1(x) \leq F_2(x)$ for all x , then $\int \psi(x) dF_2(x) \leq \int \psi(x) dF_1(x)$.

Lemma 1.2.2. Let X_1, \dots, X_n be i.i.d. with distribution $F(x)$. Let ψ be a function of X_1, \dots, X_n which is nondecreasing in each of its arguments. Then $E\psi(X_1, \dots, X_n)$ is nonincreasing function of F .

Lemma 1.2.3. (Gupta and McDonald [35]) Let $\underline{X} = (X_{11}, \dots, X_{1n_1}, \dots, X_{k1}, \dots, X_{kn_k})$ be a vector valued random variable of $\sum_{i=1}^k n_i$ (≥ 1) independent components with X_{ij} having the distribution $F_i(x)$, $j = 1, \dots, n_i$, $i = 1, \dots, k$. Let ψ be a function of $X_{11}, \dots, X_{1n_1}, \dots, X_{k1}, \dots, X_{kn_k}$ which, for any fixed i , is a nondecreasing (nonincreasing) function of X_{i1}, \dots, X_{in_i} when the other components of \underline{X} are held fixed. Then $E\psi(\underline{X})$ is a nonincreasing (nondecreasing) function of F_i .

The following two lemmas are from Barlow and Proschan [8].

Lemma 1.2.4. (a) $\phi\left(\sum_{i=1}^n a_i X_i\right) - \phi(0) \leq \sum_{i=1}^n a_i [\phi(X_i) - \phi(0)]$ for all $0 \leq X_1 \leq \dots \leq X_n \leq b$ and for all convex ϕ on $[0, b]$ if and only if $0 \leq \bar{A}_i \equiv \sum_{j=i}^n a_j \leq 1$ for $i = 1, 2, \dots, n$.

(b) $\phi\left(\sum_{i=1}^n a_i X_i\right) - \phi(0) \geq \sum_{i=1}^n a_i [\phi(X_i) - \phi(0)]$ for all $0 \leq X_1 \leq \dots \leq X_n$ and for all convex ϕ on $(-\infty, \infty)$ if and only if $\bar{A}_1 \geq 1, \bar{A}_2 \geq 1, \dots, \bar{A}_k \geq 1, \bar{A}_{k+1} \leq 0, \dots, \bar{A}_n \leq 0$ for some k ($0 \leq k \leq n$).

Lemma 1.2.5. (a) $\phi\left(\sum_{i=1}^n a_i X_i\right) \leq \sum_{i=1}^n a_i \phi(X_i)$

for all star-shaped ϕ on $[0, b]$ and all $0 \leq X_1 \leq \dots \leq X_n \leq b$ if and only if there exists k ($1 \leq k \leq n$) such that

$$0 \leq \bar{A}_1 \leq \dots \leq \bar{A}_k \leq 1 \quad \text{and} \quad \bar{A}_{k+1} = \dots = \bar{A}_n = 0.$$

(b) $\phi\left(\sum_{i=1}^n a_i X_i\right) \geq \sum_{i=1}^n a_i \phi(X_i)$ for all $0 \leq X_1 \leq \dots \leq X_n$ and for all star-shaped ϕ on $(-\infty, \infty)$ if and only if there exists k ($1 \leq k \leq n$) such that

$$\bar{A}_1 \geq \dots \geq \bar{A}_k \geq 1; \quad \bar{A}_{k+1} = \dots = \bar{A}_n = 0.$$

Lemma 1.2.6. Let F_1, F_2 be two distribution functions such that $F_1(x) \geq F_2(x) \forall x$.

$$\text{Let } T_i = \sum_{j \in \Delta} b_j X_{i;j,n} \quad i = 1, 2$$

where $b_j > 0$ for $j \in \Delta, \Delta \subset \{1, 2, \dots, n\}$ and $X_{i;j,n}$ is the j -th order statistic from $F_i, i = 1, 2$.

$$\text{Then } P[T_1 \leq x] \geq P[T_2 \leq x].$$

Proof.

$$\text{Let } \psi_i (X_{i1}, \dots, X_{in}) = \begin{cases} 1 & \text{if } T_i \geq x \\ 0 & \text{otherwise} \end{cases}$$

where X_{i1}, \dots, X_{in} are n observations from F_i ($i = 1, 2$).

Since $\psi_i (X_{i1}, \dots, X_{in})$ is nondecreasing in each of its arguments, by Lemma 1.2.2 we have

$$E \psi_1 (X_{11}, \dots, X_{1n}) \leq E \psi_2 (X_{21}, \dots, X_{2n})$$

$$\text{That is } P[T_1 \geq x] \leq P[T_2 \geq x]$$

This proves the Lemma.

We now state and prove following theorem which is more general than that of Patel [58].

Theorem 1.2.1. Let $F_i, G \in \mathfrak{F}$, $F_i(x) \geq F_{[k]}(x) \forall x$ and $i = 1, 2, \dots, k$,

$F_{[k]}(0) = 0$ and $F_{[k]} \underset{c}{<} G$.

If $a_j \geq 0$ for $j = 1, 2, \dots, r$, $G^{-1}(0) \leq 0$, $g G^{-1}(0) \leq 1$ and $a_r \geq c_1$, then

$$(1.2.9) \quad P[CS|R_1] \geq \int_{G^{-1}(0)}^{\infty} G_T^{k-1} \left(\frac{x}{c_1} \right) dG_T(x)$$

where $G_T(x)$ is the c.d.f. of T .

Proof. $P[CS|R_1] = P[T_{(k)} \geq c_1 T_{(i)}, i \neq k]$

$$= \int_0^{\infty} \prod_{i=1}^{k-1} W_i^{k-1} \left(\frac{x}{c_1} \right) dW_k(x)$$

$$\geq \int_0^{\infty} W_k^k \left(\frac{x}{c_1} \right) dW_k(x) \quad (\text{By Lemma 1.2.6})$$

$$= P[Z_k \geq c_1 Z_j, j \neq k]$$

where Z_1, \dots, Z_k are i.i.d. with c.d.f. $W_k(x)$.

Let $\varphi(x) = G^{-1}F_{[k]}(x)$.

Note that

$$(1.2.10) \quad Z_i \stackrel{\text{st}}{=} \sum_{j=1}^r a_j X_{i;j,n}^* \quad i = 1, \dots, k,$$

where $X_{i;j,n}^*$ is the j -th order statistic in a sample of size n from $F_{[k]}$, $i = 1, \dots, k$.

$$(1.2.11) \quad P[Z_k \geq c_1 \max_{1 \leq j \leq k} Z_j] = P[\varphi(\frac{1}{c_1} Z_k) \geq \varphi(Z_i), i = 1, \dots, k-1]$$

Since $\sum_{j=1}^r a_j = g G^{-1}(0) \leq 1$ and $a_j \geq 0 \forall j = 1, \dots, r$, then, by Lemma 1.2.4 (a) and (1.2.10),

$$(1.2.12) \quad \varphi(Z_i) \leq \sum_{j=1}^r a_j \varphi(X_{i;j,n}^*).$$

Since $\frac{1}{c_1} a_r \geq 1$ and $\frac{1}{c_1} \sum_{j=i}^r a_j \geq 1$ for $i = 1, \dots, r$, we have, by Lemma 1.2.4 (b) and (1.2.10),

$$(1.2.13) \quad \varphi(\frac{1}{c_1} Z_k) \geq \frac{1}{c_1} \sum_{j=1}^r a_j \varphi(X_{k;j,n}^*).$$

$$(1.2.14) \quad \varphi(X_{i;j,n}^*) \stackrel{\text{st}}{=} Y_{i;j,n}$$

where $Y_{i;j,n}$ is the j -th order statistic from G , $i = 1, 2, \dots, k$.

Thus from (1.2.11), (1.2.12), (1.2.13) and (1.2.14),

$$\begin{aligned} P[Z_k \geq c_1 \max_{1 \leq i \leq k} Z_i] &\geq P[\sum_{j=1}^r a_j Y_{k;j,n} \geq c_1 \sum_{j=1}^r a_j Y_{i;j,n}, \\ &\quad i = 1, 2, \dots, k-1] \\ &= \int_{G^{-1}(0)}^{\infty} G_T^{k-1}(\frac{x}{c_1}) dG_T(x) \end{aligned}$$

This completes the proof.

The constant $c_1 = c_1(k, P^*, n, r)$ satisfying (1.2.8) is determined by

$$\int_{G^{-1}(0)}^{\infty} G_T^{k-1} \left(\frac{x}{c_1} \right) dG_T(x) \geq P^* \quad \text{and}$$

$$g G^{-1} \left(\frac{r-1}{n} \right) \geq c_1.$$

We now consider two specific distributions $G(x)$. If $G(x) = 1 - e^{-x}$, $x \geq 0$, then we have following result which generalizes the result of Patel [58].

Corollary 1.2.1. If $F_i(x) \geq F_{[k]}(x) \forall x$ and $i = 1, \dots, k$, $F_{[k]}(0) = 0$, $F_{[k]} \prec_c G$, $G(x) = 1 - e^{-x}$, $x \geq 0$ and $n \geq \max\{r, \frac{r-1}{1-c_1}\}$, then

$$(1.2.15) \quad \inf_{\Omega} P[CS | R_1] = \int_0^{\infty} H^{k-1} \left(\frac{x}{c_1} \right) dH(x)$$

where $H(x)$ is the c.d.f. of a χ^2 random variable with $2r$ d.f.

Proof. If $G(x) = 1 - e^{-x}$ then $a_j = \frac{1}{n}$ for $j = 1, 2, \dots, r-1$

$$\text{and } a_r = \frac{1}{n} (n - r + 1).$$

$$\text{Also } \frac{1}{c_1} a_r \geq 1 \quad \text{iff} \quad n \geq \frac{r-1}{1-c_1}.$$

By Theorem 1.2.1 and the fact that $2nT$ is distributed as χ^2 with $2r$ d.f., the result follows.

If $G(x) = x$ for $0 < x < 1$, then we have the following result which is a special case of Theorem 2.1 of Barlow and Gupta [5].

Corollary 1.2.2. If $F_i(x) \geq F_{[k]}(x) \forall x$ and $i = 1, \dots, k$, $F_{[k]}(0) = 0$,

$$F_{[k]} \prec_c G \text{ and } G(x) = x \text{ for } 0 < x < 1,$$

then

$$(1.2.16) \quad \inf_{\Omega} P[CS|R_1] = \int_0^1 H_r^{k-1} \left(\frac{x}{c_1} \right) dH_r(x)$$

where $H_r(x)$ is the c.d.f. of the r -th order statistic from G .

Proof. If $G(x) = x$ $0 \leq x \leq 1$ then $a_j = 0$ for $j = 1, \dots, r-1$ and $a_r = 1$. By Theorem 1.2.1, the result follows.

We state and prove the following theorem about the asymptotic evaluations of the probability of a correct selection.

Theorem 1.2.2. If $F_i, G \in \mathfrak{F}$ for all $i = 1, \dots, k$ and

- (i) $F_i(x) \geq F_{[k]}(x) \forall x, i = 1, \dots, k$, and $F_{[k]} < G$,
- (ii) $G(x)$ has a differentiable density g in a neighborhood of its α -quantile η_α and $g(\eta_\alpha) \neq 0$,
- (iii) $g G^{-1}$ is uniformly continuous on $[0,1)$, $G^{-1}(x)$ is convex and there exists an $\eta, 0 < \eta < 1$, such that for $\eta \leq y < 1$, $g G^{-1}(y)$ is nonincreasing and $\frac{g G^{-1}(y)}{1-y}$ is nondecreasing in y , then as $n \rightarrow \infty$

$$(1.2.17) \quad P[CS|R_1] \geq \int_{-\infty}^{\infty} \phi^{k-1} \left[\frac{x}{c_1} + \frac{1-c_1}{c_1} \eta_\alpha g(\eta_\alpha) \left(\frac{n}{\alpha \bar{\alpha}} \right)^{\frac{1}{2}} \right] d\phi(x)$$

where $\frac{r}{n} \rightarrow \alpha$ as $n \rightarrow \infty$, $\bar{\alpha} = 1-\alpha$ and $\phi(x)$ is the standard normal c.d.f.

Proof. See the proof of Theorem 1.2.1, we have

$$(1.2.18) \quad P[CS|R_1] \geq P[Z_k \geq c_1 \max_{1 \leq j \leq k} Z_j]$$

where Z_1, \dots, Z_k are i.i.d. with c.d.f. $W_k(x)$ and $W_k(x)$ is the c.d.f. of $T_{(k)}$.

By Theorem 2.2 of Barlow and Van Zwet [11] and (iii),

$$(1.2.19) \quad \sup_{x \geq 0} \left| \int_{F_{[k]}^{-1}(0)}^x g[G^{-1}F_{[k]n}(u)]du - \int_{F_{[k]}^{-1}(0)}^x g[G^{-1}F_{[k]}(u)]du \right| \rightarrow 0 \text{ a.s.}$$

where $F_{[k]n}$ is the empirical distribution of $F_{[k]}$.

Then we have (see Barlow and Doksum [3]), for n large,

$$(1.2.20) \quad Z_i \stackrel{\sim}{st} Y_{i;r,n}$$

where $Y_{i;r,n}$ is the r -th order statistic from $H_{F_{[k]}}$ and $H_{F_{[k]}}^{-1}$ (the inverse of $H_{F_{[k]}}$) is defined in (1.2.1). Since $F_{[k]} \prec_c G$, therefore $H_{F_{[k]}}$ is convex. Since $G^{-1}(x)$ is increasing and convex, therefore $G^{-1}H_{F_{[k]}}(x)$ is convex. Since $H_{F_{[k]}} \prec_c G$ and $G^{-1}(0) \leq 0$, therefore $H_{F_{[k]}}^* \prec_c G$. In a manner similar to the Theorem 2.1 of Barlow and Gupta [5], we have

$$(1.2.21) \quad P[Y_{k;r,n} \geq c_1 \max_{1 \leq i \leq k} Y_{i;r,n}] \geq P[Y_{k;r,n}^* \geq c_1 Y_{i;r,n}^*, i \neq k]$$

where $Y_{i;r,n}^*$ is the r -th order statistic from G , $i = 1, \dots, k$. From (1.2.18), (1.2.20), (1.2.21) and using the fact that

$$Y_{i;r,n}^* \sim N\left(\eta_\alpha, \frac{\alpha \bar{\alpha}}{ng^2(\eta_\alpha)}\right),$$

the theorem follows.

Consistent with the basic probability requirement, we would like the size of the selected subset to be small. Let S be the size of the selected subset. One criterion of the efficiency of the procedure R_1 is the expected value of the size of the subset.

If in addition to the assumptions of the Theorem 1.2.1,

we assume that (a) $F_{[1]}(x) \geq F_{[i]}(x) \geq F_{[1]}(\frac{x}{\delta})$ for all $x \geq 0$,
 $i = 1, 2, \dots, k$, where $\delta > 1$ is given and (b) $G \prec_c F_{[1]}$.

We have the following theorem.

Theorem 1.2.3.

$$(1.2.22) \quad E(S|R_1) \leq k \int_{G^{-1}(0)}^{\infty} G_T^{k-1}(\frac{\delta}{c_1} x) dG_T(x)$$

where $G_T(x)$ is the c.d.f. of T .

$$\text{Proof. } E(S|R_1) = \sum_{i=1}^k P[T_i \geq c_1 T_j, \text{ for } j \neq i].$$

By Lemma 1.2.6, we have

$$P[T_i \geq c_1 T_j, j \neq i] \leq P[T_k^* \geq c_1 T_j^*, j = 1, \dots, k-1]$$

where T_j^* has c.d.f. $W_1(x)$ for $j = 1, \dots, k-1$ and $\frac{T_k^*}{\delta}$ has c.d.f. $W_1(x)$.

$$\text{So that } E(S|R_1) \leq k P[T_k^* \geq c_1 T_j^*, j = 1, \dots, k-1]$$

$$= k P[Z_k \delta \geq c_1 Z_j, j = 1, \dots, k-1]$$

where Z_1, \dots, Z_k are i.i.d. with cdf $W_1(x)$.

Hence

$$E(S|R_1) \leq k P[\frac{\delta}{c_1} Z_k \geq Z_j, j = 1, 2, \dots, k]$$

$$\text{Let } \varphi(x) = G^{-1} F_{[1]}(x)$$

since $G \prec_c F_{[1]}$ then $\varphi(x)$ is convex.

Using the same approach as in Theorem 1.2.1

and $\frac{\delta}{c_1} a_r \geq \frac{1}{c_1} a_r$, we have

$$\begin{aligned} P\left[\frac{\delta}{c_1} Z_k \geq Z_j, j = 1, 2, \dots, k\right] &\leq P\left[\frac{\delta}{c_1} Z_k^* \geq Z_j^*, j = 1, \dots, k\right] \\ &= \int_{G^{-1}(0)}^{\infty} G_T^{k-1}\left(\frac{\delta}{c_1} x\right) dG_T(x) \end{aligned}$$

where Z_1^*, \dots, Z_k^* are i.i.d. random variables with c.d.f. $G_T(x)$.

This proves the theorem.

Before discussing properties of the selection rule R_1 , we give some preliminary definitions. Let Ω denote the set of all k -tuples $\underline{F} = (F_1, \dots, F_k)$. Let $P_{\underline{F}}(i)$ denote

$$P_{\underline{F}}(i) = p_{\underline{F}}[\pi(i) \text{ is selected} | R]$$

when $\pi(i)$ is associated with $F_{[i]}$.

Definition 1.2.1. A rule R is strongly monotone in $\pi(i)$ if

$$P_{\underline{F}}(i) \text{ is } \begin{cases} \downarrow & \text{in } F_{[i]} & \text{when all other components of } \underline{F} \\ & & \text{are fixed} \\ \uparrow & \text{in } F_{[j]} (i \neq j) & \text{when all other components of} \\ & & \underline{F} \text{ are fixed.} \end{cases}$$

Definition 1.2.2. A rule R is monotone means

$$P_{\underline{F}}(i) \leq P_{\underline{F}}(j) \quad \text{for all } \underline{F} \in \Omega \text{ with } F_{[i]}(x) \geq F_{[j]}(x).$$

Definition 1.2.3. A rule R is unbiased if $P_{\underline{F}}(i) \leq P_{\underline{F}}(k)$ for all $\underline{F} \in \Omega$ with $F_{[i]}(x) \geq F_{[k]}(x)$.

Definition 1.2.4. A rule R is consistent with respect to Ω' means $\inf_{\Omega'} P[CS|R] \rightarrow 1$ as $n \rightarrow \infty$.

Theorem 1.2.4. If $a_i \geq 0$ for $i = 1, \dots, r$, then R_1 is strongly monotone in $\pi(i)$.

Proof.

$$\text{Let } \psi(\underline{x}) = \begin{cases} 1 & \text{if } T(i) \geq c \max_{1 \leq j \leq k} T(j) \\ 0 & \text{otherwise} \end{cases}$$

since $a_i \geq 0$, $T(i)$ is nondecreasing in each of its arguments for $i = 1, \dots, k$.

So that $\psi(\underline{x})$ is nondecreasing function of $X_{(i)1}, \dots, X_{(i)n}$ when the other components of \underline{x} are held fixed and $\psi(\underline{x})$ is nonincreasing function of $X_{(j)1}, \dots, X_{(j)n}$ ($j \neq i$) when the other components of \underline{x} are held fixed where $X_{(j)1}, \dots, X_{(j)n}$ are n observations from $F_{[j]}$ ($j = 1, \dots, k$). Since $E\psi(\underline{X}) = P_{\underline{F}}(i)$, by Lemma 1.2.3, it follows that R_1 is strongly monotone in $\pi(i)$.

Remark 1.2.1. (1) If a rule R is strongly monotone in $\pi(i)$ for all $i = 1, \dots, k$, then R is monotone and

$$\inf_{\Omega} P[CS|R] = \inf_{\Omega_0} P[CS|R]$$

where $\Omega_0 = \{\underline{F} = (F_1, \dots, F_k) : F_1 = \dots = F_k\}$.

(2) If R is monotone, then it is unbiased.

(3) If $F_i(x) = F(x, \theta_i)$, $i=1, \dots, k$ and T_i is a consistent estimator of θ_i , then R_1 is consistent.

(4) If $F_i, G \in \mathfrak{F}$, $F_i \preceq G$, $i=1, \dots, k$ and the condition (iii) (excluding $G^{-1}(x)$ is convex) of Theorem 1.2.2 is satisfied, we can show that R_1 is consistent.

The selection of the population with largest F_i ($i=1, \dots, k$) can be handled analogously. We assume $F_{[i]}(x) \leq F_{[1]}(x)$, $i=1, \dots, k$, and $F_{[1]} \prec_c G$. The rule for selecting the population with $F_{[1]}$ is

R_2 : Select population π_i if and only if

$$(1.2.23) \quad c_2 T_i \leq \min_{1 \leq j \leq k} T_j$$

where c_2 ($0 < c_2 \leq 1$) is determined so as to satisfy the basic requirement. In a manner similar to the proof of Theorem 1.2.1, if $F_i, G \in \mathfrak{F}$, $F_{[i]}(x) \leq F_{[1]}(x) \forall x$ and $i = 1, \dots, k$, $F_{[1]}(0) = 0$ and $F_{[1]} \prec_c G$ and if $a_j \geq 0$ for $j = 1, \dots, r$, $G^{-1}(0) \leq 0$, $g G^{-1}(0) \leq 1$ and $a_r \geq c_2$, then

$$(1.2.24) \quad P[CS|R_2] \geq \int_{G^{-1}(0)}^{\infty} \bar{G}_T^{k-1}(c_2 x) dG_T(x)$$

where $\bar{G}_T(x) = 1 - G_T(x)$.

(B) Efficiency of procedure R_1 under slippage configuration.

We consider slippage configuration $F_{[i]}(x) = F(\frac{x}{\delta})$, $i = 1, 2, \dots, k-1$, and $F_{[k]}(x) = F(x)$, $0 < \delta < 1$. Let $E(S|R)$ denote the expected subset size using the rule R . Then $E(S|R) - P[CS|R]$ is the expected number of non-best populations included in the selected subset. For a given $\epsilon > 0$, let $n_R(\epsilon)$ be the asymptotic sample size for which $E(S|R) - P[CS|R] = \epsilon$. We define the asymptotic relative efficiency $A R E(R, R^*, \delta)$ of R relative to R^* to be the limit as $\epsilon \rightarrow 0$ of the

$$\text{ratio } \frac{\eta_R(\epsilon)}{\eta_{R^*}(\epsilon)} \text{ i.e. ARE } (R, R^*; \delta) = \lim_{\epsilon \rightarrow 0} \frac{\eta_R(\epsilon)}{\eta_{R^*}(\epsilon)}.$$

Under the slippage configuration, for large n , we have (see Theorem 1.2.3)

$$(1.2.25) \quad E(S|R_1) = P[CS|R_1] + (k-1)P[T_{(1)} \geq c_1 \max_{i \neq 1} T_{(i)}]$$

$$(1.2.26) \quad P[T_{(1)} \geq c \max_{i \neq 1} T_{(i)}] \approx P[Y_1 \geq c_1 \max_{i \neq 1} Y_i]$$

where Y_1, \dots, Y_k are independent and Y_i is the r -th order statistic from $H_{F[i]}$ for $i = 1, \dots, k$. The right-hand side of (1.2.26) is equal to

$$(1.2.27) \quad \int_{-\infty}^{\infty} \phi\left(\frac{\delta x}{c_1} - a_\alpha h(a_\alpha)\left(1 - \frac{\delta}{c_1}\right)\left(\frac{n}{\alpha\alpha}\right)^{\frac{1}{2}}\right) \phi^{k-2}\left(\frac{x}{c_1} - a_\alpha h(a_\alpha)\left(1 - \frac{1}{c_1}\right)\left(\frac{n}{\alpha\alpha}\right)^{\frac{1}{2}}\right) d\phi(x)$$

where c_1 is the constant used in defining R_1 , a_α is the (unique) α -quantile of $H_{F[k]}(x)$ and $h(x)$ is the density function of $H_{F[k]}(x)$.

From now on, it is assumed that $k = 2$. Therefore (1.2.25) reduces to

$$(1.2.28) \quad E(S|R_1) - P[CS|R_1] = \phi(-h(a_\alpha)a_\alpha\left(1 - \frac{\delta}{c_1}\right)\left(\frac{n}{\alpha\alpha}\right)^{\frac{1}{2}})\left(1 + \frac{\delta^2}{c_1^2}\right)^{-\frac{1}{2}}.$$

$$\text{Let } \int_{-\infty}^{\infty} \phi^{k-1}\left(\frac{x}{c_1} + (1-c_1)\eta_\alpha g(\eta_\alpha)\right) \frac{1}{c_1}\left(\frac{n}{\alpha\alpha}\right)^{\frac{1}{2}} d\phi(x) = P^*.$$

Barlow and Gupta [5] have proved that

$$c_1 = 1 - \frac{2^{\frac{1}{2}}D}{n^{\frac{1}{2}}} + \frac{D}{n} - \frac{3}{2^{3/2}} \frac{D^3}{n^{3/2}} + O\left(\frac{1}{n^2}\right)$$

$$\text{where } D = \frac{\phi^{-1}(P^*) (\alpha\bar{\alpha})^{\frac{1}{2}}}{\eta_\alpha g(\eta_\alpha)}.$$

Now, setting the right hand side of (1.2.28) equal to ϵ and using

$$c_1 \approx 1 - \frac{2\sqrt{\delta}}{n^{\frac{1}{2}}}, \text{ we obtain}$$

$$(1.2.29) \quad n_{R_1}(\epsilon) \approx [-(\alpha\bar{\alpha})^{\frac{1}{2}\Phi^{-1}(\epsilon)}(1+\delta^2)^{\frac{1}{2}+\sqrt{2}} D\delta a_\alpha h(a_\alpha)]^2 \cdot [a_\alpha^2 h^2(a_\alpha)(1-\delta)^2]^{-1}.$$

Comparison with Barlow-Gupta Procedure

Barlow and Gupta [5] propose a procedure R_3 , for a quantile selection problem.

R_3 : Select population π_i if and only if

$$(1.2.30) \quad T_{r,i} \geq c_3 \max_{1 \leq j \leq k} T_{r,j}$$

where c_3 ($0 < c < 1$) is chosen to satisfy $P[CS|R] \geq P^*$ and $T_{r,i}$ is the r -th order statistic from F_i where $r \leq (n+1)\alpha < r+1$. A similar expression for $n_{R_3}(\epsilon)$ (see Barlow and Gupta [5]) is

$$n_{R_3}(\epsilon) \approx [-(\alpha\bar{\alpha})^{\frac{1}{2}\Phi^{-1}(\epsilon)}(1+\delta^2)^{\frac{1}{2}+\sqrt{2}} D\delta \xi_\alpha f(\xi_\alpha)]^2 [\xi_\alpha^2 f^2(\xi_\alpha)(1-\delta)^2]^{-1}$$

where f is the density of F with unique α -quantile, ξ_α .

$$(1.2.31) \quad ARE(R_1, R_3; \delta) = \lim_{\epsilon \rightarrow 0} \frac{n_{R_1}(\epsilon)}{n_{R_3}(\epsilon)} = \frac{\xi_\alpha^2 f^2(\xi_\alpha)}{a_\alpha^2 h^2(a_\alpha)}.$$

If $G(x) = 1 - e^{-x}$, $x > 0$ and $F(x) = 1 - e^{-x}$, $x > 0$, then

$$(1.2.32) \quad ARE(R_1, R_3; \delta) = \frac{(1-\alpha)^2 \log^2(1-\alpha)}{\alpha^2} \leq 1. \\ = 0.4803, \quad \alpha = \frac{1}{2}.$$

Comparison with Gupta Procedure

Gupta [30] gave a selection procedure for gamma populations

π_i 's with densities $\frac{1}{\Gamma(a)\theta_i^a} x^{a-1} e^{-\frac{x}{\theta_i}}$ $x > 0, \theta_i > 0, i = 1, 2, \dots, k$.

The procedure R_4 is

R_4 : Select population π_i if and only if

$$(1.2.33) \quad \bar{X}_i \geq c_4 \max_{1 \leq j \leq k} \bar{X}_j$$

where \bar{X}_i is the sample mean of size n from π_i and c_4 is the largest constant ($0 < c_4 \leq 1$) chosen so that $P[CS|R_4] \geq P^*$.

For $k = 2$, $\theta_{[1]} = \delta$ and $\theta_{[2]} = 1$, Barlow and Gupta [5] have prove that

$$(1.2.34) \quad \begin{aligned} \text{ARE}(R_3, R_4; \delta) &= \lim_{\epsilon \rightarrow 0} \frac{n_{R_3}(\epsilon)}{n_{R_4}(\epsilon)} \\ &= \frac{a(\log \delta)^2 \alpha \bar{\alpha} (1 + \delta^2)}{2(1-\delta)^2 [\epsilon_{\alpha} f(\epsilon_{\alpha})]^2} \end{aligned}$$

It is easy to show that

$$(1.2.35) \quad \begin{aligned} \text{ARE}(R_1, R_4; \delta) &= \text{ARE}(R_1, R_3; \delta) \text{ARE}(R_3, R_4; \delta) \\ &= \left\{ \frac{\sqrt{a} \log \delta \sqrt{\alpha \bar{\alpha}} \sqrt{1 + \delta^2}}{\sqrt{2} (1-\delta) a_{\alpha} h(a_{\alpha})} \right\}^2 \end{aligned}$$

If $G(x) = 1 - e^{-x}$ for $x > 0$ and $a = 1$,

then

$$(1.2.36) \quad \text{ARE}(R_1, R_4; \delta) = \frac{(1-\alpha)(1+\delta^2) \log^2 \delta}{2(1-\delta)^2 \alpha}$$

Therefore as $\delta \rightarrow 1$, we get

$$(1.2.37) \quad \text{ARE}(R_1, R_4; \delta+1) = \frac{1-\alpha}{\alpha}.$$

It is easy to see that

$$(1.2.38) \quad \text{ARE}(R_1, R_4; \delta+1) \begin{cases} < 1, & \alpha > \frac{1}{2} . \\ = 1, & \alpha = \frac{1}{2} . \\ > 1, & \alpha < \frac{1}{2} . \end{cases}$$

Comparison of R_1 and R_5 from uniform distribution

Suppose π_1 and π_2 are two independent uniform populations with distribution functions F_i ($i = 1, 2$).

$$F_i(x) = \begin{cases} 0 & x < 0 \\ \frac{x}{\theta_i} & 0 \leq x \leq \theta_i \\ 1 & x > \theta_i \end{cases}$$

where $\delta = \theta_{[1]} < \theta_{[2]} = 1$.

A sample size n is drawn from each of the two populations. Let T_i^* be the total life statistic until r -th failure from π_i ($i = 1, 2$) where $r \leq (n+1)\alpha < r+1$.

The procedure R_5 is given by

R_5 : Select population π_i if and only if

$$(1.2.39) \quad T_i^* \geq c_5 \max_{1 \leq j \leq k} T_j^*$$

where c_5 is chosen so that $P[CS|R_5] \geq P^*$. Let $T_{(i)}^*$ be associated with $\theta_{[i]}$.

$$(1.2.40) \quad E(S|R_5) - P[CS|R_5] = P[T_{(1)}^* \geq c_5 T_{(2)}^*] = P[T_1' \geq \frac{c_5}{\delta} T_2']$$

where T_1', T_2' are two independent total life statistic until r -th failure from uniform distribution over $(0,1)$. By Gupta and Sobel [42],

$$(1.2.41) \quad \frac{T_1' - u}{\sigma} \rightarrow N(0,1) \quad \text{as } n \rightarrow \infty,$$

$$\text{where } u = \frac{n\alpha(2n-\alpha n+1)}{2n+1} \approx u' = \frac{n\alpha(2-\alpha)}{2}$$

$$\sigma^2 = \frac{\alpha(1-\alpha)n^2(2-\alpha)^2n^2}{4n^3} + \frac{\alpha^3n^3}{12n^2}$$

$$= A n \quad \text{where } A = \frac{\alpha(1-\alpha)(2-\alpha)^2}{4} + \frac{\alpha^3}{12}.$$

$$\text{Hence } \frac{u}{\sigma} \approx \frac{u'}{\sigma} = B\sqrt{n} \quad \text{where } B = \frac{\alpha(2-\alpha)}{2\sqrt{A}}.$$

From (1.2.40), we have

$$\begin{aligned} E(S|R_5) - P[CS|R_5] &= P\left[\frac{T_1' - u'}{\sigma} \geq \frac{c_5}{\delta} \left(\frac{T_2' - u'}{\sigma}\right) + \left(\frac{c_5}{\delta} - 1\right) \frac{u'}{\sigma}\right] \\ &\approx P\left[Z_1 \geq \frac{c_5}{\delta} Z_2 + \left(\frac{c_5}{\delta} - 1\right) B\sqrt{n}\right] \end{aligned}$$

where Z_1, Z_2 are i.i.d. with $N(0,1)$.

Hence

$$E(S|R_5) - P[CS|R_5] = \int_{-\infty}^{\infty} \Phi\left[\frac{\delta}{c_5} x - \left(1 - \frac{\delta}{c_5}\right) B\sqrt{n}\right] d\Phi(x)$$

$$= \phi\left[\frac{-(1 - \frac{\delta}{c_5})B\sqrt{n}}{\sqrt{1 + (\frac{\delta}{c_5})^2}}\right].$$

Let $E(S|R_5) - P[CS|R_5] = \epsilon > 0$, we obtain

$$(1.2.42) \quad \left(\frac{1}{c_5} - \frac{1}{\delta}\right) \sqrt{n} = \sqrt{\frac{1}{\delta^2} + \frac{1}{c_5^2}} \cdot \frac{\phi^{-1}(\epsilon)}{B}$$

Note that

$$\inf_{\Omega} P[CS|R_5] = P[T_1' \geq c_5 T_2']$$

where T_1' , T_2' are defined as above.

$$\begin{aligned} P[T_1' \geq cT_2'] &= P\left[\frac{T_1' - u'}{\sigma} \geq c_5 \left[\frac{T_2' - u'}{\sigma}\right] + (c_5 - 1) \frac{u'}{\sigma}\right] \\ &\approx P[Z_1 \geq c_5 Z_2 + (c_5 - 1)B\sqrt{n}] \end{aligned}$$

where Z_1 , Z_2 are i.i.d. with $N(0,1)$.

Hence

$$\begin{aligned} P[T_1' \geq cT_2'] &= \int_{-\infty}^{\infty} \phi\left(\frac{1}{c_5} x - (1 - \frac{1}{c_5})B\sqrt{n}\right) d\phi(x) \\ &= \phi\left[\frac{-(1 - \frac{1}{c_5})B\sqrt{n}}{\sqrt{1 + \frac{1}{c_5^2}}}\right] \end{aligned}$$

Setting $\inf_{\Omega} P[CS|R_5] = P^*$, we obtain

$$\begin{aligned} -(1 - \frac{1}{c_5})B\sqrt{n} &= \phi^{-1}(P^*) \sqrt{1 + \frac{1}{c_5^2}} \\ (1 - c_5)\sqrt{n} &= D \sqrt{1 + c_5^2} \quad \text{where } D = \frac{\phi^{-1}(P^*)}{B}. \end{aligned}$$

We see that $c_5 \approx 1 - \frac{\sqrt{2} D}{\sqrt{n}}$,

$$\frac{1}{c_5} \approx 1 + \frac{\sqrt{2} D}{\sqrt{n}}.$$

From (1.2.42),

$$\begin{aligned} \left(1 + \frac{\sqrt{2D}}{\sqrt{n}} - \frac{1}{\delta}\right)\sqrt{n} &= \frac{\phi^{-1}(\epsilon)}{B} \left\{ \frac{1}{\delta^2} + \left[1 + \frac{2\sqrt{2}D}{\sqrt{n}} + \frac{2D^2}{n}\right]^{\frac{1}{2}} \right\} \\ \sqrt{n}\left(1 - \frac{1}{\delta}\right) + \sqrt{2}D &\approx \frac{\phi^{-1}(\epsilon)}{B} \left\{ \frac{1}{\delta^2} + 1 \right\}^{\frac{1}{2}} \end{aligned}$$

Thus

$$(1.2.43) \quad n_{R_5}(\epsilon) \approx \left\{ \frac{\phi^{-1}(\epsilon)\sqrt{1+\delta^2} - \sqrt{2}\delta\phi^{-1}(P^*)}{B(1-\delta)} \right\}^2.$$

If we assume that $G(x) = x$ for $0 < x < 1$, then the $n_{R_1}(\epsilon)$ is given by (1.2.29) with $a_\alpha = \alpha$, $h(a_\alpha) = 1$, $\eta_\alpha = \alpha$ and $g(\eta_\alpha) = 1$.

Hence

$$\begin{aligned} (1.2.44) \quad \text{ARE}(R_1, R_5; \delta) &= \lim_{\epsilon \rightarrow 0} \frac{n_{R_1}(\epsilon)}{n_{R_5}(\epsilon)} \\ &= \frac{B^2(1-\alpha)}{\alpha} = \frac{3(1-\alpha)(2-\alpha)^2}{3(1-\alpha)(2-\alpha)^2 + \alpha^2} < 1 \\ &= 0.931, \quad \alpha = \frac{1}{2}. \\ &= 0.675, \quad \alpha = \frac{3}{4}. \\ &= 0.574, \quad \alpha = \frac{4}{5}. \end{aligned}$$

(C) Selection procedure for distribution \prec ordered with respect to Weibull distribution
C

Assumed that the specified distribution $G(x)$ is a Weibull distribution. In other words, $G(x)$ is given by

$$G(x) = \begin{cases} 1 - e^{-\lambda x^\alpha} & \text{for } x \geq 0 \\ 0 & \text{for } x < 0 \end{cases}$$

where $\lambda > 0$ and attention is restricted to $\alpha \geq 1$ which is assumed known.

In this case, we use the statistic T_i^* which is defined by

$$(1.2.45) \quad T_i^* = \sum_{j=1}^r X_{i;j,n}^\alpha + (n-r)X_{i;r,n}^\alpha, \quad i = 1, \dots, k,$$

(as before, $X_{i;j,n}$ denote the j -th order statistic from F_i , $i=1, \dots, k$).

Since $G(x)$ is convex with respect to the exponential distribution if $\alpha \geq 1$ and since the convex ordering is transitive, the family of distributions which are convex with respect to Weibull ($\alpha \geq 1$) will have IFR distribution. Thus our interest here is in a special subclass of IFR distributions.

The rule for selecting the population which is associated with $F_{[k]}$ is as follows,

R_6 : Select population π_i if and only if

$$(1.2.46) \quad T_i^* \geq c_6 \max_{1 \leq j \leq k} T_j^*$$

where c_6 ($0 < c_6 \leq 1$) is determined so as to satisfy the basic probability requirement.

Using the fact that if $F <_c G$ and $F(0) = G(0) = 0$

then

$$F_\alpha <_c G_\alpha \quad \text{for } \alpha \geq 1,$$

where F_α is the c.d.f. of X^α , $F(x)$ is the c.d.f. of X , G_α is the c.d.f. of Y^α and $G(y)$ is the c.d.f. of Y .

Also,

$G_\alpha^{-1} F_\alpha(X_{i;n}^\alpha) =_{st}$ i -th order statistic from $G^*(x) = 1 - e^{-\lambda x}$, $x \geq 0$ where $X_{1,n} \leq \dots \leq X_{n,n}$ are order statistics from F . In a

manner similar to the proof of Theorem 1.2.1, one can prove the following theorem.

Theorem 1.2.3. If $F_i(x) \geq F_{[k]}(x) \quad \forall x$ and $i = 1, \dots, k$,

$$F_{[k]}(0) = 0, F_{[k]} \prec_c G, G(x) = 1 - e^{-\lambda x^\alpha}, x > 0, \lambda > 0 \text{ and } \alpha (\geq 1)$$

is known. If $n \geq \max\{r, \frac{r-1}{1-c_6}\}$, then

$$(1.2.47) \inf_{\Omega} P[CS | R_6] = \int_0^{\infty} G_T^{k-1} \left(\frac{x}{c_6} \right) dG_T(x)$$

where $G_T(x)$ is the c.d.f. of a χ^2 random variable with $2r$ d.f.

(D) Selection with respect to the means for Gamma populations

Let π_1, \dots, π_k be k populations with densities

$$r_i(x) = \frac{\beta^{\alpha_i}}{\Gamma(\alpha_i)} x^{\alpha_i-1} e^{-\beta x}, x \geq 0, \beta > 0, \alpha_i \geq 1, i = 1, \dots, k.$$

Let $R_i(x)$ be the distribution function of π_i , $i = 1, \dots, k$. We are given a sample of size n from each π_i . Let T_i^* be total life statistic until r -th failure from π_i .

Let $\alpha_{[1]} \leq \dots \leq \alpha_{[k]}$ be the ordered values of α_i 's. We are interested in selecting the population with the largest value $\alpha_{[k]}$ (unknown). Since the mean of π_i is $\frac{\alpha_i}{\beta}$, selection of the population with largest mean is equivalent to selecting the population with largest value, $\alpha_{[k]}$. The subset selection rule based on T_i is:

R_7 : Select population π_i if and only if

$$(1.2.48) \quad T_i^* \geq c_7 \max_{1 \leq j \leq k} T_j^*,$$

where c_7 ($0 < c_7 < 1$) is chosen to satisfy

$$P[CS|R_7] \geq P^*.$$

Since the rule R_7 is scale invariant, we can assume $\beta = 1$.

Case 1: All α_i are unknown and ≥ 1 . Let $\Omega_1 = \{\underline{\alpha} = (\alpha_1, \dots, \alpha_k) : \alpha_i \geq 1 \forall i\}$

$$R_i \prec G(x) = 1 - e^{-x}, \quad x > 0, \quad i = 1, \dots, k.$$

In this case, by Corollary 1.2.2, we have the following result.

$$\text{If } n \geq \max\{r, \frac{r-1}{1-c}\}, \text{ then } \inf_{\Omega_1} P[CS|R_7] = \int_0^{\infty} H^{k-1}\left(\frac{x}{c}\right) dH(x),$$

where $H(x)$ is the c.d.f. of a χ^2 r.v. with $2r$ d.f.

Case 2: α_i are unknown but assume $1 \leq \alpha_i \leq \Delta$, $i = 1, \dots, k$ and Δ is known.

Let $R_{\Delta}(x)$ be the c.d.f. of X with density function

$$r_{\Delta}(x) = \frac{\beta^{\Delta}}{\Gamma(\Delta)} x^{\Delta-1} e^{-\beta x}, \quad x > 0, \quad \beta > 0.$$

Let $h(x)$ be the density function of a χ^2 r.v. with $2r$ d.f.

Theorem 1.2.4.

$$(1.2.49) \quad P[CS|R_7] \geq \int_0^{\infty} H^{k-1}\left(\frac{2n}{c_7} x\right) \frac{2nh(2ny)}{r_{\Delta}(y)} e^{-x} dx,$$

where $y = R_{\Delta}^{-1}(1 - e^{-x})$.

$$\text{Proof.} \quad P[CS|R_7] = P[T_{(k)}^* \geq c_7 \max_{1 \leq j \leq k-1} T_{(j)}^*],$$

where $T_{(i)}^*$ is associated with $\alpha_{[i]}$, $i = 1, \dots, k$.

Since $R_{\Delta}(x) \leq R_j(x) \leq G(x) = 1 - e^{-x}$, by Lemma 1.2.6, then

$$(1.2.50) \quad P[CS|R_7] \geq P[T_k^{**} \geq c_7 \max_{1 \leq j \leq k-1} T_j^{**}]$$

where T_k^{**} is the total life statistic until r -th failure from $G(x)$ and T_j^{**} ($j = 1, \dots, k-1$) is the total life statistic until r -th failure from $R_{\Delta}(x)$.

Since $\Delta \geq 1$ then $R_{\Delta} \prec_c G$.

Let $\varphi(x) = G^{-1}R_{\Delta}(x)$.

$$(1.2.51) \quad P[T_k^{**} \geq c_7 T_j^{**}, j=1, \dots, k-1] = P[\varphi(\frac{1}{n} T_k^{**}) \geq \varphi(\frac{c_7}{n} T_j^{**}), j=1, \dots, k-1]$$

By Lemma 1.2.4 (a) with $a_1 = \dots = a_{r-1} = \frac{c_7}{n}$, $a_r = \frac{(n-r+1)c_7}{n}$

$$a_i = 0 \quad \text{for } i \geq r+1$$

and $\varphi(x) \stackrel{st}{=} Y$ where X is a r.v. with distribution R_{Δ}

Y is a r.v. with distribution G ,

we have

$$(1.2.52) \quad P[\varphi(\frac{1}{n} T_k^{**}) \geq \varphi(\frac{c_7}{n} T_j^{**}), j=1, \dots, k-1] \geq P[\varphi(\frac{1}{n} T_k^{**}) \geq \frac{c_7}{2n} Y_j, \\ j = 1, \dots, k-1]$$

where Y_j ($j = 1, \dots, k-1$) is a r.v. with χ^2 with $2r$ d.f. From

(1.2.50), (1.2.51) and (1.2.52), we have

$$P[CS|R_7] \geq \int_0^{\infty} H^{k-1}(\frac{2n}{c_7} x) dB(x),$$

where $B(x) = P[\varphi(\frac{1}{n} T_k^{**}) \leq x]$.

Since $\varphi(x) = -\ln(1 - R_{\Delta}(x))$, then $\varphi^{-1}(x) = R_{\Delta}^{-1}(1 - e^{-x})$.

Thus

$$\begin{aligned} B(x) &= P[T_k^{**} \leq n \varphi^{-1}(x)] = H(2n \varphi^{-1}(x)) \\ &= H[2n R_{\Delta}^{-1}(1-e^{-x})]. \end{aligned}$$

Now

$$\begin{aligned} \frac{dB(x)}{dx} &= h[2ny] \frac{d(2ny)}{dx} \\ &= h(2ny)2n \cdot \frac{e^{-x}}{r_{\Delta}(y)} \quad \text{where } y = R_{\Delta}^{-1}(1-e^{-x}). \end{aligned}$$

So that,

$$\int_0^{\infty} H^{k-1}\left(\frac{2n}{c_7} x\right) dB(x) = \int_0^{\infty} H^{k-1}\left(\frac{2n}{c_7} x\right) \frac{2nh(2ny)}{r_{\Delta}(y)} e^{-x} dx.$$

This completes the proof.

Let S denote the size of the selected subset. The expected value of S when R_7 is used is given by

$$(1.2.53) \quad E(S|R_7) = \sum_{i=1}^k P[T_i^* \geq c_7 \max_{1 \leq j \leq k} T_j^*].$$

Let $\Omega' = \{\underline{\alpha} = (\alpha_1, \dots, \alpha_k) : 1 \leq \alpha_i \leq \Delta, i = 1, \dots, k\}$.

For $\underline{\alpha} \in \Omega'$, since $R_{\Delta}(x) \leq R_i(x) \leq G(x) = 1-e^{-x}$, then

$$E(S|R_7) \leq \sum_{i=1}^k P[T_1^{**} \geq c_7 \max_{2 \leq j \leq k} T_j^{**}]$$

where T_1^{**} is the total life statistic until r -th failure from $R_{\Delta}(x)$ and T_j^{**} ($j = 2, \dots, k$) is the total life statistic until r -th failure from $G(x)$.

$$\text{Hence } E(S|R_7) \leq k P[T_1^{**} \geq c_7 \max_{2 \leq j \leq k} T_j^{**}]$$

so that,

$$(1.2.54) \quad \sup_{\Omega} E(S|R_7) = k \int R^{k-1} \left(\frac{x}{c_7}\right) dS(x)$$

where $R(x)$ ($S(x)$) is the c.d.f. of total life statistic until r -th failure from $G(x)$ ($R_{\Delta}(x)$).

1.3. Selecting a best population - using indifference zone approach.

Let π_1, \dots, π_k be k populations. The random variable X_i associated with π_i has an absolutely continuous distribution F_i . We assume there exists a $F_{[k]}(x)$ such that $F_{[i]}(x) \geq F_{[k]}(\frac{x}{\delta})$ for all x , $i=1, \dots, k-1$ and δ ($0 < \delta \leq 1$) is specified. Let

$$(1.3.1) \quad \Omega = \{ \underline{F} = (F_1, \dots, F_k) : \exists a_j \text{ such that}$$

$$F_i(x) \geq F_j(\frac{x}{\delta}) \quad \forall i \neq j \} .$$

The correct selection is the choice of any population which is associated with $F_{[k]}$. We propose the selection rule R_g : Select population π_i if and only if

$$(1.3.2) \quad T_i = \max_{1 \leq j \leq k} T_j \quad \text{where } T_i \text{ is defined as in Section 1.2.}$$

We want the $P[CS|R_g] \geq P^*$, for all $\underline{F} \in \Omega$, where P^* ($\frac{1}{k} < P^* < 1$) is specified.

Theorem 1.3.1. If $F_i, G \in \mathfrak{B}$, $i = 1, \dots, k$, $F_{[k]}(0) = 0$, $F_{[k]} <_c G$ and $G^{-1}(0) \leq 0$. If $a_j \geq 0$, $j = 1, \dots, r$,

$$gG^{-1}(0) \leq 1 \text{ and } a_r \geq \delta,$$

then

$$(1.3.3) \quad P[CS|R_\delta] \geq \int_0^\infty G_T^{k-1}\left(\frac{x}{\delta}\right) dG_T(x)$$

where $G_T(x)$ is the c.d.f. of T .

Proof. $P[CS|R_\delta] = P[T_{(k)} \geq \max_{1 \leq j \leq k} T_{(j)}]$.

Since $F_{[i]}(\delta x) \geq F_{[k]}(x)$, $i=1, \dots, k-1$, it follows from Lemma 1.2.6 that

$$\begin{aligned} P[CS|R_\delta] &= P[T_{(k)} \geq \delta \frac{T_{(j)}}{\delta} \quad \forall j \neq k] \\ &\geq P[T_{(k)} \geq \delta T_j^* \quad \forall j \neq k] \end{aligned}$$

where $T_1^*, \dots, T_{j-1}^*, T_{(k)}$ are i.i.d. with c.d.f. $W_k(x)$. Using the same argument as in Theorem 1.2.1, we have our theorem.

Remark 1.3.1. $\inf_{\Omega} P[CS|R] = \int_{G^{-1}(0)}^\infty G_T^{k-1}\left(\frac{x}{\delta}\right) dG_T(x)$ if $G^{-1}(0) = 0$.

For given k , δ , P^* and $G(x)$, we can possibly find the values of the pair (n, r) , $(n \geq r)$ which satisfy

$$(1.3.4) \quad a_r \geq \delta \text{ and } \int_{G^{-1}(0)}^\infty G_T^{k-1}\left(\frac{x}{\delta}\right) dG_T(x) \geq P^*.$$

If $G(x) = x$ for $0 < x < 1$, we can always find the values of the pair (n, r) , $(n \geq r)$ which satisfy

$$\int_0^\infty G_T^{k-1}\left(\frac{x}{\delta}\right) dG_T(x) \geq P^*.$$

In this case, $G_T(x)$ is the c.d.f. of r -th order statistic based on a sample of size n from G .

If $G(x) = 1 - e^{-x}$ for $x \geq 0$ (see Patel [58]), we can find the smallest integer r which satisfies

$\int_0^{\infty} G_T^{k-1}(\frac{x}{\delta}) dG_T(x) \geq P^*$ where $G_T(x)$ is the c.d.f. of a χ^2 random variable with $2r$ d.f. Since $\frac{1}{\delta} a_r \geq 1$ iff $n \geq \frac{r-1}{1-\delta}$, the minimum n satisfies $n \geq \max\{r, \frac{r-1}{1-\delta}\}$.

1.4. Selection procedure for distribution \prec^* ordered with respect to G .

Let $t(> 0)$ be a given number. Let $N_i(t)$ be the number of failures in time t among the n units on life test from π_i which has a continuous distribution F_i , $i = 1, \dots, k$. We assume that

$$(1.4.1) \quad N_i(t) \geq 1, \quad i = 1, \dots, k,$$

and there exists a $F_{[k]}(x)$ such that $F_i(x) \geq F_{[k]}(x)$ for all x , $i = 1, \dots, k$. The correct selection is the choice of any population which is associated with $F_{[k]}$. Let T_i be the total life statistic until $N_i(t)$ -th failure from population π_i . Let $T_{(i)}$ be associated with $F_{[i]}$. We propose the rule

$$(1.4.2) \quad R_g: \text{Select } \pi_i \Leftrightarrow T_i \geq c_g \max_{1 \leq j \leq k} T_j$$

where $c_g (0 < c_g \leq 1)$ is chosen so that $P[CS|R_g] \geq P^*$.

Theorem 1.4.1. If $F_i(x) \geq F_{[k]}(x) \quad \forall x, \quad i = 1, \dots, k$ and $F_{[k]} \prec^* G$, then

$$(1.4.3) \quad P[CS|R_g] \geq \int_0^{\infty} A_1^{k-1}(\frac{x}{c_g}) dA_2(x)$$

where $A_1(x)$ ($A_2(x)$) is the c.d.f. of total life statistic until n -th (first) failure from G .

Proof. From (1.4.1), we have

$$\begin{aligned} P[CS|R_9] &= P[T_{(k)} \geq c_9 \max T_{(j)}] \\ &\geq P[T_k^* \geq c_9 T_j^* \text{ for } j = 1, \dots, k-1] \end{aligned}$$

where $\frac{T_i^*}{n}$ is the mean of a sample of size n from $F_{[i]}$, $i=1, \dots, k-1$

and $\frac{T_k^*}{n}$ is the first order statistic from $F_{[k]}$.

Let

$$(1.4.4) \quad T_j^{**} = \sum_{\ell=1}^n X_{j\ell} \quad \text{for } j = 1, \dots, k-1.$$

where X_{j1}, \dots, X_{jn} are i.i.d. from $F_{[j]}$, $j = 1, \dots, k-1$.

since $F_{[j]}(x) \geq F_{[k]}(x)$ then $T_j^{**} \geq T_j^*$. Hence

$$(1.4.5) \quad P[T_k^* \geq c_9 T_j^*, j \neq k] \geq P[T_k^* \geq c_9 T_j^{**}, j \neq k].$$

Let $\varphi(x) = G^{-1}F_{[k]}(x)$.

$$(1.4.6) \quad P[T_k^* \geq c_9 T_j^{**}, j \neq k] = P[\varphi(\frac{T_k^*}{nc_9}) \geq \varphi(\frac{1}{n} T_j^{**}), j \neq k]$$

Since φ is starshaped, then

$$(1.4.7) \quad \varphi(\frac{1}{c_9} \frac{T_k^*}{n}) \geq \frac{1}{c_9} \varphi(\frac{T_k^*}{n}).$$

By Lemma 1.2.5 (a) and (1.4.4),

$$(1.4.8) \quad \varphi(\frac{1}{n} T_j^{**}) \leq \sum_{\ell=1}^n \frac{1}{n} \varphi(X_{j\ell}).$$

Note that $\varphi(\frac{T_k^*}{n}) \stackrel{st}{=} Y_k$ where Y_k is the first order statistic of size n from G and $\varphi(X_{j\ell})$ has distribution G , $j = 1, \dots, k-1$, $\ell = 1, \dots, n$.

Let $Y_j = \sum_{\ell=1}^n \varphi(X_{j\ell})$, $j = 1, \dots, k-1$.

From (1.4.7) and (1.4.8), the right-hand side of (1.4.6) is greater than or equal to

$$\begin{aligned} P\left[\frac{1}{c_9} Y_k \geq \frac{1}{n} Y_j, j=1, \dots, k-1\right] &= P\left[\frac{1}{c_9}(nY_k) \geq Y_j, j=1, \dots, k-1\right] \\ &= \int_0^{\infty} A_1^{k-1}\left(\frac{x}{c_9}\right) dA_2(x). \end{aligned}$$

The proof is complete.

1.5. Selection with respect to the means for distributions s-ordered with respect to a specified distribution G.

Let the random variable X_i associated with π_i have an absolutely continuous distribution F_i . Assume that $F_i(x) = F\left(\frac{x}{\theta_i}\right)$, $i=1, \dots, k$, $F < G$ and F and G are symmetric about 0. As before, let

$0 < \theta_{[1]} \leq \dots \leq \theta_{[k]}$ be ordered values of unknown θ_i ($i = 1, \dots, k$).

A best population is defined as the one having $\theta_{[k]}$. We are given a sample of size n from each of the k populations. Assume that n is large. Let

$$(1.5.1) \quad T_i = -r'X_{i;r',n} - X_{i;r'+1,n} - \dots - X_{i;l_i,n} + X_{i;l_i+1,n} + \dots + (n-r+1)X_{i;r,n}$$

where $X_{i;1,n} \leq \dots \leq X_{i;l_i,n} < 0 \leq X_{i;l_i+1,n} \leq \dots \leq X_{i;n,n}$ are order statistics from $F_i(x)$, and r', r are given integers such that $1 \leq r' \leq l_i$, $n \geq r > l_i$, $i = 1, \dots, k$.

Similarly, let

$$(1.5.2) \quad T = -r'Y_{r',n} - Y_{r'+1,n} - \dots - Y_{\ell,n} + Y_{\ell+1,n} + \dots + (n-r+1)Y_{r,n}$$

where $Y_{1,n} \leq \dots \leq Y_{\ell,n} < 0 < Y_{\ell+1,n} \leq \dots \leq Y_{n,n}$ are order statistics from G . Then, for selecting a subset containing $\theta_{[k]}$, we propose the following selection rule:

R_{10} : Select population π_i if and only if

$$(1.5.3) \quad T_i \geq c_{10} \max_{1 \leq j \leq k} T_j$$

where c_{10} ($0 < c_{10} < 1$) is determined as to satisfy the probability requirement, $\inf_{\Omega} P[CS|R_{10}] \geq P^*$, where CS stands for a correct selection, i.e., the selection of any subset which contains the population with $\theta_{[k]}$ and $\Omega = \{(\theta_1, \dots, \theta_k) : \theta_i > 0, i=1, \dots, k\}$.

Let $\pi(i)$ be associated with $\theta_{[i]}$, $i = 1, \dots, k$. Now, we give a theorem pertaining to the computation of the minimum probability of a correct selection.

Theorem 1.5.1. If $F_i(x) = F\left(\frac{x}{\theta_i}\right) \forall x$ and $i = 1, \dots, k$, $F \prec_s G$ and

F and G are symmetric about 0.

If $\lim_{n \rightarrow \infty} \frac{r'}{n} = t'$, $\lim_{n \rightarrow \infty} \frac{r}{n} = t$ and $c_{10} \leq \min\{t', 1-t\}$,

then,

$$(1.5.4) \quad \lim_{n \rightarrow \infty} \inf_{\Omega} P[CS|R_{10}] = \int_0^{\infty} H^{k-1}\left(\frac{x}{c_{10}}\right) dH(x)$$

where $H(x)$ is the c.d.f. of T defined in 1.5.2.

Proof. $P[CS|R_{10}] = P[T_{(k)} \geq c_{10}T_{(j)}, j=1, \dots, k-1]$

where $T_{(j)} = -r'X_{(i)r',n} - X_{(i)r'+1,n} - \dots - X_{(i)\ell(i),n} + X_{(i)\ell(i)+1,n} + \dots + (n-r+1)X_{(i)r,n}$

and $X_{(i),1,n} \leq \dots \leq X_{(i)\ell(i),n} < 0 < X_{(i)\ell(i)+1,n} < \dots < X_{(i)n,n}$ are order statistics from $\pi(i)$.

It is easy to see that

$$(1.5.5) \quad \inf_{\Omega} P[CS|R_{10}] = P[T'_k \geq c_{10} T'_j, j=1, \dots, k-1]$$

where $T'_i = -r'Y_{i;r',n} - \dots - Y_{i;\ell(i),n} + Y_{i;\ell(i)+1,n} + \dots + (n-r+1)Y_{i;r,n}$

and $Y_{i;1,n} \leq \dots \leq Y_{i;\ell(i),n} < 0 < Y_{i;\ell(i)+1,n} \leq \dots \leq Y_{i,n,n}$ are order statistics from $F(x)$.

Let $\phi(x) = G^{-1}F(x)$.

$$(1.5.6) \quad P[T'_k \geq c_{10} T'_j] = P[\phi(\frac{1}{c_{10}n} T'_k) \geq \phi(\frac{1}{n} T'_j), j=1, \dots, k-1].$$

By Corollary 2.9 of Lawrence [48] with $i_0 = \ell(i)$, $j_0 = \ell(i)+1$,

and

$$a_j = \begin{cases} 0 & 1 \leq j < r' \\ -\frac{r'}{n} & j = r' \\ -\frac{1}{n} & r' < j \leq \ell(i) \\ \frac{1}{n} & \ell(i) < j < r \\ \frac{n-r+1}{n} & j = r \\ 0 & n \geq j > r \end{cases},$$

then

$$(1.5.7) \quad \phi(\frac{1}{n} T'_i) \leq \frac{-r'}{n} \phi(Y_{i;r',n}) - \dots - \frac{1}{n} \phi(Y_{i;\ell(i),n}) \\ + \frac{1}{n} \phi(Y_{i;\ell(i)+1,n}) + \dots + \frac{n-r+1}{n} \phi(Y_{i;r,n}), i=1, \dots, k-1.$$

Also by Corollary 2.10 of Lawrence [48] with $i_0 = r'-1$, $j_0 = n+1$,

$$a_j = \begin{cases} 0 & 0 \leq j < r' \\ \frac{-r'}{nc_{10}} & j = r' \\ \frac{-1}{nc_{10}} & r' < j \leq \ell(k) \\ \frac{1}{nc_{10}} & \ell(k) < j < r \\ \frac{n-r+1}{nc_{10}} & j = r \\ 0 & j > r \end{cases}$$

$$\frac{-r'}{nc_{10}} \leq -1 \text{ and } \frac{n-r+1}{nc_{10}} \geq 1,$$

then

$$(1.5.8) \quad \phi\left(\frac{1}{nc_{10}} T'_k\right) \geq \frac{-r'}{nc_{10}} \phi(Y_{k;r',n}) - \dots - \frac{1}{nc_{10}} \phi(Y_{k;\ell(k),n}) \\ + \frac{1}{nc_{10}} \phi(Y_{k;\ell(k)+1,n}) + \dots + \frac{n-r+1}{nc_{10}} \phi(Y_{k;r,n}).$$

Now $\phi(Y_{j;i,n}) \stackrel{\text{st}}{=} V_{j;i,n}$, $j = 1, \dots, k$, $i = 1, \dots, n$, where $V_{j;i,n}$ is the i -th order statistic from $G(x)$, $j = 1, \dots, k$.

Let

$$T''_i = -r'V_{i;r',n} - \dots - V_{i;\ell(i),n} + V_{i;\ell(i)+1,n} + \dots \\ + (n-r+1)V_{i;r,n}, \quad i = 1, \dots, k.$$

From (1.5.6), (1.5.7) and (1.5.8), we obtain

$$P[T'_k \geq c_{10} T'_j, j \neq k] \geq P[T''_k \geq c_{10} T''_j, j = 1, \dots, k-1].$$

Since G is symmetric about 0, as n becomes large,

$$T''_j \stackrel{\text{st}}{=} T, \quad j = 1, \dots, k.$$

Thus

$$P[CS|R_{10}] \geq \int_0^{\infty} H^{k-1} \left(\frac{x}{c_{10}} \right) dH(x) \text{ if } c_{10} \leq \min\left\{ \frac{r'}{n}, \frac{n-r'+1}{n} \right\}.$$

This proves the theorem.

Corollary 1.5.1. If $F_i(x) = F\left(\frac{x}{\theta_i}\right) \forall x$ and $i = 1, \dots, k$, $F < G$, F is symmetric about 0 and $g(x) = \frac{dG(x)}{dx} = \frac{1}{2} e^{-|x|}$, $-\infty < x < \infty$.

If $\lim_{n \rightarrow \infty} \frac{r'}{n} = t' < \frac{1}{2}$, $\lim_{n \rightarrow \infty} \frac{r}{n} = t > \frac{1}{2}$ and $c_{10} \leq \min\{t', 1-t\}$, then, as n is large,

$$(1.5.9) \quad \inf_{\Omega} P[CS|R_{10}] \approx \int_{-\infty}^{\infty} \phi^{k-1} \left[\frac{x}{c_{10}} + \left(\frac{1-c_{10}}{c_{10}} \right) \sqrt{r-r'+1} \right] d\phi(x)$$

Proof. Considering

$$T^* = \sum_{i=r'}^{\ell-1} -i(Y_{i,n} - Y_{i+1,n}) + \sum_{\ell+1}^r (n-i+1)(Y_{i,n} - Y_{i-1,n})$$

where $Y_{1,n} \leq \dots \leq Y_{\ell,n} < 0 < Y_{\ell+1,n} < \dots < Y_{n,n}$ are order statistics from $G(x)$.

Since $2 T^*$ is distributed as a χ^2 r.v. with $2(r-r'+1)$ d.f. (see Lawrence [47]) and T^* has same distribution as T when n is large, then $2 T$ is distributed as χ^2 with $2(r-r'+1)$ d.f.

By Theorem 1.5.1,

$$P[CS|R_{10}] \geq P[V_k \geq c_{10} \max_{1 \leq j \leq k} V_j]$$

where V_1, \dots, V_k are i.i.d. from χ^2 with $2(r-r'+1)$ d.f.

Since $V_i \sim N(2(r-r'+1), 4(r-r'+1))$, then

$$P[V_k \geq c \max_{1 \leq j \leq k} V_j] \approx \int_{-\infty}^{\infty} \phi^{k-1} \left(\frac{x}{c_{10}} + \frac{1-c_{10}}{c_{10}} \sqrt{r-r'+1} \right) d\phi(x)$$

This proves the corollary.

Assume $X_{1,n} \leq \dots \leq X_{k,n} \leq 0 \leq X_{k+1,n} \leq \dots \leq X_{n,n}$ are the order statistics from distribution G where $\frac{d}{dx} G(x) = g(x) = \frac{1}{2} e^{-|x|}$, $-\infty < x < \infty$.

Define

$$(1.5.10) \quad V_i = \frac{\sum_{j=1}^{i-1} H_n^{-1}\left(\frac{j}{n}\right)}{H_n^{-1}\left(\frac{i}{n}\right)}, \quad i = 1, \dots, n,$$

where $H_n^{-1}\left(\frac{i}{n}\right) = \sum_{j=1}^i gG^{-1}\left(\frac{j-1}{n}\right) (X_{j,n} - X_{j-1,n})$.

Now we want to get the limiting distribution of V_n (as n is large).

Let $T_n = \sqrt{n}\left[\frac{1}{n} V_n - \frac{1}{2}\right]$. To obtain the asymptotic distribution of T_n ,

we use the following result. Let $X_{1,n} < \dots < X_{n,n}$ be the order statistics

from G and $S_n = \frac{1}{n} \sum_{i=1}^n L\left(\frac{i}{n}\right) X_{i,n}$.

$$\sigma^2 = \sigma^2(G) = 2 \int \int_{s < t} L(G(s))L(G(t))G(s)[1-G(t)]dsdt.$$

where L satisfies the condition (ii) of following lemma.

Lemma 1.5.1. (Moore [53])

If $\sigma^2 < \infty$ and

$$(i) \quad E|X| = \int_0^1 |G^{-1}(u)| du < \infty$$

(ii) L is continuous on $[0,1]$ except for jump discontinuities at a_1, \dots, a_M , and L' is continuous and of bounded variation on $[0,1] - \{a_1, \dots, a_M\}$,

then

$$(1.5.12) \quad \mathcal{L}\left\{\sqrt{n}\left[S_n - \int_{-\infty}^{\infty} xL(G(x))dG(x)\right]\right\} \rightarrow N(0, \sigma^2).$$

Let

$$(1.5.11) \quad S_n = \frac{1}{n} \sum_{i=1}^{n-1} H_n^{-1}\left(\frac{i}{n}\right) - \frac{1}{2} H_n^{-1}(1).$$

Then using an argument similar to the one in Barlow and Doksum [3]

$$S_n = \frac{1}{n} \sum_{i=1}^n L\left(\frac{i}{n}\right) X_{i,n}$$

$$\text{where } L\left(\frac{i}{n}\right) = g G^{-1}\left(\frac{i-1}{n}\right) - n\left(\frac{1}{2} - \frac{i+1}{n}\right) \left[g G^{-1}\left(\frac{i}{n}\right) - g G^{-1}\left(\frac{i-1}{n}\right) \right].$$

$$\text{Since } g^{-1}G(x) = \begin{cases} x & \text{for } 0 < x \leq \frac{1}{2} \\ 1-x & \text{for } \frac{1}{2} < x \leq 1 \end{cases},$$

then

$$L\left(\frac{i}{n}\right) = \begin{cases} 2\left(\frac{i}{n}\right) - \frac{1}{2} & \text{for } \frac{i}{n} \leq \frac{1}{2} \\ \frac{3}{2} - 2\left(\frac{i}{n}\right) & \text{for } \frac{i}{n} > \frac{1}{2} \end{cases}.$$

Thus we can consider

$$(1.5.12) \quad S_n = \frac{1}{n} \sum_{i=1}^n L\left(\frac{i}{n}\right) X_{i,n}$$

$$\text{where } L(x) = \begin{cases} 2x - \frac{1}{2} & \text{for } 0 < x \leq \frac{1}{2} \\ \frac{3}{2} - 2x & \text{for } \frac{1}{2} < x \leq 1 \end{cases}.$$

We can show that

$$\int_{-\infty}^{\infty} xL(G(x))dG(x) = 0.$$

It follows from Lemma 1.5.1,

$$\mathcal{L}(\sqrt{n} S_n) \rightarrow N(0, \sigma^2(G)).$$

where

$$\begin{aligned}
(1.5.13) \quad \sigma^2(G) &= 2 \int \int_{s < t} L(G(s))L(G(t))G(s)(1-G(t))dsdt. \\
\sigma^2(G) &= 2 \int_0^1 \left[\int_0^t \frac{L(s)}{g G^{-1}(s)} s ds \right] \frac{L(t)}{g G^{-1}(t)} (1-t) dt \\
&= 2 \int_0^{\frac{1}{2}} \left[\int_0^t \frac{L(s)}{g G^{-1}(s)} s ds \right] \frac{L(t)}{g G^{-1}(t)} (1-t) dt \\
&\quad + 2 \int_{\frac{1}{2}}^1 \left[\int_0^{\frac{1}{2}} \frac{L(s)}{g G^{-1}(s)} s ds + \int_{\frac{1}{2}}^t \frac{L(s)}{g G^{-1}(s)} s ds \right] \frac{L(t)(1-t)}{g G^{-1}(t)} dt \\
&= 2 \cdot \frac{1}{48} + 2 \cdot \frac{5}{48} = \frac{1}{4}.
\end{aligned}$$

By Theorem 2.1 of Barlow and Van Zwet [11], $H_n^{-1}(1) \rightarrow G(X_{n,n})$ a.s.

Since $G(X_{n,n}) \rightarrow 1$, therefore $H_n^{-1}(1) \rightarrow 1$ a.s.

Now $T_n = \frac{\sqrt{n}S_n}{H_n^{-1}(1)}$, hence by Slutsky's theorem,

we have the following result.

Theorem 1.5.2.

$$(1.5.14) \quad \mathcal{L} \left(\sqrt{n} \left(\frac{V_n}{n} - \frac{1}{2} \right) \right) \rightarrow N \left(0, \frac{1}{4} \right)$$

1.6. Selecting a subset which contains all populations better than a control.

In addition to π_1, \dots, π_k , we have a control population π_0 . Let π_i have a continuous distribution F_i , $i = 0, 1, \dots, k$. We are given a sample size n from each of the $(k+1)$ populations π_i , $i = 0, 1, \dots, k$. Let $\xi_{\alpha i}$ be the unique α -quantile, $i = 0, 1, \dots, k$. A population π_i ($i = 1, 2, \dots, k$) is defined as better than the control π_0 if $\xi_{\alpha i} \geq \xi_{\alpha 0}$.

We want to select a subset such that the probability is at least p^* ($\frac{1}{k} < p^* < 1$) that all populations π_i ($i = 1, \dots, k$) for which $\xi_{\alpha i} \geq \xi_{\alpha 0}$ will be included in the subset. We regard any such selection a correct selection (CS). We assume that for a fixed i , $i = 1, \dots, k$, either $F_i(x) \geq F_0(x)$ or $F_i(x) \leq F_0(x) \forall x > 0$. Thus π_i is better than π_0 iff $F_i(x) \leq F_0(x) \forall x > 0$. Here π_0 is not known and we assume $F_0 \prec_* G$. Let T_i be the r -th order statistic from F_i where $r \leq (n+1)\alpha < r+1$ and let H_i be the c.d.f. of the r -th order statistic from F_i , $i = 0, 1, \dots, k$. We propose the rule R_{11} as follows,

R_{11} : Select π_i if and only if

$$(1.6.1) \quad T_i \geq c_{11} T_0$$

where c_{11} ($0 < c_{11} < 1$) is determined to satisfy the basic probability requirement. Let t be the number of π_i 's better than π_0 and let Δ be the index set for those π_i , t is unknown. Let $G_r(x)$ be the c.d.f. of the r -th order statistic of sample size r from G . Let Ω be the set of $(k+1)$ -tuples (F_0, F_1, \dots, F_k) .

Theorem 1.6.1. If $F_0 \prec_* G$ and for a fixed i , $i = 1, \dots, k$, either $F_i(x) \geq F_0(x)$ or $F_i(x) \leq F_0(x)$, $x > 0$, then

$$(1.6.2) \quad \inf_{\Omega} P[CS|R_{11}] = \int_0^{\infty} \bar{G}_r^k(c_{11}x) dG_r(x) \text{ where } \bar{G}_r(x) = 1 - G_r(x).$$

Proof. $P[CS|R_{11}] = P[T_i \geq c_{11} T_0 \quad i \in \Delta]$

$$= \int_0^{\infty} \prod_{i \in \Delta} [1 - H_i(c_{11}x)] dH_0(x)$$

Since $F_i(x) \leq F_0(x)$ for $i \in \Delta$, then $H_i(x) \leq H_0(x)$.

Thus

$$\begin{aligned} P[CS|R_{11}] &\geq \int_0^{\infty} \pi_{i \in \Delta} [1-H_0(c_{11}x)] dH_0(x) \\ &= \int_0^{\infty} [1-H_0(c_{11}x)]^t dH_0(x) \geq \int_0^{\infty} [1-H_0(c_{11}x)]^k dH_0(x) \\ &= P[\min_{1 \leq i \leq k} Z_i \geq c_{11} Z_0] \end{aligned}$$

where Z_0, Z_1, \dots, Z_k are i.i.d. with c.d.f. $H_0(x)$. Since $F_0 \prec^* G$ (see Barlow and Gupta [5]), then

$$P[\min_{1 \leq i \leq k} Z_i \geq c_{11} Z_0] \geq \int_0^{\infty} [\bar{G}_r(c_{11}x)]^k dG_r(x).$$

This completes the proof.

1.7. Quantile selection rule for a restricted family of distributions

We assume that each $F_i(G)$ has a unique α -quantile $\xi_{i\alpha}(\eta_\alpha)$ and $F_i(0) = 0 = G(0)$, $i = 1, 2, \dots, k$.

Let $\xi_{[1]\alpha} \leq \dots \leq \xi_{[k]\alpha}$ be ordered values of $\xi_{i\alpha}$, $i = 1, \dots, k$. A best population is defined as the one having $\xi_{[k]\alpha}$. Let T_i denote the r -th order statistic from F_i , where $r \leq (n+1)\alpha < r+1$.

The rule we propose is

R_{12} : Select π_i if and only if

$$(1.7.1) \quad T_i \geq \max_{1 \leq j \leq k} T_j - c_{12}$$

where c_{12} ($0 < c_{12} \leq 1$) is determined so as to satisfy the probability requirement $P[CS|R_{12}] \geq P^*$.

As before let Ω be the set of all k -tuples (F_1, \dots, F_k) . Also if F has a unique α -quantile ξ_α , we define $F \underset{S^*}{<} G$ if and only if $F(0) = 0 = G(0)$ and $G^{-1}F(x)$ is concave-convex about the ξ_α , on the support of F .

For example, if $G(x) = 1 - e^{-x}$, then $F \underset{S^*}{<} G$ is equivalent to saying that $F(x)$ is DFR for $x \leq \xi_\alpha$ and $F(x)$ is IFR for $x > \xi_\alpha$, on the support of F . We are interested in this "turning point" ξ_α . Let $H_{r,i}(x)$ be the c.d.f. of the r -th order statistic from $F_{[i]}$ and let $G_r(x)$ be the c.d.f. of the r -th order statistic from G .

Theorem 1.7.1. If $F_{[i]}(x) \geq F_{[k]}(x)$, $x \geq 0$ and $i = 1, \dots, k$, $F_{[k]} \underset{S^*}{<} G$ and $F_{[k]}(y + \xi_{[k]\alpha}) \geq G(y + \eta_\alpha)$, for $y > 0$, then

$$(1.7.2) \quad \text{Inf}_\Omega P[CS|R_{12}] = \int_0^\infty G_r^{k-1}(x+c_{12}) dG_r(x)$$

where $G_r(x)$ is the cdf of r -th order statistic from G .

Proof.

$$(1.7.3) \quad P[CS|R_{12}] = \int_0^{k-1} \prod_{i=1}^r H_{r,i}(x+c_{12}) dH_{r,k}(x) \\ \geq \int_0^\infty [H_{r,k}(x+c_{12})]^{k-1} dH_{r,k}(x) = P[X_{r,k} \geq \max_{1 \leq i \leq k-1} X_{r,i} - c_{12}]$$

where $X_{r1}, \dots, X_{r,k}$ are i.i.d. r.v.'s with distribution $H_{r,k}$.

Let $\varphi(x) = G^{-1}F_{[k]}(x) = G_r^{-1}H_{r,k}(x)$. Since $G^{-1}F_{[k]}(x)$ is concave-convex about $\xi_{[k]\alpha}$, then $\varphi'(x)$ is decreasing in x for $0 < x < \xi_{[k]\alpha}$.

and $\varphi'(x)$ is increasing in x for $x > \xi_{[k]\alpha}$.

So that $\varphi'(x) \geq \varphi'(\xi_{[k]\alpha}) \forall x > 0$. Since $F_{[k]}(y + \xi_{[k]\alpha}) \geq G(y + \eta_\alpha)$ for $y > 0$, then $F_{[k]}(x) \geq G(x + \eta_\alpha - \xi_{[k]\alpha})$ for $x \geq \xi_{[k]\alpha}$.

Hence
$$\frac{G^{-1}F_{[k]}(x) - \eta_\alpha}{x - \xi_{[k]\alpha}} \geq 1.$$

Let $A(x) = G^{-1}F_{[k]}(x) - G^{-1}F_{[k]}(\xi_{[k]\alpha}) = G^{-1}F_{[k]}(x) - \eta_\alpha$, then

$$\frac{A(x) - A(\xi_{[k]\alpha})}{x - \xi_{[k]\alpha}} \geq 1.$$

Let $x \rightarrow \xi_{[k]\alpha}$, then $A'(\xi_{[k]\alpha}) \geq 1$.

Since
$$\frac{d}{dx} A(x) = A'(x) = \frac{f_{[k]}(x)}{g[G^{-1}F_{[k]}(x)]},$$

then
$$A'(\xi_{[k]\alpha}) = \frac{f_{[k]}(\xi_{[k]\alpha})}{g(\eta_\alpha)} \geq 1.$$

Hence $\varphi'(x) \geq 1 \quad \forall x > 0$.

Since $\frac{\varphi(\max_{1 \leq i \leq k} X_{r,i}) - \varphi(X_{r,k})}{\max_{1 \leq i \leq k} X_{r,i} - X_{r,k}} = \varphi'(t)$ for some t such that

$X_{r,k} \leq t \leq \max_{1 \leq i \leq k} X_{r,i}$ and $\varphi(X_{r,i}) = Y_{r,i}$ has distribution G_r ,

then

$$\max_{1 \leq i \leq k} Y_{r,i} - Y_{r,k} \geq \max_{1 \leq i \leq k} X_{r,i} - X_{r,k}.$$

From (1.7.3), we obtain

$$\begin{aligned} P[CS|R_{12}] &\geq P[\max_{1 \leq i \leq k} Y_{r,i} - Y_{r,k} \leq c_{12}] \\ &= \int_0^\infty G_r^{k-1}(x + c_{12}) dG_r(x). \end{aligned}$$

This completes the proof.

1.8. Estimation of ordered parameters

We assume that $F_i(x) = F\left(\frac{x}{\theta_i}\right) \forall x$ and $i = 1, \dots, k$, where F_i and θ_i are unknown and $\theta_i > 0$, $i = 1, \dots, k$. Let $\theta_{[1]} \leq \dots \leq \theta_{[k]}$ be ordered values of θ_i 's. We assume that each $F_i(F)$ has a unique α -quantile $\xi_i(\xi)$, $i = 1, \dots, k$. Let $\xi_{[1]} \leq \dots \leq \xi_{[k]}$ be ordered values of ξ_i 's. We are given a sample of size n from each of the k populations. Let T_i be the total life statistic until r -th failure from F_i , $i=1, \dots, k$, $T^*(T)$ be the total life statistic until r -th failure from $F(G)$ and let T_i^* be the r -th order statistic from F_i , $i = 1, \dots, k$, where $r \leq (n+1)\alpha < r+1$.

Let $T_{[1]} \leq \dots \leq T_{[k]}$ be order values of T_i 's and let $T_{[1]}^* \leq \dots \leq T_{[k]}^*$ be order values of T_i^* 's. Let $T_{(i)}$ and $T_{(i)}^*$ be associated with $\theta_{[i]}$, $i = 1, \dots, k$.

(A) Estimation of $\theta_{[i]}$ based on $T_{[i]}$.

Theorem 1.8.1. If $F < G$, $F(0) = G(0) = 0$ and if $\theta_0 = \int_0^{\infty} x dF(x) = \int_0^{\infty} x dG(x)$ where θ_0 is known, then

$$(1.8.1) \quad P\left[\theta_{[i]} \leq \frac{1}{\Delta} T_{[i]}\right] \geq \beta \quad \text{if } \Delta \leq \theta_0(n-r+1),$$

where $\Delta = W^{-1}\left(1 - \beta^{\frac{1}{k-i+1}}\right)$ and $W(x)$ is the c.d.f. of T from G .

Proof.
$$P\left[\theta_{[i]} \leq \frac{1}{\Delta} T_{[i]}\right] = P\left[T_{[i]} \geq \Delta \theta_{[i]}\right].$$

Since $P[T_{[i]} \leq t]$ is a non-increasing function of θ_{ℓ} ($1 \leq \ell \leq k$) (see Chen and Dudewicz [20]), then

$$\begin{aligned} P[T_{[i]} \geq \Delta \theta_{[i]}] &\geq p^{k-i+1} [T_{(i)} \geq \Delta \theta_{[i]}] \\ &= p^{k-i+1} [T^* \geq \Delta]. \end{aligned}$$

By Theorem 4.6 of Barlow and Proschan [8],

$$\begin{aligned} p^{k-i+1} [T^* \geq \Delta] &\geq p^{k-i+1} [T \geq \Delta] \\ &= \{1 - W(\Delta)\}^{k-i+1} = \{1 - (1 - \beta)^{\frac{1}{k-i+1}}\}^{k-i+1} = \beta. \end{aligned}$$

Let $\chi_{\alpha}^2(2r)$ denote the 100α percent point of a chi-square distribution with $2r$ d.f. Again, by Theorems 3.3 and 2.4 of Barlow and Proschan [9], we have the following theorems.

Theorem 1.8.2. If F is IFRA and $\int_0^{\infty} x dF(x) = \theta_0$ where θ_0 is known, then

$$(1.8.2) \quad P[\theta_{[i]} \leq c_1 T_{[i]}] \geq \beta_1 \text{ where}$$

$$c_1 = \begin{cases} \frac{2}{\theta_0 \chi_{1-\beta'}^2(2r)} & \text{if } \chi_{1-\beta'}^2(2r) \leq 2(n-r+1) \\ \frac{1}{(n-r+1)\theta_0} & \text{if } \chi_{1-\beta'}^2(2r) \geq 2(n-r+1) \end{cases}$$

$$\text{where } \beta' = \beta_1^{\frac{1}{k-i+1}}$$

Theorem 1.8.3. If F is IFR and $\int_0^{\infty} x dF(x) = \theta_0$ where θ_0 is known, then

$$(1.8.3) \quad P[\theta_{[i]} \geq c_2 T_{[i]}] \geq \beta_2 \text{ where } c_2 = \frac{2nk}{\theta_0^2 \beta''(2r)}$$

$$\text{and } \beta'' = \beta_2^{\frac{1}{i}}.$$

Let β_1, β_2 ($0 < \beta_1, \beta_2 < 1$) be such that $\beta_1 + \beta_2^{-1} = \gamma$ where γ is fixed and $0 < \gamma < 1$. By Theorems 1.8.2, and 1.8.3 and the fact that $P(AB) \geq P(A)+P(B)-1$ with $A = \{\theta_{[i]} \leq c_1 T_{[i]}\}$ and $B = \{\theta_{[i]} \geq c_2 T_{[i]}\}$, we have the following theorem for two-sided confidence interval of $\theta_{[i]}$.

Theorem 1.8.4. If F is IFR and $\int_0^{\infty} x dF(x) = \theta_0$ where θ_0 is known, then

$$(1.8.4) \quad P[c_2 T_{[i]} \leq \theta_{[i]} \leq c_1 T_{[i]}] \geq \gamma$$

where c_1, c_2 are defined as above.

(B) Estimation of $\xi_{[i]}$ based on $T_{[i]}$

Theorem 1.8.5. If $F < G$, then

$$(1.8.6) \quad P[\xi_{[i]} \leq c_1 T_{[i]}] \geq \beta_1 \text{ where } c_1 = \begin{cases} \Delta_1 & \text{if } \Delta_1(n-r+1) \geq 1 \\ \frac{1}{n-i+1} & \text{if } \Delta_1(n-r+1) \leq 1 \end{cases}$$

$$\text{where } \Delta_1 = \frac{G^{-1}(\alpha)}{W^{-1}(1-\beta_1^{\frac{1}{k-i+1}})} \text{ and}$$

$W(x)$ is the c.d.f. of T from G .

$$\begin{aligned}
\text{Proof. } & P[\xi_{[i]} \leq c_1 T_{[i]}] \\
& \geq p^{k-i+1} [T_{(i)} \geq \frac{1}{c_1} \xi_{[i]}] \quad (\text{see Theorem 1.8.1}) \\
& = p^{k-i+1} [T_{(i)} \geq \frac{1}{c_1} \cdot \theta_{[i]} \xi] \\
& = p^{k-1+1} [T^* \geq \frac{1}{c_1} \xi] \\
& = p^{k-i+1} [T^* \geq \frac{1}{c_1} F^{-1}(\alpha)] = p^{k-i+1} [F(c_1 T^*) \geq \alpha].
\end{aligned}$$

In a manner similar to the proof of Theorem 3.1 of Barlow and Proschan [9], we can show that

$$P[F(c_1 T^*) \geq \alpha] \geq \beta_1 \frac{1}{k-i+1}.$$

Hence

$$P[\xi_{[i]} \leq c_1 T_{[i]}] \geq \beta_1.$$

Looking at Theorem 2.1 of Barlow and Proschan [9], we can have the following theorem.

Theorem 1.8.6. If $F \underset{c}{\prec} G$ and $F(0) = 0 = G(0)$, then

$$(1.8.7) \quad P[\xi_{[i]} \geq c_2 T_{[i]}] \geq \beta_2 \text{ where } c_2 = \begin{cases} \Delta_2 & \text{if } \Delta_2^n \leq 1 \\ \frac{1}{n} & \text{if } \Delta_2^n \geq 1 \end{cases},$$

$$\Delta_2 = \frac{G^{-1}(\alpha)}{W^{-1}(\beta_2)} \text{ and }$$

$W(x)$ is the c.d.f. of T from G .

Let β_1, β_2 ($0 < \beta_1, \beta_2 < 1$) be such that $\beta_1 + \beta_2 - 1 = \gamma$ where γ is fixed and $0 < \gamma < 1$. By Theorem 1.8.5 and Theorem 1.8.6, and the fact that if $F(0) = 0 = G^{-1}(0) = 0$, then $F \underset{c}{\prec} G \Rightarrow F \underset{*}{\prec} G$ we

have the following theorem for two-sided confidence interval of $\xi[i]$.

Theorem 1.8.7. If $F \prec_c G$ and $F(0) = 0 = G^{-1}(0)$, then

$$(1.8.8) \quad P[c_2 T[i] \leq \xi[i] \leq c_1 T[i]] \geq \gamma$$

(c) Estimation of $\xi[i]$ based on $T^*[i]$.

Theorem 1.8.8. If $F \prec^* G$, then

$$(1.8.9) \quad (1) \quad P\left[\xi[i] \leq \frac{T^*[i]}{\Delta_1}\right] \geq \beta \text{ if } \Delta_1 = \frac{G_j^{-1}(1-\beta^{\frac{1}{k-i+1}})}{G^{-1}(\alpha)} \leq 1$$

$$(2) \quad P\left[\xi[i] \geq \frac{T^*[i]}{\Delta_2}\right] \geq \beta \text{ if } \Delta_2 = \frac{G_j^{-1}(\beta^{\frac{1}{i}})}{G^{-1}(\alpha)} \geq 1.$$

Proof. First, we want to prove (1).

$$\begin{aligned} P\left[\xi[i] \leq \frac{T^*[i]}{\Delta_1}\right] &= P[T^*[i] \geq \Delta_1 \xi[i]] \\ &\geq P^{k-i+1}[T^*_{(i)} \geq \Delta_1 \xi[i]] \quad (\text{see Theorem 1.8.5}) \\ &= \{1 - P[T^*_{(i)} \leq \Delta_1 \xi[i]]\}^{k-i+1} \\ &= \{1 - F_j(\Delta_1 \xi[i] \cdot \frac{\xi}{\xi[i]})\}^{k-i+1} = \{1 - F_j(\Delta_1 \xi)\}^{k-i+1} \end{aligned}$$

where $F_j(x)$ is the c.d.f. of the j -th order statistic from F .

Since $G^{-1}F(x) = G_j^{-1}F_j(x)$, it follows that $F_j(x) = G_j G^{-1}F(x)$. Hence

$$\begin{aligned}
F_j(\Delta_1 \xi) &= G_j G^{-1} F(\Delta_1 \xi) \\
&\leq G_j [\Delta_1 G^{-1} F(\xi)] \quad (\text{since } F \prec_* G \text{ and } 0 < \Delta_1 < 1) \\
&= G_j [\Delta_1 G^{-1}(\alpha)] \\
&= G_j [G_j^{-1} (1 - \beta^{\frac{1}{k-i+1}})] \\
&= 1 - \beta^{\frac{1}{k-i+1}}
\end{aligned}$$

so that

$$\{1 - F_j(\Delta_1 \xi)\}^{k-i+1} \geq \{1 - (1 - \beta^{\frac{1}{k-i+1}})\}^{k-i+1} = \beta.$$

Now, we want to prove (2).

$$\begin{aligned}
P[\xi_{[i]} \geq \frac{T_{[i]}^*}{\Delta_2}] &= P[T_{[i]}^* \leq \Delta_2 \xi_{[i]}] \\
&\geq P^i [T^*(i) \leq \Delta_2 \xi_{[i]}] \\
&= F_j^i(\Delta_2 \xi) \\
&= \{G_j G^{-1} F(\Delta_2 \xi)\}^i \\
&\geq G_j [\Delta_2 G^{-1} F(\xi)]^i \quad (\text{since } F \prec_* G \text{ and } \Delta_2 \geq 1) \\
&= \{G_j (\Delta_2 G^{-1}(\alpha))\}^i = \{G_j G_j^{-1}(\beta^{\frac{1}{i}})\}^i = \beta.
\end{aligned}$$

This completes the proof.

CHAPTER II
INTERVAL ESTIMATION AND SELECTION PROCEDURES
FOR A SET OF GOOD POPULATIONS

2.1. Introduction

In practice, one is always faced with the problem of selecting the better ones from a group of populations. Here, we consider such kind of problems. Suppose that we are given k ($k \geq 2$) populations π_1, \dots, π_k with distributions $F(x, \theta_i)$ where θ_i lies in an interval Λ on the real line, $i = 1, \dots, k$. The quality of the i -th population is characterized by the real-valued parameter θ_i . The population with the largest θ -value is called the best population. A population is considered a good one if its quality does not fall too much below that of the best population. If $d(\theta_i, \theta_j)$ is a suitable distance measure between θ_i and θ_j and if $\theta_{[k]} = \max(\theta_1, \dots, \theta_k)$, population π_i is

$$(2.1.1) \quad \begin{aligned} &\text{good,} && \text{if } d(\theta_{[k]}, \theta_i) \leq \Delta \\ &\text{bad,} && \text{if } d(\theta_{[k]}, \theta_i) > \Delta \end{aligned}$$

where Δ is a given positive constant. Let $\underline{X}_i = (X_{i1}, \dots, X_{in})$, $i = 1, \dots, k$ be mutually independent random samples, each of size n , from population π_i , $i = 1, \dots, k$ respectively. Let $T_i = T(\underline{X}_i)$ be

an appropriate estimator of θ_i . Let $g_n(t, \theta_i)$ and $G_n(t, \theta_i)$ be the density function and the distribution function of T_i , respectively. The results of this chapter relate to two cases,

$$(2.1.2) \quad (i) \quad G_n(t, \theta_i) = G_n(t - \theta_i), \quad -\infty < t < \infty, \quad -\infty < \theta_i < \infty, \quad \text{and}$$

$$(ii) \quad G_n(t, \theta_i) = G_n\left(\frac{t}{\theta_i}\right), \quad t > 0, \quad \theta_i > 0.$$

If the distribution function $F(x, \theta_i)$ of π_i belongs to a location parameter family, then $F(x, \theta_i) = F(x - \theta_i)$. The distribution function $G_n(t, \theta_i)$ of T_i which is based on a random sample size n is not necessarily of the form $G_n(t - \theta_i)$, but we can always find a statistic T_i such that T_i has the distribution function of the form $G_n(t - \theta_i)$. For example, let π_1, \dots, π_k be k normal populations with unknown means $\theta_1, \dots, \theta_k$ and a common known variance σ^2 . Let $T_i = \bar{X}_i$ be the sample mean of π_i based on a sample of size n . In this case, $G_n(x, \theta_i)$ (the c.d.f. of \bar{X}_i) is of the form $G_n(x, \theta_i) = G_n(x - \theta_i)$, where $G_n(x) = \Phi\left(\frac{\sqrt{n}}{\sigma} x\right)$.

If the distribution function $F(x, \theta_i)$ of π_i belongs to a scale parameter family, then $F(x, \theta_i) = F\left(\frac{x}{\theta_i}\right)$, $x > 0$, $\theta_i > 0$. As before we can find a statistic T_i such that T_i has the distribution function of the form $G_n\left(\frac{t}{\theta_i}\right)$. The results of this chapter deal with the cases when $G_n(t, \theta_i)$ is one of the two forms in (2.1.2).

In Definition (2.1.1) we take d as d_L or d_S , respectively, whenever θ_i is a location parameter or is a scale parameter for the density of T_i . d_L and d_S are defined as $d_L(a, b) = a - b$, $d_S(a, b) = \frac{a}{b}$.

Estimation of $\theta_{[k]}$ based on (T_1, \dots, T_k) has been considered by several authors. Construction of two-sided and one-sided confidence intervals for $\theta_{[k]}$ based on

$$(2.1.1) \quad T_{[k]} = \max_{1 \leq i \leq k} T_i$$

was considered by Dudewicz [24], Dudewicz and Tong [26], Saxena and Tong [65] and others.

For problem of selecting the good populations from a given collection was considered by Desu [21], Carroll, Gupta and Huang [19] and others.

We consider confidence interval I which is based on $T_{[k]}$. In this chapter, we are interested in finding the smallest sample size N such that the probability that I contains at least one good population is at least P^* where P^* is a specified number, $0 < P^* < 1$. The probability that I contains all good populations and the probability that I excludes all the bad population are discussed.

In Section 2.2, we investigate the above problems for location parameter case. The infima of coverage probabilities are obtained. Some special cases are discussed. In Section 2.3, we investigate the above problems for scale parameter case. In Section 4, we discuss the above problem for the means of normal populations and for the scale parameters of the gamma populations.

2.2. Results for the location parameter case.

In this section, we assume that θ_i is a location parameter for T_i , $i = 1, \dots, k$. It is assumed that there is no a priori

knowledge about the θ_i 's. For arbitrary fixed $d_1, d_2 \ni d_1 + d_2 > 0$, the confidence interval is defined as

$$(2.2.1) \quad I_1 = (T_{[k]} - d_1, T_{[k]} + d_2)$$

where $T_{[k]} = \max_{1 \leq i \leq k} T_i$.

For the location parameter case, we say that population π_i is

$$(2.2.2) \quad \begin{array}{ll} \text{good} & \text{if } \theta_i \geq \theta_{[k]}^{-\Delta}, \\ \text{bad} & \text{if } \theta_i < \theta_{[k]}^{-\Delta}, \end{array}$$

where $\theta_{[k]} = \max_{1 \leq i \leq k} \theta_i$.

For given $\underline{\theta} = (\theta_1, \dots, \theta_k)$ and $\Delta > 0$, let $\alpha_{\underline{\theta}}(1)$ denote the probability that I_1 contains at least one good population, let $\alpha_{\underline{\theta}}(2)$ denote the probability that I_1 contains all good populations and let $\alpha_{\underline{\theta}}(3)$ denote the probability that I_1 excludes all the bad populations.

Let us denote the ordered θ -values by $\theta_{[1]} \leq \dots \leq \theta_{[k]}$ and let Ω be the parameter space which is the collection of all possible parameter vectors $\underline{\theta} = (\theta_1, \dots, \theta_k)$. For given $\Delta > 0$, let m denote the unknown number of bad populations in the given collection of k populations. Clearly we have $0 \leq m \leq k-1$.

Let

$$(2.2.3) \quad \Omega_m = \{\underline{\theta} : \theta_{[1]} \leq \dots \leq \theta_{[m]} < \theta_{[k]}^{-\Delta} \leq \theta_{[m+1]} \leq \dots \leq \theta_{[k]}\}.$$

Then

$$(2.2.4) \quad \Omega = \bigcup_{m=0}^{k-1} \Omega_m.$$

We need the following lemma before we prove some theorems.

Lemma 2.2.1. Define $f(r) = AC^r - BD^r$, $r = 0, 1, \dots, m$ where $A \geq B > 0$.

If either (i) $1 < C \leq D$ or (ii) $1 > C \geq D > 0$, then

$$(2.2.4) \quad \min_{0 \leq r \leq m} f(r) = \min[f(0), f(m)].$$

Proof. Assume $1 < C \leq D$. For $r = 0, 1, \dots, m-1$,

$$f(r+1) - f(r) = AC^r(C-1) - BD^r(D-1)$$

$$f(r+1) \leq f(r) \text{ if and only if } \frac{A}{B} \left(\frac{C}{D}\right)^r \leq \frac{D-1}{C-1}, \quad r = 0, 1, \dots, m-1.$$

$$\text{Since } \frac{C}{D} \leq 1 \Rightarrow \frac{A}{B} \left(\frac{C}{D}\right)^{r+1} \leq \frac{D-1}{C-1}.$$

Hence

$$f(r+2) \leq f(r+1).$$

Similarly, one can show that if $f(r+1) \geq f(r)$ then $f(r) \geq f(r-1)$.

Hence the lemma follows.

If $1 > C \geq D > 0$, then by a similar argument as above, the result follows.

This completes the proof of the lemma.

Now we want to discuss the probability that I_1 contains at least one good population and the infimum of this probability over Ω . We need the following lemma.

Lemma 2.2.2. If the family of density functions $\{g_n(t-\theta): \theta \in \Lambda_1\}$ has a monotone likelihood ratio, for every $n \geq 1$, then for arbitrary fixed d_1, d_2 satisfying $d_1 + d_2 > \Delta$,

$$(2.2.5) \quad \inf_{\Omega_m} \alpha_{\theta}^*(1) = \min\{G_n^m(\Delta + d_1)G_n^{k-m}(d_1) - G_n^m(\Delta - d_2)G_n^{k-1}(-d_2), \\ G_n^{k-m}(d_1) - G_n^{k-m}(-d_2)\}.$$

Proof. For $\underline{\theta} \in \Omega_m$,

$$\begin{aligned}
 (2.2.6) \quad \alpha_{\underline{\theta}}(1) &= P\{T_{[k]}-d_1 < \theta_{[i]} < T_{[k]}+d_2, \text{ for some } i, i=m+1, \dots, k\} \\
 &= 1-P\{\theta_{[i]} \notin I, \text{ for each } i = m+1, \dots, k\} \\
 &= 1-P\{T_{[k]}+d_2 \leq \theta_{[m+1]}\} - P\{T_{[k]}-d_1 \geq \theta_{[k]}\} \text{ (since} \\
 &\quad d_1+d_2 > \Delta) \\
 &= P\{T_{[k]}-d_1 < \theta_{[k]}\} - P\{T_{[k]}+d_2 \leq \theta_{[m+1]}\} \\
 &= \prod_{j=1}^k G_n[\theta_{[k]}+d_1-\theta_{[j]}] - \prod_{j=1}^k G_n[\theta_{[m+1]}-d_2-\theta_{[j]}]
 \end{aligned}$$

Let $\delta_i = \theta_{[k]} - \theta_{[i]}$, $i = 1, \dots, k$, then $0 = \delta_k < \delta_{k-1} < \dots < \delta_{m+1} < \Delta < \delta_m < \dots < \delta_1$.

Hence

$$(2.2.7) \quad \alpha_{\underline{\theta}}(1) = \prod_{j=1}^k G_n(\delta_j+d_1) - \prod_{j=1}^k G_n(\delta_j-\delta_{m+1}-d_2).$$

It is easy to see that $\alpha_{\underline{\theta}}(1)$ is nondecreasing in δ_{m+1} , where

$\delta_{m+2} < \delta_{m+1} < \Delta$. Let $\delta_{m+1} = \delta_{m+2}$, then

$$\alpha_{\underline{\theta}}(1) = \prod_{\substack{j \neq m+1 \\ j \neq m+2}} G_n(\delta_j+d_1) G_n^2(\delta_{m+2}+d_1) - \prod_{\substack{j \neq m+1 \\ j \neq m+2}} G_n(\delta_j-\delta_{m+2}-d_2) G_n^2(-d_2).$$

In this case, also we can see that $\alpha_{\underline{\theta}}(1)$ is nondecreasing in δ_{m+2} ,

where $\delta_{m+3} < \delta_{m+2} < \Delta$. Let $\delta_{m+2} = \delta_{m+3}$. Repeating the process with each δ_j , $j = m+1, \dots, k-1$, thus we have

$$(2.2.8) \quad Q \equiv \inf_{H_1} \alpha_{\underline{\theta}}(1) = \prod_{j=1}^m G_n(\delta_j+d_1) G_n^{k-m}(d_1) - \prod_{j=1}^m G_n(\delta_j-d_2) G_n^{k-m}(-d_2)$$

where $H_1 = \{(\delta_{m+1}, \dots, \delta_{k-1}) : 0 < \delta_{k-1} < \dots < \delta_{m+1} < \Delta\}$.

For $i = 1, 2, \dots, m$,

$$\frac{\partial Q}{\partial \delta_i} = g(\delta_i + d_1) \left\{ A - B \frac{g(\delta_i - d_2)}{g(\delta_i + d_1)} \right\}$$

where $A = G_n^{k-m}(d_1) \prod_{\substack{j=1 \\ j \neq i}}^m G_n(\delta_j + d_1)$ and $B = G_n^{k-m}(-d_2) \prod_{\substack{j=1 \\ j \neq i}}^m G_n(\delta_j - d_2)$.

Since $A > B$ and $\frac{g(\delta_i - d_2)}{g(\delta_i + d_1)}$ is non-increasing in $\delta_i > \Delta$, $i = 1, \dots, m$, then Q is greater than or equal to the right hand side of equation (2.2.8) with either $\delta_i = \Delta$ or $\delta_i = \infty$, $i = 1, \dots, m$.

Thus

$$(2.2.9) \quad Q \geq \min_{0 \leq r \leq m} \{G_n^r(\Delta + d_1) G_n^{k-m}(d_1) - G_n^r(\Delta - d_2) G_n^{k-m}(-d_2)\}.$$

By Lemma 2.2.1, we get

$$(2.2.10) \quad \inf_{\Omega_m} \alpha_{\underline{\theta}}(1) = \min \{G_n^{k-m}(d_1) - G_n^{k-m}(-d_2), G_n^m(\Delta + d_1) G_n^{k-m}(d_1) - G_n^m(\Delta - d_2) G_n^{k-m}(-d_2)\}.$$

This completes the proof.

Theorem 2.2.1. If the family of density functions $\{g_n(t-\theta): \theta \in \Lambda_1\}$ has monotone likelihood ratio, for every $n \geq 1$, then for arbitrary fixed d_1, d_2 satisfying $d_1 + d_2 > \Delta$,

$$(2.2.11) \quad \inf_{\Omega} \alpha_{\underline{\theta}}(1) = \min \{G_n^k(d_1) - G_n^k(-d_2), G_n^{k-m}(d_1) G_n^m(\Delta + d_1) - G_n^{k-m}(-d_2) G_n^m(\Delta - d_2), m=1, \dots, k-1\}.$$

Proof. Since $\Omega = \bigcup_{m=0}^{k-1} \Omega_m$, by Lemma 2.2.2, we have

$$(2.2.12) \quad \inf_{\Omega} \alpha_{\underline{\theta}}(1) = \min_{0 \leq m \leq k-1} \{\min[A^{k-m} B^{k-B}, A^{k-m} C^{m-B} D^{k-m}]\}$$

where $A = G_n(d_1)$, $B = G_n(-d_2)$, $C = G_n(\Delta + d_1)$ and $D = G_n(\Delta - d_2)$.

By Lemma 2.2.1, $\min_{0 \leq m \leq k-1} [A^{k-m} B^m] = \min[A-B, A^k - B^k]$. From

(2.2.12), we get

$$(2.2.13) \quad \inf_{\underline{\theta}} \alpha_{\underline{\theta}}(1) = \min\{A-B, A^k - B^k, A^{k-m} C^m - B^{k-m} D^m, m=1, 2, \dots, k-1\}.$$

This completes the proof.

Corollary 2.2.1. Let the family of density functions $\{g_n(t-\theta) : \theta \in \Lambda_1\}$ have a monotone likelihood ratio, for every $n \geq 1$. For arbitrary fixed d_1, d_2 satisfying $g(-d_2) \geq g(d_1)$ and $d_1 + d_2 > \Delta$, then

$$(2.2.14) \quad \inf_{\Omega} \alpha_{\underline{\theta}}(1) = \min\{G_n(d_1) - G_n(-d_2), G_n^k(d_1) - G_n^k(-d_2), \\ G_n(d_1) G_n^{k-1}(\Delta + d_1) - G_n(-d_2) G_n^{k-1}(\Delta - d_2)\}.$$

Proof. By Theorem 2.2.1, we get

$$(2.2.15) \quad \inf_{\Omega} \alpha_{\underline{\theta}}(1) = \min\{A-B, A^k - B^k, A^{k-m} C^m - B^{k-m} D^m, m=1, \dots, k-1\}$$

where $A = G_n(d_1)$, $B = G_n(-d_2)$, $C = G_n(\Delta + d_1)$ and $D = G_n(\Delta - d_2)$.

Consider

$$(2.2.16) \quad f(m) = A^{k-m} C^m - B^{k-m} D^m \\ = A^k \left(\frac{C}{A}\right)^m - B^k \left(\frac{D}{B}\right)^m, \quad m = 0, 1, \dots, k-1.$$

Let

$$\psi(x) = \frac{G_n(x+d_1)}{G_n(x-d_2)} \\ \frac{d\psi(x)}{dx} = \frac{g_n(x+d_1)}{G_n^2(x-d_2)} \{G_n(x-d_2) - G_n(x+d_1)\} \frac{g_n(x-d_2)}{g_n(x+d_1)}$$

since $d_2 \geq -d_1$, $\frac{g(x-d_2)}{g(x-d_1)}$ is nondecreasing in $x > 0$ and $g(-d_2) \geq g(d_1)$,

then $\frac{d\psi(x)}{dx} \leq 0$ for $x > 0$. Thus $\psi(\Delta) = \frac{C}{D} \leq \psi(0) = \frac{A}{B}$.

By Lemma 2.2.1 and $1 < \frac{C}{A} \leq \frac{D}{B}$, then

$$(2.2.17) \quad \min_{0 \leq m \leq k-1} f(m) = \min\{f(0), f(k-1)\}.$$

Hence from (2.2.15) and (2.2.17),

$$\begin{aligned} \inf_{\Omega} \alpha_{\underline{\theta}}(1) &= \min\{A-B, A^k-B^k, f(0), f(k-1)\} \\ &= \min\{A-B, A^k-B^k, AC^{k-1}-BD^{k-1}\}. \end{aligned}$$

This completes the proof of the corollary.

Now we wish to discuss the probability that I_1 contains all good populations and the infimum of this probability over Ω .

Lemma 2.2.3. If the family of density functions $\{g_n(t-\theta)$:

$\theta \in \Lambda_1\}$ has a monotone likelihood ratio, for every $n \geq 1$, then for arbitrary fixed d_1, d_2 satisfying $d_1+d_2 > \Delta$,

$$(2.2.18) \quad \inf_{\Omega_m} \alpha_{\underline{\theta}}(2) \equiv \min_{0 \leq r \leq k-m-2} \min_{0 \leq \ell \leq m} \{ [G_n(d_1)]^{k+\ell-m-r-1} G_n^{r+1}(-\Delta+d_1) - [G_n(\Delta-d_2)]^{k+\ell-m-r-1} G_n^{r+1}(-d_2) \}.$$

Proof. For $\theta \in \Omega_m$,

$$\begin{aligned} (2.2.19) \quad \alpha_{\underline{\theta}}(2) &= P\{\theta[i] \in [T[k]-d_1, T[k]+d_2], i = m+1, \dots, k\} \\ &= P\{\theta[m+1] \geq T[k]-d_1, \theta[k] \leq T[k]+d_2\} \\ &= \prod_{j=1}^k G_n(\theta[m+1]-\theta[j]+d_1) - \prod_{j=1}^k G_n(\theta[k]-\theta[j]-d_2) \\ &= \prod_{j=1}^k G_n(\delta_j - \delta_{m+1} + d_1) - \prod_{j=1}^k G_n(\delta_j - d_2) \end{aligned}$$

where $\delta_j = \theta[k] - \theta[j]$, $j = 1, \dots, k$ and

$$0 = \delta_k < \delta_{k-1} < \dots < \delta_{m+1} < \Delta < \delta_m < \dots < \delta_1.$$

Since $\alpha_{\theta}(2)$ is nonincreasing in δ_{m+1} , where $\delta_{m+2} < \delta_{m+1} < \Delta$, then $\alpha_{\theta}(2) \geq Q$ where

$$(2.2.20) \quad Q = A \prod_{j=m+2}^{k-1} G_n(\delta_j - \Delta + d_1) - B \prod_{j=m+2}^{k-1} G_n(\delta_j - d_2),$$

$$A = \prod_{j=1}^m G_n(\delta_j - \Delta + d_1) G_n(d_1) G_n(-\Delta + d_1) \text{ and}$$

$$B = \prod_{j=1}^m G_n(\delta_j - d_2) G_n(\Delta - d_2) G_n(-d_2).$$

For $i = m+2, \dots, k-1$,

$$\frac{\partial Q}{\partial \delta_i} = g_n(\delta_i - \Delta + d_1) \left[A \prod_{\substack{j=m+2 \\ j \neq i}}^{k-1} G_n(\delta_j - \Delta + d_1) - B \prod_{\substack{j=m+2 \\ j \neq i}}^{k-1} G_n(\delta_j - d_2) \right] \frac{g_n(\delta_i - d_2)}{g_n(\delta_i - \Delta + d_1)}.$$

Again, since $d_2 \geq \Delta - d_1$ and $\frac{g_n(\delta_i - d_2)}{g_n(\delta_i - \Delta + d_1)}$ is nondecreasing in δ_i ,

$i = m+2, \dots, k-1$, then Q is greater than or equal to the right hand side of equation (2.2.20) with either $\delta_i = 0$ or $\delta_i = \Delta$, $i = m+2, \dots, k-1$.

Let $H_1 = \{(\delta_{m+1}, \delta_{m+2}, \dots, \delta_{k-1}) : 0 < \delta_{k-1} < \dots < \delta_{m+1} < \Delta\}$.

Hence,

$$(2.2.21) \quad \inf_{H_1} \alpha_{\theta}(2) = \min_{0 \leq r \leq k-m-2} [A G_n^r(-\Delta + d_1) G_n^{k-m-2-r}(d_1) - B G_n^r(-d_2) G_n^{k-m-2-r}(\Delta - d_2)].$$

Let $H_2 = \{(\delta_1, \dots, \delta_m) : \Delta < \delta_m < \dots < \delta_1\}$.

In a manner similar as above, for $r = 0, 1, \dots, k-m-2$, we can show that

$$\begin{aligned}
(2.2.22) \quad & \inf_{H_2} [AG_n^r(-\Delta+d_1)G_n^{k-m-2-r}(d_1) - BG_n^r(-d_2)G_n^{k-m-2-r}(\Delta-d_2)] \\
& = \min_{0 \leq \ell \leq m} \{G_n^{r+1}(-\Delta+d_1)[G_n(d_1)]^{k-m-1-r+\ell} - [G_n(\Delta-d_2)]^{k-m-1-r+\ell} \\
& \quad \quad \quad G_n^{r+1}(-d_2)\}.
\end{aligned}$$

From (2.2.21) and (2.2.22), we have our lemma.

Since $\Omega = \bigcup_{m=0}^{k-1} \Omega_m$, by Lemma 2.2.3, we can have the following theorem.

Theorem 2.2.2. If the family of density functions $\{g_n(t-\theta): \theta \in \Lambda_1\}$ has a monotone likelihood ratio, for every $n \geq 1$, then for arbitrary fixed d_1, d_2 satisfying $d_1 + d_2 > \Delta$,

$$(2.2.23) \quad \inf_{\Omega} \alpha_{\theta}(2) = \min_{0 \leq m \leq k-1} \min_{0 \leq r \leq k-m-2} \min_{0 \leq \ell \leq m} \{[G_n(d_1)]^{k+\ell-m-r-1} \\
G_n^{r+1}(-\Delta+d_1) - [G_n(\Delta-d_2)]^{k+\ell-m-r-1} G_n^{r+1}(-d_2)\}.$$

Lemma 2.2.4. Suppose that (i) $g_n(t) = g_n(-t) > 0$ for all t and (ii) the family of density functions $\{g_n(t-\theta): \theta \in \Lambda_1\}$ has a monotone likelihood ratio, for every $n \geq 1$. For arbitrary fixed d_1, d_2 satisfying $d_1 = d_2 = d > \Delta$,

$$(2.2.24) \quad \inf_{\Omega_m} \alpha_{\theta}(2) = \min\{A^{k-m-1}B^{-C^{k-m-1}}D, A^{k-1}B^{-C^{k-1}}D, AB^{k-m-1} \\
CD^{k-m-1}, A^{m+1}B^{k-m-1}C^{m+1}D^{k-m-1}\}$$

where $A = G_n(d_1)$, $B = G_n(-\Delta+d_1)$, $C = G_n(\Delta-d_2)$ and $D = G_n(-d_2)$.

Proof. By Lemma 2.2.3, we have

$$(2.2.25) \quad \inf_{\Omega_m} \alpha_{\theta}(2) = \min_{0 \leq r \leq k-m-2} \min_{0 \leq \ell \leq m} \{A^{k+\ell-m-r-1}B^{r+1}C^{k+\ell-m-r-1}D^{r+1}\}$$

where A , B , C and D are defined as above.

Consider

$$(2.2.26) \quad Q = \min_{0 \leq r \leq k-m-2} \{A^{k+\ell-m} \left[\frac{B}{A}\right]^{r+1} - C^{k+\ell-m} \left[\frac{D}{C}\right]^{r+1}\}.$$

Now we want to show $\frac{B}{A} \geq \frac{D}{C}$.

Let $f(x) = G_n(x-d) - G_n^2(x-d)$ for $0 \leq x \leq d$.

$$\frac{df(x)}{dx} = g_n(x-d)[1 - 2G_n(x-d)] \geq 0, \quad 0 < x < d.$$

Since $0 < \Delta < d$, $f(\Delta) \geq f(0)$ and $G_n(t) = 1 - G_n(-t)$, we can see that $\frac{B}{A} \geq \frac{D}{C}$.

By Lemma 2.2.1 and from (2.2.26),

$$(2.2.27) \quad Q = \min\{A^{k+\ell-m-1} B - C^{k+\ell-m-1} D, A^{\ell+1} B^{k-m-1} - C^{\ell+1} D^{k-m-1}\}.$$

From (2.2.25) and (2.2.27),

$$(2.2.28) \quad \inf_{\Omega_m} \alpha_{\theta}(2) = \min\left\{ \min_{0 \leq \ell < m} [A^{k+\ell-m-1} B - C^{k+\ell-m-1} D], \right. \\ \left. \min_{0 \leq \ell < m} [A^{\ell+1} B^{k-m-1} - C^{\ell+1} D^{k-m-1}] \right\}.$$

By Lemma 2.2.1 and (2.2.28), we get

$$(2.2.29) \quad \inf_{\Omega_m} \alpha_{\theta}(2) = \min\left\{ \min[A^{k-m-1} B - C^{k-m-1} D, A^{k-1} B - C^{k-1} D], \right. \\ \left. \min[AB^{k-m-1} - CD^{k-m-1}, A^{m+1} B^{k-m-1} - C^{m+1} D^{k-m-1}] \right\}$$

This completes the proof.

Since $\Omega = \bigcup_{m=0}^{k-1} \Omega_m$, by Lemma 2.2.4 and Lemma 2.2.1, we can obtain the following result.

Corollary 2.2.2. Suppose that (i) $g_n(t) = g_n(-t) > 0$ for all $t > 0$ and (ii) the family of density functions $\{g_n(t-\theta) : \theta \in \Lambda_1\}$ has a

monotone likelihood ratio, for every $n \geq 1$. For arbitrary fixed d_1, d_2 satisfying $d_1 = d_2 = d > \Delta$,

$$(2.2.30) \quad \inf_{\Omega} \alpha_{\underline{\theta}}(2) = \min\{A-C, A^k-B^k, AB^{k-1}-CD^{k-1}, A^{k-1}B-C^{k-1}D\}$$

where A, B, C and D are defined as in Lemma 2.2.4.

Now we wish to discuss the probability that I_1 excludes all the bad populations and the infimum of this probability over Ω .

Lemma 2.2.5. For $\underline{\theta} \in \Omega_m$,

$$(2.2.31) \quad \alpha_{\underline{\theta}}(3) \geq 1 - G_n^{k-m}(d_1)G(-\Delta+d_1)$$

Proof. For $\underline{\theta} \in \Omega_m$,

$$\begin{aligned} (2.2.32) \quad \alpha_{\underline{\theta}}(3) &= P[\theta_{[i]} \notin I_1, i = 1, \dots, m] \\ &\geq P[\theta_{[1]} \geq T_{[k]}+d_2] + P[\theta_{[m]} \leq T_{[k]}-d_1] \\ &= \prod_{j=1}^k G_n(\theta_{[1]} - \theta_{[j]} - d_2) + 1 - \prod_{j=1}^k G_n(\theta_{[m]} - \theta_{[j]} + d_1). \end{aligned}$$

Let $\delta_i = \theta_{[k]} - \theta_{[i]}$, $i = 1, 2, \dots, k$.

$$(2.2.33) \quad \alpha_{\underline{\theta}}(3) = 1 + \prod_{j=1}^k G_n(\delta_j - \delta_1 - d_2) - \prod_{j=1}^k G_n(\delta_j - \delta_m + d_1)$$

since $\frac{\partial \alpha_{\underline{\theta}}(3)}{\partial \delta_1} \leq 0$ for $\delta_1 > \delta_2$, by letting $\delta_1 = \infty$, then

$$(2.2.34) \quad \alpha_{\underline{\theta}}(3) \geq 1 - \prod_{j=2}^k G_n(\delta_j - \delta_m + d_1).$$

Since the right hand side of (2.3.34) is nonincreasing in δ_j for $j = 2, 3, \dots, m-1, m+1, \dots, k-1, k$ and is nondecreasing in δ_m , it follows

that the infimum of $1 - \prod_{j=2}^k G_n(\delta_j - \delta_m + d_1)$ over the set $\{(\delta_2, \dots, \delta_{k-1}) : 0 < \delta_{k-1} < \dots < \delta_{m+1} < \Delta < \delta_m < \dots < \delta_2\}$ is achieved at $\delta_{m+1} = \dots = \delta_{k-1} = \Delta$, $\delta_m = \Delta$ and $\delta_2 = \dots = \delta_{m-1} = \infty$. Thus

$$(2.2.35) \quad \alpha_{\underline{\theta}}(3) \geq 1 - G_n^{k-m}(d_1)G_n(d_1 - \Delta)$$

The proof is complete.

Since $\Omega = \bigcup_{m=0}^{k-1} \Omega_m$, by Lemma 2.2.5, we can easily lead to the following theorem.

Theorem 2.2.3. For arbitrary fixed d_1, d_2 satisfying $d_1 + d_2 > 0$,

$$(2.2.36) \quad \alpha_{\underline{\theta}}(3) \geq 1 - G_n(d_1)G_n(-\Delta + d_1).$$

Remark 2.2.1. (i) The right hand side of (2.2.36) is independent of d_2 . That is, for a given d_1 and for any fixed d_2 satisfying $d_1 + d_2 > 0$, $\alpha_{\underline{\theta}}(3)$ is bounded below by $1 - G_n(d_1)G_n(-\Delta + d_1)$. If the confidence interval is defined as $I_1 = (T_{[k]} - d_1, \infty)$ where d_1 is any given number, then $\inf_{\Omega} \alpha_{\underline{\theta}}(3)$ can be obtained as follows.

$$(2.2.37) \quad \inf_{\Omega} \alpha_{\underline{\theta}}(3) = 1 - G_n(d_1)G_n(-\Delta + d_1).$$

(ii) If we assume $\alpha_{\underline{\theta}}(4)$ is the probability that I_1 contains at least one good population and also I_1 excludes all bad populations, then $\alpha_{\underline{\theta}}(4) \geq \alpha_{\underline{\theta}}(1) + \alpha_{\underline{\theta}}(3) - 1$. From (2.2.11) and (2.2.36), we get

$$\alpha_{\underline{\theta}}(4) \geq \min\{G_n(d_1) - G_n(-d_2), G_n^k(d_1) - G_n^k(-d_2), G_n^{k-m}(d_1)G_n^m(\Delta + d_1) - G_n^{k-m}(-d_2)G_n^m(\Delta - d_2), m=1, \dots, k-1\} - G_n(d_1)G_n(-\Delta + d_1).$$

Similarly, we can find the lower bound of $\alpha_{\underline{\theta}}(5)$ which is defined as the probability that I_1 contains all good populations and also I_1 excludes all bad populations.

Let S' be the number of non-best (bad) populations that enter I_1 . We are interested in $E(S')$, the expected size (over Ω) of non-best populations that enter I_1 .

Theorem 2.2.4.

$$(2.2.38) \quad \sup_{\Omega} E(S') \leq (k-1)G_n(d_1)$$

Proof. For $\underline{\theta} \in \Omega_m$,

$$\begin{aligned} (2.2.39) \quad E(S) &= \sum_{i=1}^m P[\theta_{[i]} \in I] \\ &= \sum_{i=1}^m \left\{ \prod_{j=1}^k G_n(\theta_{[i]} - \theta_{[j]} + d_1) - \prod_{j=1}^k G_n(\theta_{[i]} - \theta_{[j]} - d_2) \right\} \\ &\leq m \left\{ \prod_{j=1}^k G_n(\theta_{[k]} - \theta_{[j]} + d_1) - \prod_{j=1}^k G_n(\theta_{[1]} - \theta_{[j]} - d_2) \right\} \\ &\leq m \left\{ \prod_{j=1}^m G_n(\theta_{[k]} - \theta_{[j]} + d_1) \prod_{j=m+1}^{k-1} G_n(\theta_{[k]} - \theta_{[j]} + d_1) G_n(d_1) - \right. \\ &\quad \left. G_n(-d_2) G_n^{k-1}[\theta_{[1]} - \theta_{[k]} - d_2] \right\} \\ &\leq m \{ G_n(d_1) G_n^{k-m-1}(\Delta + d_1) G_n^m(\theta_{[k]} - \theta_{[1]} + d_1) - G_n(-d_2) \cdot \\ &\quad G_n^{k-1}[\theta_{[1]} - \theta_{[k]} - d_2] \} \\ &\leq m G_n(d_1) G_n^{k-m-1}(\Delta + d_1). \end{aligned}$$

since $\Omega = \bigcup_{m=0}^{k-1} \Omega_m$, then

$$\begin{aligned} (2.2.40) \quad \sup_{\Omega} E(S') &\leq \max\{m G_n(d_1) G_n^{k-m-1}(\Delta + d_1) \mid m=0, 1, \dots, k-1\} \\ &= (k-1)G_n(d_1). \end{aligned}$$

In the following, we deal with special values of d_1, d_2 and Δ .

Case (i): Arbitrary but fixed d_1, d_2 satisfying $d_1 + d_2 > 0$ and $\Delta = 0$. Since $\Delta = 0$, the only good population is the one associated with $\theta_{[k]}$. Applying the earlier results,

$$(2.2.41) \quad \inf_{\Omega} \alpha_{\underline{\theta}}(1) = \inf_{\Omega} \alpha_{\underline{\theta}}(2) = \min\{G_n(d_1) - G_n(-d_2), G_n^k(d_1) - G_n^k(-d_2)\},$$

a result obtained by Dudewicz and Tong [26]. Also, in this case

$$(2.2.42) \quad \alpha_{\underline{\theta}}(3) \geq 1 - G_n^2(d_1).$$

Case (ii): One side (lower) confidence interval. Arbitrary but fixed d_1 and $d_2 = \infty$ with $\Delta > 0$. Our confidence interval is now $I_1 = (T_{[k]} - d_1, \infty)$. We can show that

$$(2.2.43) \quad \inf_{\Omega} \alpha_{\underline{\theta}}(1) = G_n^k(d_1),$$

$$(2.2.44) \quad \inf_{\Omega} \alpha_{\underline{\theta}}(2) = G_n(d_1)G_n^{k-1}(-\Delta + d_1) \text{ and}$$

$$(2.2.45) \quad \inf_{\Omega} \alpha_{\underline{\theta}}(3) = 1 - G_n(d_1)G_n(-\Delta + d_1).$$

Case (iii): One side (upper) confidence interval. Arbitrary fixed d_2 , $d_1 = \infty$ and $\Delta > 0$. Now our confidence interval is defined as $I_1 = (-\infty, T_{[k]} + d_2)$, we can show that

$$(2.2.45) \quad \inf_{\Omega} \alpha_{\underline{\theta}}(1) = 1 - G(-d_2),$$

$$(2.2.46) \quad \inf_{\Omega} \alpha_{\underline{\theta}}(2) = 1 - G_n^{k-1}(\Delta - d_2)G(-d_2) \text{ and}$$

$$(2.2.47) \quad \inf_{\Omega} \alpha_{\underline{\theta}}(3) = 0.$$

2.3. Result for the scale parameter case.

In this section, we assume θ_i is a scale parameter for T_i , $i = 1, \dots, k$. It is assumed that there is no a priori knowledge about the θ_i 's.

$$(2.3.1) \quad G_n(t, \theta_i) = \begin{cases} G_n\left(\frac{t}{\theta_i}\right) & \text{for } t \geq 0 \\ 0 & \text{for } t < 0, \end{cases}$$

for $\theta_i > 0$, $k = 1, \dots, k$.

For arbitrary fixed a, b satisfying $a \geq 1$ and $b \geq 1$, if the confidence interval is defined as

$$(2.3.2) \quad I_2 = \left(\frac{1}{a} T_{[k]}, b T_{[k]}\right)$$

where $T_{[k]} = \max_{1 \leq i \leq k} T_i$.

Let Δ be a given positive constant, $\Delta \geq 1$. We say population π_i is

$$(2.3.3) \quad \begin{array}{ll} \text{good} & \text{if } \theta_{[k]} \leq \Delta \theta_i \\ \text{bad} & \text{if } \theta_{[k]} > \Delta \theta_i \end{array}$$

where $\theta_{[k]} = \max_{1 \leq i \leq k} \theta_i$.

For given $\underline{\theta} = (\theta_1, \dots, \theta_k)$ and $\Delta \geq 1$, let $\beta_{\underline{\theta}}(1)$ denote the probability that I_2 contains at least one good population, let $\beta_{\underline{\theta}}(2)$ denote the probability that I_2 contains all good population and let $\beta_{\underline{\theta}}(3)$ denote the probability that I_2 excludes all the bad populations. Using the same arguments as in Section 2.2, we obtain the following results.

Theorem 2.3.1. If the family of density functions $\{g_n(\frac{x}{\theta}): \theta \in \Lambda_2\}$ has monotone likelihood ratio, for every $n \geq 1$, then for arbitrary fixed a, b satisfying $a \geq 1, b \geq 1$ and $ab > \Delta$, then

$$(2.3.4) \quad \inf_{\Omega} \beta_{\underline{\theta}}(1) = \min\{G_n(a) - G_n(\frac{1}{b}), G_n^k(a) - G_n^k(\frac{1}{b}), \\ G_n^{k-m}(a)G_n^m(a\Delta) - G_n^{k-m}(\frac{1}{b})G_n^m(\frac{\Delta}{b}), m=1, \dots, k-1\}.$$

Corollary 2.3.1. Let the family of density functions $\{g_n(\frac{x}{\theta}): \theta \in \Lambda_2\}$ has monotone likelihood ratio, for every $n \geq 1$. For arbitrary fixed a, b satisfying $a \geq 1, b \geq 1, g_n(\frac{1}{b}) \geq g_n(a)$ and $ab > \Delta$,

$$(2.3.5) \quad \inf_{\Omega} \beta_{\underline{\theta}}(1) = \min\{G_n(a) - G_n(\frac{1}{b}), G_n^k(a) - G_n^k(\frac{1}{b}), G_n(a)G_n^{k-1}(a\Delta) - \\ G_n(\frac{1}{b})G_n^{k-1}(\frac{\Delta}{b})\}.$$

Theorem 2.3.2. If the family of density functions $\{g_n(\frac{x}{\theta}): \theta \in \Lambda_2\}$ has a monotone likelihood ratio, for every $n \geq 1$, then for arbitrary fixed a, b satisfying $a \geq 1, b \geq 1$ and $ab > \Delta$,

$$(2.3.6) \quad \inf_{\Omega} \beta_{\underline{\theta}}(2) = \min_{0 \leq m \leq k-1} \min_{0 \leq r \leq k-m-2} \min_{0 \leq \ell \leq m} \{[G_n(a)]^{k+\ell-m-r-1} \\ G_n^{r+1}(\frac{a}{\Delta}) - [G_n(\frac{\Delta}{b})]^{k+\ell-m-r-1} G_n^{r+1}(\frac{1}{b})\}.$$

Theorem 2.3.3. For arbitrary fixed $a \geq 1$ and $b \geq 1$,

$$(2.3.7) \quad \beta_{\underline{\theta}}(3) \geq 1 - G_n(a)G_n(\frac{a}{\Delta}).$$

Let S' be the number of non-best (bad) populations that enter I_2 .

Also we can obtain the following result.

Theorem 2.3.4.

$$(2.3.8) \quad \sup_{\Omega} E(S') \leq (k-1)G_n(a).$$

In the following, we deal with some special values of d_1 , d_2 and Δ .

Case (i): Arbitrary fixed a, b satisfying $a \geq 1$ and $b \geq 1$ and $\Delta = 1$.

Since $\Delta = 1$ the only good population is the one associated with $\theta_{[k]}$. Applying the earlier results,

$$(2.3.9) \quad \inf_{\Omega} \beta_{\underline{\theta}}(1) = \inf_{\Omega} \beta_{\underline{\theta}}(2) = \min\{G_n(a) - G_n\left(\frac{1}{b}\right), G_n^k(a) - G_n^k\left(\frac{1}{b}\right)\},$$

a result obtained by Saxena and Tong [66]. Also, in this case

$$(2.3.10) \quad \beta_{\underline{\theta}}(3) \geq 1 - G_n^2(a).$$

Case (ii): One side (lower) confidence interval. Arbitrary but fixed $a \geq 1$ and $b = \infty$ with $\Delta \geq 1$. Our confidence interval is defined as $I_2 = (a T_{[k]}, \infty)$. We can show that

$$(2.3.11) \quad \inf_{\Omega} \beta_{\underline{\theta}}(1) = G_n^k(a),$$

$$(2.3.12) \quad \inf_{\Omega} \beta_{\underline{\theta}}(2) = G_n(a)G_n^{k-1}\left(\frac{a}{\Delta}\right) \text{ and}$$

$$(2.3.13) \quad \inf_{\Omega} \beta_{\underline{\theta}}(3) = 1 - G_n(a)G_n\left(\frac{a}{\Delta}\right).$$

Case (iii): One side (upper) confidence interval. For arbitrary fixed b , $a = \infty$ and $\Delta \geq 1$. Now our confidence interval is defined as

$$I_2 = (0, b T_{[k]}).$$

We can show that

$$(2.3.14) \quad \inf_{\Omega} \beta_{\underline{\theta}}(1) = 1 - G_n\left(\frac{1}{b}\right),$$

$$(2.3.15) \quad \inf_{\Omega} \beta_{\underline{\theta}}(2) = 1 - G_n\left(\frac{1}{b}\right) G_n^{k-1}\left(\frac{\Delta}{b}\right) \text{ and}$$

$$(2.3.16) \quad \inf_{\Omega} \beta_{\underline{\theta}}(3) = 0.$$

2.4. Some examples.

We shall discuss the above problem for the means of normal populations and for the scale parameter of the gamma population.

Let $\pi_1, \pi_2, \dots, \pi_k$ be k normal populations with unknown means $\theta_1, \theta_2, \dots, \theta_k$ and a common known variance σ^2 . Let \bar{X}_i be the sample mean from π_i based on a sample of size n . Let

$$(2.4.1) \quad T_i = \bar{X}_i \text{ and } T_{[k]} = \max_{1 \leq i \leq k} \bar{X}_i.$$

Let $G_n(x, \theta_i)$ be the c.d.f. of \bar{X}_i from π_i , $i = 1, \dots, k$. In this case, $G_n(x, \theta_i) = G_n(x - \theta_i)$ and

$$(2.4.2) \quad G_n(x) = \Phi\left(\frac{\sqrt{n}}{\sigma} x\right)$$

where Φ is the standard normal distribution function. The confidence interval I_1 is of the form $(T_{[k]} - d_1, T_{[k]} + d_2)$, where $d_1 + d_2 > \Delta$. In this case, applying Theorem 2.2.1, we get

$$(2.4.3) \quad \inf_{\Omega} \alpha_{\underline{\theta}}(1) = \min\{A - B, A^k - B^k, A^{k-m} C^m - B^{k-m} D^m, m=1, 2, \dots, k-1\}$$

where

$$A = \Phi\left(\frac{\sqrt{n} d_1}{\sigma}\right), B = \Phi\left(\frac{-\sqrt{n} d_2}{\sigma}\right), C = \Phi\left(\frac{\sqrt{n}(\Delta + d_1)}{\sigma}\right) \text{ and } D = \Phi\left(\frac{\sqrt{n}(\Delta - d_2)}{\sigma}\right).$$

For a given P^* , $\Delta > 0$, $d_1 > 0$, $d_2 > 0$ and $d_1 + d_2 > \Delta$, since

$\lim_{n \rightarrow \infty} A = 1$, $\lim_{n \rightarrow \infty} B = 0$, $\lim_{n \rightarrow \infty} C = 1$ and $\lim_{n \rightarrow \infty} D = 0$, there exists a

smallest integer N such that for every $n \geq N$, we have $\inf_{\Omega} \alpha_{\underline{\theta}}(1) \geq P^*$.

Let x_1 be the unique solution of $\phi(d_1 x) - \phi(-d_2 x) = P^*$. Let x_2 be the unique solution of $\phi^k(d_1 x) - \phi^k(-d_2 x) = P^*$. Let y_m be the unique solution of

$$\phi^{k-m}(d_1 x) \phi^m((\Delta + d_1)x) - \phi^{k-m}(-d_2 x) \phi^m((\Delta - d_2)x) = P^*, \quad m=1, 2, \dots, k-1.$$

Let

$$N_i = [\sigma^2 x_i^2], \quad i = 1, 2 \quad \text{and} \quad M_j = [\sigma^2 y_j^2], \quad j = 1, \dots, k-1,$$

where $[x]$ denotes the smallest integer $\geq x$. It follows that

the smallest sample size N required to satisfy $\inf_{\Omega} \alpha_{\underline{\theta}}(1) \geq P^*$ is given by

$$(2.4.4) \quad N = \min\{N_i, M_j, \quad i=1, 2, \quad j=1, \dots, k-1\}.$$

Consider the probability that I_1 contains all good populations.

For given P^* , $\Delta > 0$, $d_1 > \Delta$ and $d_2 > 0$, then

$$\lim_{n \rightarrow \infty} G_n(d_1) = 1, \quad \lim_{n \rightarrow \infty} G_n(-\Delta + d_1) = 1 \quad \text{and} \quad \lim_{n \rightarrow \infty} G_n(-d_2) = 0.$$

By Theorem 2.2.2, we can always find the smallest sample size N which satisfies

$$(2.4.5) \quad \inf_{\Omega} \alpha_{\underline{\theta}}(2) \geq P^*.$$

Consider the probability that I_1 excludes all good populations.

For given P^* , $\Delta > 0$, d_1, d_2 such that $d_1 + d_2 > 0$ and $d_1 < \Delta$, then

$$\lim_{n \rightarrow \infty} G(-\Delta + d_1) = 0.$$

By Theorem 2.2.3, we can always find the smallest sample size N required to satisfy

$$(2.4.6) \quad \inf_{\Omega} \alpha_{\underline{\theta}}(3) \geq P^*.$$

Now let us consider the problem in relation to the scale parameters of the populations. Let $\pi_1, \pi_2, \dots, \pi_k$ be k gamma distributions. The distribution associated with π_i has the density function $g(x, \alpha, \theta_i)$ where

$$(2.4.7) \quad g(x, \alpha, \theta_i) = \begin{cases} \frac{1}{\Gamma(\alpha)\theta_i^\alpha} x^{\alpha-1} e^{-\frac{x}{\theta_i}} & \text{for } x > 0 \\ 0 & \text{otherwise.} \end{cases}$$

We assume that the shape parameter α is the same for all populations and α is known. Let T_i be the sample sum from π_i based on n observations and let $T_{[k]} = \max_{1 \leq i \leq k} T_i$. Then the confidence interval is

given by

$$(2.4.8) \quad I_2 = \left(\frac{1}{a} T_{[k]}, b T_{[k]} \right) \text{ where } ab > \Delta.$$

Let $G_n(x, \theta_i)$ be the c.d.f. of T_i from $i = 1, \dots, k$.

In this case, $G_n(x, \theta_i) = G_n\left(\frac{x}{\theta_i}\right)$ and

(2.4.9) $G_n(x)$ is the c.d.f. of a gamma distribution with scale parameter 1 and shape parameter $n\alpha$. Since the condition in Theorem 2.3.1 holds, we get

$$(2.4.10) \quad \inf_{\Omega} \beta_{\underline{\theta}}(1) = \min\{G(a) - G\left(\frac{1}{b}\right), G^k(a) - G^k\left(\frac{1}{b}\right), G^{k-m}(a)G^m(a\Delta) - G^{k-m}\left(\frac{1}{b}\right)G^m\left(\frac{\Delta}{b}\right), m=1, \dots, k-1\}.$$

For given P^* , $\Delta > 1$, $a > 1$, $b > 1$ with $ab > \Delta$, since $\lim_{n \rightarrow \infty} G_n(a) = 1$, $\lim_{n \rightarrow \infty} G_n(\frac{1}{b}) = 0$ and $\lim_{n \rightarrow \infty} G_n(a\Delta) = 1$, there exists a smallest integer N such that for every $n \geq N$, we have $\inf_{\Omega} \beta_{\underline{\theta}}(1) \geq P^*$.

Let N_1 be the smallest positive integer such that $G_n(a) - G_n(\frac{1}{b}) \geq P^*$. Let N_2 be the smallest positive integer such that $G_n^k(a) - G_n^k(\frac{1}{b}) \geq P^*$. Let M_m be the smallest positive integer such that $G_n^{k-m}(a) G_n^m(a\Delta) - G_n^{k-m}(\frac{1}{b}) G_n^m(\frac{\Delta}{b})$, $m = 1, 2, \dots, k-1$.

It follows that the smallest sample size N required to satisfy $\inf_{\Omega} \alpha_{\underline{\theta}}(1) \geq P^*$ is given by

$$(2.4.11) \quad N = \min\{N_i, M_j, i=1, 2, j=1, 2, \dots, k-1\}.$$

Consider the probability that I_2 contains all good populations. For a given P^* , $\Delta > 1$, $a > \Delta$ and $b > 1$, then

$$\lim_{n \rightarrow \infty} G_n(a) = 1, \lim_{n \rightarrow \infty} G_n(\frac{a}{\Delta}) = 1 \text{ and } \lim_{n \rightarrow \infty} (\frac{1}{b}) = 0.$$

By Theorem 2.3.2, we can always find the smallest sample size N required to satisfy $\inf_{\Omega} \beta_{\underline{\theta}}(2) \geq P^*$.

Consider the probability that I_2 excludes all good populations. For given P^* , $\Delta > 1$, $b > 1$ and $a > 1$ with $a < \Delta$, then $\lim_{n \rightarrow \infty} G_n(\frac{a}{\Delta}) = 0$. By Theorem 2.3.3, we can always find the smallest sample size N required to satisfy $\inf_{\Omega} \beta_{\underline{\theta}}(3) \geq P^*$.

Remark 2.3.1. (i) If the sample size N is preassigned and P^* and Δ are given, then the confidence interval I_1 can always be found from (2.2.11), (2.2.23) and (2.2.36) with $G_n(x)$ as defined in (2.4.2) so that $\inf_{\Omega} \alpha_{\underline{\theta}}(i) \geq P^*$, $i = 1, 2, 3$.

Similarly, the confidence interval I_2 can always be obtained from (2.3.4), (2.3.6) and (2.3.7) with $G_n(x)$ as defined in (2.4.9) so that $\inf_{\Omega} \beta_{\underline{\theta}}(i) \geq P^*$, $i = 1, 2, 3$.

(ii) In location parameter case, if the population with the smallest θ -value is called the best population. Let Δ be given, we say population π_j is

$$(2.4.12) \quad \begin{array}{ll} \text{good} & \text{if } \theta_j \leq \theta_{[1]} + \Delta, \\ \text{bad} & \text{if } \theta_j > \theta_{[1]} + \Delta \end{array}$$

where $\theta_{[1]} = \min_{1 \leq i \leq k} \theta_i$.

Then the confidence interval is given by

$$(2.4.13) \quad I_1 = (T_{[1]} - d_1, T_{[1]} + d_2)$$

where $T_{[1]} = \min_{1 \leq i \leq k} T_i$.

The calculation of the coverage probabilities $\alpha_{\underline{\theta}}(i)$, $i = 1, 2, 3$, and their infima can be handled similarly as in Section 2.2.

The scale parameter case can be handled in a similar manner.

CHAPTER III
SELECTION PROCEDURES IN TERMS OF
MAJORIZATION AND WEAK MAJORIZATION

3.1. Introduction

Over the last fifty years, majorization and Schur functions have been applied to develop many useful inequalities in many branches of mathematics. Some of the references are: Schur [67], Ostrowski [57], Hardy, Littlewood and Pólya [45], Beckenbach and Bellman [17], Marshall and Proschan [52] and Marshall and Olkin [50]. In recent times the techniques of majorization and Schur functions have been applied to probability and statistics (see, for example, Marshall, Olkin and Proschan [51], Rinott [62], Proschan and Sethuraman [61], Nevius, Proschan and Sethuraman [55] and Gupta and Wong [44]).

Recently, Nevius, Proschan and Sethuraman [56] introduced a stochastic version of weak majorization. They have discussed some properties and made some applications.

Gupta [32] defined a class of selection procedures and considered some of its properties. Some additional results concerning the properties of this class of procedures were obtained by Gupta and Panchapakesan [36], Gupta and

Panchapkesan [37] defined a class of selection procedures which is a natural generalization of the class considered by Gupta [32]. In that paper, they obtained a sufficient condition for the probability of a correct selection to be nondecreasing in λ when $\lambda_1 = \dots = \lambda_k = \lambda$, where λ_i is the parameter associated with π_i , $i = 1, \dots, k$. They also obtained the supremum of the expected size of the selected subset and showed that if the sufficient condition holds, it takes place when the λ_i 's are equal.

In this chapter, we order the parameter space by means of majorization or weak majorization and propose some selection procedures when the parameter $\underline{\lambda}_i$ associated with π_i , $i=1, \dots, k$ is a vector. Section 3.2 defines a class of procedures R_h for selecting the population associated with vector $\underline{\lambda}_{[k]}$ (see Section 3.2 for definitions). A sufficient condition is obtained for the infimum of the probability of a correct selection to be Schur-convex in $\underline{\lambda}$. Also another sufficient condition is obtained for the supremum of the expected size of the selected subset to take place when $\underline{\lambda}_1 = \dots = \underline{\lambda}_k$. Some special cases of interest are discussed. In Section 3.3, a sufficient condition is obtained for the same infimum of the probability of a correct selection to be nondecreasing and Schur-convex in $\underline{\lambda}$. Section 3.4 defines a class of procedure R_H for the selection of the population with vector $\underline{\lambda}_{[1]}$. Some properties of the selection procedure are briefly discussed. Section 3.5 and 3.6 deal with selection procedures for multivariate normal distributions in terms of majorization

and weak majorization of mean vectors. Various cases corresponding to the known or unknown common covariance matrix Σ are studied. Properties of these selection procedures are also established.

3.2 A class of selection procedures R_h (and some properties) for vector-valued λ_j in terms of majorization

First we give the definitions of majorization, Schur-concave and Schur-convex functions.

Definition 3.2.1. A vector $\underline{a} = (a_1, \dots, a_p)$ is said to majorize a vector $\underline{b} = (b_1, \dots, b_p)$, if $a_1 \geq \dots \geq a_p$, $b_1 \geq \dots \geq b_p$, and $\sum_{i=1}^r a_i \geq \sum_{i=1}^r b_i$, $r = 1, \dots, p-1$, while $\sum_{i=1}^p a_i = \sum_{i=1}^p b_i$; we write, $\underline{a} \underset{m}{>} \underline{b}$.

The above definition is according to Beckenbach and Bellman [17] and differs slightly from that of Hardy, Littlewood and Pólya [45].

Definition 3.2.2. H is called a Schur-concave (Schur-convex) function if $H(\underline{a}) \leq H(\underline{b})$ ($H(\underline{a}) \geq H(\underline{b})$) whenever $\underline{a} \underset{m}{>} \underline{b}$. A Schur function is a function which is either Schur-concave or Schur-convex.

From above definitions, we know that majorization is a partial ordering in R^p , the p -dimensional Euclidean space and Schur functions are functions that are monotone with respect to this partial ordering. Now we state the following theorem.

Theorem 3.2.1. (Ostrowski [57])

Assume H is defined for $Z_1 \geq \dots \geq Z_p$ and first partial derivatives of H exist. Then $H(\underline{Z}) \geq H(\underline{Z}')$ for all $\underline{Z} > \underline{Z}'$ if and only if

$$(3.2.1) \quad \frac{\partial H}{\partial Z_i} - \frac{\partial H}{\partial Z_j} \geq 0, \text{ for } i < j, i, j=1, \dots, p.$$

Let π_1, \dots, π_k be k populations. Let Λ be an interval on the real line. Associated with π_i is a vector $\underline{\lambda}_i = (\lambda_{i1}, \dots, \lambda_{ip})$, $i = 1, \dots, k$ where X_{ij} has density $\phi(\lambda_{ij}, x)$, $i = 1, \dots, k$, $j=1, \dots, p$. We assume that $\lambda_{ij} \in \Lambda$ and $\lambda_{i1} \geq \dots \geq \lambda_{ip}$, $i=1, \dots, k$. We say that π_i is better than π_j if $\underline{\lambda}_i = (\lambda_{i1}, \dots, \lambda_{ip}) > \underline{\lambda}_j = (\lambda_{j1}, \dots, \lambda_{jp})$. It is assumed that among the k given populations, there always exists one population which is better than others. This is equivalent to saying that there exists a $\underline{\lambda}_{[k]}$ such that $\underline{\lambda}_{[k]} > \underline{\lambda}_i$ for all $i = 1, \dots, k$. This population is called the best population. If there are more than one "best" population, then we assume that one of them is tagged as the best. Based on one observation X_{i1}, \dots, X_{ip} from population π_i , we construct a suitable statistic $T_i = g(X_{i1}, \dots, X_{ip})$, $i = 1, \dots, k$. Let $F_{\underline{\lambda}_i}(x) = P[T_i \leq x]$ be the distribution function of T_i from population π_i . Based on the values of T_i from π_i , $i = 1, \dots, k$, we wish to define a class of procedures for selecting a non-empty subset of the k populations such that the probability is at least P^* ($\frac{1}{k} < P^* < 1$) that the population associated with $\underline{\lambda}_{[k]}$ is included in the selected subset.

Let

$$(3.2.2) \quad E = \{\underline{\lambda} = (\lambda_1, \dots, \lambda_p): \lambda_1 \geq \dots \geq \lambda_p, \lambda_i \in \Lambda, i=1, \dots, p\}$$

and

$$(3.2.3) \quad \Omega = \{\underline{\omega} = (\underline{\lambda}_1, \dots, \underline{\lambda}_k): \underline{\lambda}_i \in E, i=1, \dots, k \text{ and there exists some } i \text{ such that } \underline{\lambda}_i > \underline{\lambda}_j \text{ for each } j\}.$$

We wish to define a selection rule R such that

$$(3.2.4) \quad \inf_{\Omega} P[CS|R] \geq P^*.$$

Let $h \equiv h_{c,d}$; $c \in [1, \infty)$, $d \in [0, \infty)$ be a function defined on the real line satisfying the following properties: for every real x ,

$$(3.2.5) \quad \begin{aligned} & \text{(i)} \quad h_{c,d}(x) \geq x \\ & \text{(ii)} \quad h_{1,0}(x) = x \\ & \text{(iii)} \quad h_{c,d}(x) \text{ is continuous in } c \text{ and } d \\ & \text{(iv)} \quad h_{c,d}(x) \uparrow \infty \text{ as } d \rightarrow \infty \text{ and/or } xh_{c,d}(x) \uparrow \infty \text{ as } c \rightarrow \infty, x \neq 0. \end{aligned}$$

Some functions satisfying these properties that will be of interest are cx , $x+d$ and $cx+d$. Now, we define a class of procedures R_h as follows.

R_h : Select population π_i if and only if

$$(3.2.6) \quad h(T_i) \geq \max_{1 \leq r \leq k} T_r.$$

Because of (3.2.5)-(i), the procedure R_h will always select a

non-empty subset. Let $T_{(i)}$ be the random variable associated with $\lambda_{[i]}$ and let $F_{\lambda_{[i]}}$ be the cumulative distribution function (c.d.f.) of $T_{(i)}$, $i=1, \dots, k$, we have

$$(3.2.7) \quad P[CS|R_h] = P[h(T_{(k)}) \geq T_{(i)}, i=1, \dots, k-1] \\ = \int_{-\infty}^{\infty} \prod_{i=1}^{k-1} F_{\lambda_{[i]}}(h(x)) dF_{\lambda_{[k]}}(x).$$

We now assume that the distributions are Schur-concave in $\underline{\lambda}$, i.e., for $\underline{\lambda} \succ_m \underline{\lambda}'$, $F_{\underline{\lambda}}$ and $F_{\underline{\lambda}'}$ are distinct and

$$(3.2.8) \quad F_{\underline{\lambda}}(x) \leq F_{\underline{\lambda}'}(x), \text{ for all } x.$$

Then

$$P[CS|R_h] \geq \int_{-\infty}^{\infty} F_{\lambda_{[k]}}^{k-1}(h(x)) dF_{\lambda_{[k]}}(x).$$

Hence

$$(3.2.9) \quad \inf_{\Omega} P[CS|R_h] = \inf_{\underline{\lambda}} \psi(\underline{\lambda}; c, d, k)$$

where

$$(3.2.10) \quad \psi(\underline{\lambda}; c, d, k) = \int_{-\infty}^{\infty} F_{\lambda_{[k]}}^{k-1}(h(x)) dF_{\lambda_{[k]}}(x), \underline{\lambda} \in E.$$

In the following theorem, we are interested in a sufficient condition for the Schur-convexity of $E\psi(T, \underline{\lambda})$ where $\psi(x, \underline{\lambda})$ is some real-valued function, $\underline{\lambda} = (\lambda_1, \dots, \lambda_p) \in E$ and $T = g(X_{\lambda_1}, \dots, X_{\lambda_p})$ is a function of $X_{\lambda_1}, \dots, X_{\lambda_p}$. Let $F_{\underline{\lambda}}(x) = P[T \leq x]$.

Theorem 3.2.2. Let $\{F_{\underline{\lambda}}(x), \underline{\lambda} \in E\}$ be a family of continuous distributions on the real line such that $F_{\underline{\lambda}}(x)$ is a differentiable function in x and λ_i and let $\psi(x, \underline{\lambda})$ be a bounded real-valued and differentiable function in x and λ_i , $i = 1, \dots, p$. Then $E\psi(T, \underline{\lambda})$ is Schur-convex in $\underline{\lambda} \in E$ provided that

$$(3.2.11) \quad \left| \begin{array}{cc} \frac{\partial}{\partial x} F_{\underline{\lambda}}(x) & \frac{\partial}{\partial x} \psi(x, \underline{\lambda}) \\ (\frac{\partial}{\partial \lambda_i} - \frac{\partial}{\partial \lambda_j}) F_{\underline{\lambda}}(x) & (\frac{\partial}{\partial \lambda_i} - \frac{\partial}{\partial \lambda_j}) \psi(x, \underline{\lambda}) \end{array} \right| \geq 0, \text{ for } i < j, i, \\ j=1, \dots, P.$$

Proof. We argue along the lines of the proof of Theorem 2.1 of Gupta and Panchapkesan [37]. We assume that $F_{\underline{\lambda}}(x)$, $\underline{\lambda} \in E$ has the support I . Let

$$(3.2.12) \quad A(\underline{\lambda}) = E\psi(T, \underline{\lambda}) = \int_I \psi(x, \underline{\lambda}) dF_{\underline{\lambda}}(x).$$

Consider $\underline{u}_1, \underline{u}_2 \in E$, where $\underline{u}_i = (u_{i1}, \dots, u_{ip})$, $i = 1, 2$, $u_{i1} \geq u_{i2} \geq \dots \geq u_{ip}$, $i = 1, 2$ and assume $\underline{u}_1 = \underline{u}_2$. Define

$$(3.2.13) \quad A_1(\underline{u}_1, \underline{u}_2) = \int_I \psi(x, \underline{u}_2) dF_{\underline{u}_1}(x),$$

$$(3.2.14) \quad A_2(\underline{u}_1, \underline{u}_2) = \int_I \psi(x, \underline{u}_1) dF_{\underline{u}_2}(x),$$

and

$$(3.2.15) \quad B(\underline{u}_1, \underline{u}_2) = A_1(\underline{u}_1, \underline{u}_2) + A_2(\underline{u}_1, \underline{u}_2).$$

By integrating $A_1(\underline{u}_1, \underline{u}_2)$ by part (see Gupta and Panchapkesan [37]),

$$(3.2.16) \quad B(\underline{u}_1, \underline{u}_2) = \text{a term independent of } \underline{u}_1 \\ + \int_I [\psi(x, \underline{u}_1) f_{\underline{u}_2}(x) - \psi'(x, \underline{u}_2) F_{\underline{u}_1}(x)] dx$$

where $\psi'(x, \underline{u}) = \frac{\partial}{\partial x} \psi(x, \underline{u})$ and $f_{\underline{u}}(x) = \frac{d}{dx} F_{\underline{u}}(x)$.

For $i < j$, we have

$$(3.2.17) \quad (\frac{\partial}{\partial u_{1i}} - \frac{\partial}{\partial u_{1j}}) B(\underline{u}_1, \underline{u}_2) = \int_I \{ f_{\underline{u}_2}(x) [\frac{\partial \psi(x, \underline{u}_1)}{\partial u_{1i}} - \frac{\partial \psi(x, \underline{u}_1)}{\partial u_{1j}}] \\ - \psi'(x, \underline{u}_2) [\frac{\partial F_{\underline{u}_1}(x)}{\partial u_{1i}} - \frac{\partial F_{\underline{u}_1}(x)}{\partial u_{1j}}] \} dx$$

We can show that, for $\ell = 1, \dots, p$,

$$(3.2.18) \quad \frac{\partial}{\partial \lambda_\ell} B(\underline{\lambda}, \underline{\lambda}) = \sum_{j=1}^2 \frac{\partial}{\partial u_{j\ell}} B(u_1, u_2) \Big|_{u_1=u_2=\underline{\lambda}}$$

where $\underline{\lambda} = (\lambda_1, \dots, \lambda_p)$ and $\lambda_1 \geq \dots \geq \lambda_p$.

From (3.2.18) we get

$$(3.2.19) \quad \frac{\partial}{\partial \lambda_i} B(\underline{\lambda}, \underline{\lambda}) = 2 \cdot \frac{\partial}{\partial u_{1i}} B(u_1, u_2) \Big|_{u_1=u_2=\underline{\lambda}}$$

and

$$(3.2.20) \quad \frac{\partial}{\partial \lambda_j} B(\underline{\lambda}, \underline{\lambda}) = 2 \frac{\partial}{\partial u_{1j}} B(u_1, u_2) \Big|_{u_1=u_2=\underline{\lambda}}$$

From (3.2.19) and (3.2.20), we get

$$(3.2.21) \quad \left(\frac{\partial}{\partial \lambda_i} - \frac{\partial}{\partial \lambda_j} \right) B(\underline{\lambda}, \underline{\lambda}) = 2 \left(\frac{\partial}{\partial u_{1i}} - \frac{\partial}{\partial u_{1j}} \right) B(u_1, u_2) \Big|_{u_1=u_2=\underline{\lambda}}$$

If

$$(3.2.22) \quad \begin{vmatrix} \frac{\partial}{\partial x} F_{u_2}(x) & \frac{\partial}{\partial x} \psi(x, u_2) \\ \left(\frac{\partial}{\partial u_{1i}} - \frac{\partial}{\partial u_{1j}} \right) F_{u_1}(x) & \left(\frac{\partial}{\partial u_{1i}} - \frac{\partial}{\partial u_{1j}} \right) \psi(x, u_1) \end{vmatrix} \geq 0, \quad \begin{matrix} \forall i < j, \\ i, j = 1, \dots, p, \end{matrix}$$

then from (3.2.21),

$$(3.2.23) \quad \left(\frac{\partial}{\partial u_{1i}} - \frac{\partial}{\partial u_{1j}} \right) B(u_1, u_2) \geq 0 \quad \text{for } i < j, \quad i, j = 1, \dots, p.$$

Hence if (3.2.22) holds, we have from (3.2.21)

$$(3.2.24) \quad \left(\frac{\partial}{\partial \lambda_i} - \frac{\partial}{\partial \lambda_j} \right) B(\underline{\lambda}, \underline{\lambda}) \geq 0 \quad \text{for } i < j, \quad i, j = 1, \dots, p.$$

Note that

$$(3.2.25) \quad B(\underline{\lambda}, \underline{\lambda}) = 2 A(\underline{\lambda}).$$

Applying Theorem 3.2.1 and from (3.2.24),

it follows that $B(\underline{\lambda}, \underline{\lambda})$ is Schur-convex in $\underline{\lambda}$ if (3.2.22) holds. Since $\underline{u}_1 = \underline{u}_2$, then $A(\underline{\lambda})$ is Schur-convex in $\underline{\lambda}$ if (3.2.11) holds.

This completes the proof of the theorem.

In some cases, we will be dealing with the function $\psi(x, \underline{\lambda})$ such that it satisfies

$$(3.2.26) \quad \frac{\partial}{\partial \lambda_1} \psi(x, \underline{\lambda}) = \dots = \frac{\partial}{\partial \lambda_p} \psi(x, \underline{\lambda}),$$

then $(\frac{\partial}{\partial \lambda_i} - \frac{\partial}{\partial \lambda_j})\psi(x, \underline{\lambda}) = 0$, $i < j$, $i, j = 1, \dots, p$. Hence

we have the following result.

Corollary 3.2.1. If $\psi(x, \underline{\lambda})$ satisfies (3.2.26), then $E\psi(T, \underline{\lambda})$ is Schur-convex in $\underline{\lambda}$ if

$$(3.2.27) \quad (\frac{\partial}{\partial \lambda_i} - \frac{\partial}{\partial \lambda_j})F_{\underline{\lambda}}(x) \cdot \frac{\partial}{\partial x} \psi(x, \underline{\lambda}) \leq 0, \quad i < j, \quad i, j = 1, \dots, p.$$

Corollary 3.2.2. If $F_{\underline{\lambda}}(x)$ is Schur-concave in $\underline{\lambda} \in E$, $\psi(x, \underline{\lambda})$ satisfies (3.2.26) and $\psi(x, \underline{\lambda})$ is nondecreasing in x , then $E\psi(T, \underline{\lambda})$ is Schur-convex in $\underline{\lambda} \in E$.

Proof. Since $F_{\underline{\lambda}}(x)$ is Schur-concave in $\underline{\lambda} \in E$,

$$(\frac{\partial}{\partial \lambda_i} - \frac{\partial}{\partial \lambda_j})F_{\underline{\lambda}}(x) \leq 0, \quad \text{for } i < j, \quad i, j = 1, \dots, p.$$

Also $\psi(x, \underline{\lambda})$ is nondecreasing in x , then $\frac{\partial}{\partial x} \psi(x, \underline{\lambda}) \geq 0$.

Hence by Corollary 3.2.1, the result follows.

As a special case, if $\psi(x, \underline{\lambda}) = \psi(x)$, then we have the following result.

Corollary 3.2.3. If $F_{\underline{\lambda}}(x)$ is Schur-concave in $\underline{\lambda}$ and $\psi(x)$ is nondecreasing in x , then $E\psi(T)$ is Schur-convex in $\underline{\lambda}$.

Theorem 3.2.3. For the procedure R_h , let $\{F_{\underline{\lambda}}(x), \underline{\lambda} \in E\}$ be a family of continuous distributions on the real line such that $F_{\underline{\lambda}}(x)$ and $F_{\underline{\lambda}}(h(x))$ are differentiable functions in x and λ_i , $i = 1, \dots, p$. Then $\psi(\underline{\lambda}; c, d, k)$ as defined in (3.2.10) is Schur-convex in $\underline{\lambda} \in E$ provided that

$$(3.2.28) \quad f_{\underline{\lambda}}(x) \left(\frac{\partial}{\partial \lambda_i} - \frac{\partial}{\partial \lambda_j} \right) F_{\underline{\lambda}}(h(x)) - h'(x) f_{\underline{\lambda}}(h(x)) \left(\frac{\partial}{\partial \lambda_i} - \frac{\partial}{\partial \lambda_j} \right) F_{\underline{\lambda}}(x) \geq 0,$$

for $i < j$, $i, j = 1, \dots, p$,

where $f_{\underline{\lambda}}(x) = \frac{d}{dx} F_{\underline{\lambda}}(x)$ and $h'(x) = \frac{d}{dx} h(x)$.

Proof. By letting $\psi(x, \underline{\lambda}) = F_{\underline{\lambda}}^{k-1}(h(x))$ and using Theorem 3.2.2. yields the proof.

We note that

$$(3.2.29) \quad \frac{\partial}{\partial \lambda_1} \psi(\underline{\lambda}; c, d, k) = \dots = \frac{\partial}{\partial \lambda_p} \psi(\underline{\lambda}; c, d, k), \text{ if}$$

$$(3.2.30) \quad f_{\underline{\lambda}}(x) \left(\frac{\partial}{\partial \lambda_i} - \frac{\partial}{\partial \lambda_j} \right) F_{\underline{\lambda}}(h(x)) - h'(x) f_{\underline{\lambda}}(h(x)) \left(\frac{\partial}{\partial \lambda_i} - \frac{\partial}{\partial \lambda_j} \right) F_{\underline{\lambda}}(x) = 0,$$

for $i < j$, $i, j = 1, \dots, k$.

Case (i): $F_{\underline{\lambda}}(x) = F(x - \xi(\underline{\lambda}))$, $-\infty < \xi(\underline{\lambda}) < \infty$ and $h(x) = x + d$, $d > 0$ where $\xi(\underline{\lambda})$ is a Schur-convex in $\underline{\lambda}$, then

$$(3.2.31) \quad \frac{\partial}{\partial \lambda_i} F_{\underline{\lambda}}(x) = -f(x - \xi(\underline{\lambda})) \frac{\partial \xi(\underline{\lambda})}{\partial \lambda_i} = -f_{\underline{\lambda}}(x) \frac{\partial \xi(\underline{\lambda})}{\partial \lambda_i}.$$

In this case, (3.2.30) is satisfied and $\psi(\underline{\lambda}; c, d, k)$ satisfies (3.2.29).

Case (ii): If $F_{\underline{\lambda}}(x) = F\left(\frac{x}{\eta(\underline{\lambda})}\right)$, $x > 0$, $\eta(\underline{\lambda}) > 0$ and $h(x) = cx$, $c \geq 1$ where $\eta(\underline{\lambda})$ is a Schur-convex in $\underline{\lambda}$, then

$$(3.2.32) \quad \frac{\partial F_{\underline{\lambda}}(x)}{\partial \lambda_i} = f\left(\frac{x}{\eta(\underline{\lambda})}\right) \left[-\frac{x}{\eta^2(\underline{\lambda})} \frac{\partial}{\partial \lambda_i} \eta(\underline{\lambda}) \right] = -\frac{x}{\eta(\underline{\lambda})} f_{\underline{\lambda}}(x) \frac{\partial \eta(\underline{\lambda})}{\partial \lambda_i}.$$

In this case, (3.2.30) is satisfied and $\psi(\underline{\lambda}; c, d, k)$ satisfies (3.2.29).

Actually, in the above two cases, $\psi(\underline{\lambda}; c, d, k)$ does not involve $\underline{\lambda}$, i.e., $\psi(\underline{\lambda}; c, d, k)$ is independent of $\underline{\lambda}$.

Let S denote the size of the subset selected by the procedure R_h . Let $\Omega' = \{\underline{\omega} = (\lambda_1, \dots, \lambda_k) : \lambda_i \in E, i=1, \dots, k \text{ and there exist } \ell_1, \dots, \ell_k \text{ such that } \lambda_{\ell_1} > \lambda_{\ell_2} > \dots > \lambda_{\ell_k}\}$.

Let $E_{\underline{\omega}}(S|R_h)$ be the expected size of the selected subset using R_h , $\underline{\omega} \in \Omega'$. Let $\lambda_{[1]} < \lambda_{[2]} < \dots < \lambda_{[k]}$, for $\underline{\omega} \in \Omega'$. It is easy to see that

$$(3.2.33) \quad E_{\underline{\omega}}(S|R_h) = \sum_{i=1}^k p_i,$$

where

$$(3.2.34) \quad p_i = \int \prod_{\substack{r=1 \\ r \neq i}}^k F_{\lambda_{[r]}}(h(x)) dF_{\lambda_{[i]}}(x),$$

and $F_{\lambda_{[i]}}$ is the c.d.f. of $T_{(i)}$ which is associated with $\lambda_{[i]}$, $i=1, \dots, k$. Using the same arguments as in Gupta and Panchapkesan [37] and Theorem 3.2.2, we have the following theorem.

Theorem 3.2.4. Let $\{F_{\underline{\lambda}}(x), \underline{\lambda} \in E\}$ and $\{F_{\underline{\lambda}}(h(x)), \underline{\lambda} \in E\}$ be as in the hypothesis of Theorem 3.2.3. For $\underline{\omega} \in \Omega'$, $E_{\underline{\omega}}(S|R_h)$ is Schur-convex

in $\underline{\lambda}$, where $\lambda_{[1]} = \dots = \lambda_{[m]} = \lambda < \lambda_{[m+1]} < \dots < \lambda_{[k]}$ and consequently $\sup_{\Omega'} E(S|R_h)$ takes place at \underline{v} where $\lambda_{[1]} = \dots = \lambda_{[k]} = \underline{v}$ provided that, for $\lambda_1 < \lambda_2$ and $\lambda_1, \lambda_2 \in E$, the following holds;

$$(3.2.35) \quad \left(\frac{\partial}{\partial \lambda_{1i}} - \frac{\partial}{\partial \lambda_{1j}} \right) F_{\lambda_1}(h(x)) f_{\lambda_2}(x) - \left(\frac{\partial}{\partial \lambda_{1i}} - \frac{\partial}{\partial \lambda_{1j}} \right) F_{\lambda_1}(x) f_{\lambda_2}(h(x)) \cdot h'(x) \geq 0,$$

for $i < j$, $i, j = 1, \dots, p$ where $\lambda_i = (\lambda_{i1}, \dots, \lambda_{ip})$, $i=1, 2$.

Remark 3.2.1. (i) If $\frac{f_{\lambda_1}(x)}{f_{\lambda_2}(x)}$ is nondecreasing in x for $\lambda_1 < \lambda_2$, then

(3.2.35) is satisfied, when (1) $F_{\lambda}(x) = F(x - \xi(\lambda))$, $-\infty < \xi(\lambda) < \infty$

and $h(x) = x + d$, $d > 0$ where $\xi(\lambda)$ is Schur-concave in λ , or when

(2) $F_{\lambda}(x) = F\left(\frac{x}{\eta(\lambda)}\right)$, $x \geq 0$, $\eta(\lambda) > 0$, and $h(x) = cx$, $c \geq 1$ where $\eta(\lambda)$ is Schur-concave in λ .

(ii) If (3.2.35) is satisfied, then

$$(3.2.36) \quad \sup_{\Omega'} E_{\omega} (S|R_h) = k \sup_{\Omega_0} P[CS|R_h]$$

where $\Omega_0 = \{\omega = (\lambda_1, \dots, \lambda_k) : \lambda_1 = \dots = \lambda_k, \lambda_i \in E, i=1, \dots, k\}$

(iii) If (1) $F_{\lambda}(x) = F(x - \xi(\lambda))$, $-\infty < \xi(\lambda) < \infty$ and $h(x) = x + d$

where $\xi(\lambda)$ is Schur-concave in λ or (2) $F_{\lambda}(x) = F\left(\frac{x}{\eta(\lambda)}\right)$, $x \geq 0$, $\eta(\lambda) > 0$ and $h(x) = cx$, $c \geq 1$ where $\eta(\lambda)$ is Schur-concave in λ ,

also if $\frac{f_{\lambda_1}(x)}{f_{\lambda_2}(x)}$ is nondecreasing in x for $\lambda_1 < \lambda_2$, then

$$(3.2.37) \quad \sup_{\Omega'} E_{\omega} (S|R_h) = k P^*.$$

3.3. A sufficient condition for the monotonicity in terms of weak majorization

First we give the definition of weak majorization.

Definition 3.3.1. A vector $\underline{a} = (a_1, \dots, a_p)$ is said to weakly majorize a vector $\underline{b} = (b_1, \dots, b_p)$, if $a_1 \geq \dots \geq a_p$, $b_1 \geq \dots \geq b_p$ and $\sum_{i=1}^r a_i \geq \sum_{i=1}^r b_i$, $r = 1, \dots, p$. If \underline{a} weakly majorize \underline{b} , then we write $\underline{a} \gg_m \underline{b}$.

We state the following theorem which is the characterization of weak majorization (see Nevius, Proschan and Sethuraman [56]).

Theorem 3.3.1. $\underline{Z} \gg_m \underline{Z}'$ if and only if $f(\underline{Z}) \geq f(\underline{Z}')$ for all nondecreasing Schur-convex functions, where $f(\underline{Z})$ is defined for $Z_1 \geq \dots \geq Z_p$.

Let $\underline{X}_\lambda = (X_{\lambda_1}, \dots, X_{\lambda_p})$ be a random vector of independent components where $\lambda \in E$ and the distribution function $F_{\lambda_i}(x)$ of random variable X_{λ_i} is stochastically increasing in λ_i , $i=1, \dots, p$.

Let

$$(3.3.1) \quad T = g(X_{\lambda_1}, \dots, X_{\lambda_p}) \text{ be a function of } X_{\lambda_1}, \dots, X_{\lambda_p}.$$

Let

$$(3.3.2) \quad F_\lambda(x) = P[T \leq x] = P[g(X_{\lambda_1}, \dots, X_{\lambda_p}) \leq x].$$

In the following theorem, we obtain a sufficient condition for $E\psi(T, \lambda)$ to be nondecreasing and Schur-convex in $\lambda \in E$ for some function $\psi(T, \lambda)$ as defined in Theorem 3.3.2.

Theorem 3.3.2. If $\{F_{\underline{\lambda}}(x) : \underline{\lambda} = (\lambda_1, \dots, \lambda_p) \in E\}$ is a family of continuous distributions on the real line such that $F_{\underline{\lambda}}(x)$ is a differentiable function in x and each λ_i , $F_{\lambda_i}(x)$ is stochastically increasing in λ_i and $\psi(x, \underline{\lambda})$ is a bounded real-valued and differentiable function in x and λ_i , $i = 1, \dots, p$, then $E\psi(T, \underline{\lambda}) = E\psi(g(\underline{X}_{\underline{\lambda}}), \underline{\lambda})$ is nondecreasing and Schur-convex in $\underline{\lambda} \in E$ provided

(3.3.3) (i) $\psi(g(\underline{X}_{\underline{\lambda}}), \underline{\lambda})$ is nondecreasing in $\underline{X}_{\underline{\lambda}}$ and $\underline{\lambda}$, and

$$(ii) \frac{\partial}{\partial x} F_{\underline{\lambda}}(x) \left(\frac{\partial}{\partial \lambda_i} - \frac{\partial}{\partial \lambda_j} \right) \psi(x, \underline{\lambda}) - \frac{\partial}{\partial x} \psi(x, \underline{\lambda}) \left(\frac{\partial}{\partial \lambda_i} - \frac{\partial}{\partial \lambda_j} \right) F_{\underline{\lambda}}(x) \geq 0,$$

for $i < j$, $i, j = 1, \dots, p$.

Proof. By Theorem 3.2.2. and (3.3.3)-(ii), $E\psi(T, \underline{\lambda})$ is Schur-convex in $\underline{\lambda}$.

Now we want to show that $E\psi(T, \underline{\lambda})$ is nondecreasing in $\underline{\lambda}$. Let $\underline{\lambda} = (\lambda_1, \dots, \lambda_p)$, $\lambda_1 \geq \dots \geq \lambda_p$ and $\underline{\lambda}' = (\lambda'_1, \dots, \lambda'_p)$, $\lambda'_1 \geq \dots \geq \lambda'_p$. Assume $\underline{\lambda} \geq \underline{\lambda}'$. We define $\underline{\lambda} \geq \underline{\lambda}'$ in the sense that $\lambda_i \geq \lambda'_i$, $i = 1, \dots, p$. Since $F_{\lambda_i}(x)$ associated with the random variable X_{λ_i} is stochastically increasing in λ_i , i.e. if $\lambda_i \geq \lambda'_i$, $i = 1, \dots, p$, then

$$(3.3.4) \quad X_{\lambda_i} \stackrel{st}{\geq} X_{\lambda'_i}, \quad i = 1, \dots, p.$$

Since $X_{\lambda_1}, \dots, X_{\lambda_p}$ are independent random variables, then

$$(3.3.5) \quad \underline{X}_{\underline{\lambda}} \stackrel{st}{\geq} \underline{X}_{\underline{\lambda}'}$$

By (3.3.3)-(i), since $\psi(g(\underline{X}_{\underline{\lambda}}), \underline{\lambda})$ is nondecreasing in $\underline{X}_{\underline{\lambda}}$, then

$$(3.3.6) \quad \psi(g(\underline{X}_{\underline{\lambda}}), \underline{\lambda}) \stackrel{st}{\geq} \psi(g(\underline{X}_{\underline{\lambda}'}, \underline{\lambda})).$$

Since $\psi(g(X_{\underline{\lambda}}, \cdot), \underline{\lambda}')$ is nondecreasing in $\underline{\lambda}'$ and $\underline{\lambda} \geq \underline{\lambda}'$,

$$(3.3.7) \quad \psi(g(X_{\underline{\lambda}}, \cdot), \underline{\lambda}) \geq \psi(g(X_{\underline{\lambda}}, \cdot), \underline{\lambda}').$$

From (3.3.6) and (3.3.7), we have

$$(3.3.8) \quad \psi(g(X_{\underline{\lambda}}, \cdot), \underline{\lambda}) \underset{st}{\geq} \psi(g(X_{\underline{\lambda}}, \cdot), \underline{\lambda}').$$

Hence

$$(3.3.9) \quad E\psi(T, \underline{\lambda}) \geq E\psi(T, \underline{\lambda}').$$

This completes the proof.

By applying Theorem 3.3.2, we can obtain the following results.

Corollary 3.3.1. If $F_{\underline{\lambda}}(x)$ is Schur-concave in $\underline{\lambda}$, $\psi(x, \underline{\lambda})$ satisfies (3.2.26), $\psi(x, \underline{\lambda})$ is nondecreasing in x , $\psi(g(X_{\underline{\lambda}}, \cdot), \underline{\lambda})$ is nondecreasing in $X_{\underline{\lambda}}$ and $\underline{\lambda}$ and $F_{\lambda_i}(x)$ is stochastically increasing in λ_i , $i=1, \dots, p$, then $E\psi(T, \underline{\lambda})$ is nondecreasing and Schur-convex in $\underline{\lambda}$.

Corollary 3.3.2. If $F_{\underline{\lambda}}(x)$ is Schur-concave in $\underline{\lambda}$, $\psi(x)$ is nondecreasing in x , $\psi(g(X_{\underline{\lambda}}, \cdot))$ is nondecreasing in $X_{\underline{\lambda}}$ and $F_{\lambda_i}(x)$ is stochastically increasing in λ_i , $i=1, \dots, p$, then $E\psi(T)$ is nondecreasing and Schur-convex in $\underline{\lambda}$.

By letting $\psi(x, \underline{\lambda}) = F_{\underline{\lambda}}^{k-1}(h(x))$ and using Theorem 3.3.1 and Theorem 3.3.2, we have the following theorem.

Theorem 3.3.3. Let $\{F_{\underline{\lambda}}(x), \underline{\lambda} \in E\}$ be a family of continuous distributions on the real line such that $F_{\underline{\lambda}}(x)$ and $F_{\underline{\lambda}}(h(x))$ are differentiable functions in x and λ_i and $F_{\lambda_i}(x)$ is stochastically increasing in λ_i , $i=1, \dots, p$. Let $\psi(\underline{\lambda}; c, d, k)$ be defined in (3.2.10).

Then $\psi(\underline{\lambda}; c, d, k)$ is nondecreasing in $\underline{\lambda} \in E$ in the sense of weak majorization, i.e., it is nondecreasing and Schur-convex in $\underline{\lambda}$, provided that

(3.3.10) (i) $F_{\underline{\lambda}}(h(g(X_{\underline{\lambda}})))$ is nondecreasing in $X_{\underline{\lambda}}$ and $\underline{\lambda}$ and

$$(ii) f_{\underline{\lambda}}(x) \left(\frac{\partial}{\partial \lambda_i} - \frac{\partial}{\partial \lambda_j} \right) F_{\underline{\lambda}}(h(x)) - h'(x) f_{\underline{\lambda}}(h(x)) \left(\frac{\partial}{\partial \lambda_i} - \frac{\partial}{\partial \lambda_j} \right)$$

$$F_{\underline{\lambda}}(x) \geq 0, \text{ for}$$

$$i < j, i, j = 1, \dots, p, \text{ where } f_{\underline{\lambda}}(x) = \frac{d}{dx} F_{\underline{\lambda}}(x).$$

3.4. Selection of the population associated with $\underline{\lambda}_{[1]}$.

If the best population is defined to be the one associated with $\underline{\lambda}_{[1]}$, where $\underline{\lambda}_{[1]} < \underline{\lambda}_i$, $i = 1, \dots, k$. We now define a class of procedures R_H for the selection of the population associated with $\underline{\lambda}_{[1]}$.

Let $H \equiv H_{c,d}$; $c \in [1, \infty)$, $d \in [0, \infty)$ be a function defined on the real line satisfying the following conditions. For every x

$$(3.4.1) \quad (i) H_{c,d}(x) \leq x$$

$$(ii) H_{1,0}(x) = x$$

$$(iii) H_{c,d}(x) \text{ is continuous in } c \text{ and } d$$

$$(iv) H_{c,d}(x) \uparrow \infty \text{ as } d \rightarrow \infty \text{ and/or } xH_{c,d}(x) \downarrow 0 \text{ as } c \rightarrow \infty.$$

Of particular interest are the functions $\frac{x}{c}$, $x-d$ and $\frac{x}{c} - d$.

A class of procedures R_H for selecting a subset containing the best is defined as follows.

R_H : Select population π_i if and only if

$$(3.4.2) \quad H(T_i) \leq \min_{1 \leq r \leq k} T_r.$$

The probability of a correct selection is given by

$$(3.4.3) \quad P[CS|R_H] = \int_{-\infty}^{\infty} \prod_{r=2}^k \bar{F}_{\underline{\lambda}[r]}(H(x)) dF_{\underline{\lambda}[1]}(x),$$

where $\bar{F}_{\underline{\lambda}}(x) = 1 - F_{\underline{\lambda}}(x)$.

Because of the assumption (3.2.8),

$$(3.4.4) \quad P[CS|R_H] \geq \int_{-\infty}^{\infty} \bar{F}_{\underline{\lambda}[1]}^{k-1}(H(x)) dF_{\underline{\lambda}[1]}(x).$$

Let $\Omega = \{\underline{\omega} = (\underline{\lambda}_1, \dots, \underline{\lambda}_k) : \underline{\lambda}_i \in E, i=1, \dots, k \text{ and there exists some } i \text{ such that } \underline{\lambda}_i < \underline{\lambda}_j \forall j\}$.

Hence

$$(3.4.5) \quad \inf_{\Omega} P[CS|R_H] = \inf_{\underline{\lambda}} \varphi(\underline{\lambda}; c, d, k)$$

where

$$(3.4.6) \quad \varphi(\underline{\lambda}; c, d, k) = \int \bar{F}_{\underline{\lambda}}^{k-1}(H(x)) dF_{\underline{\lambda}}(x) \text{ and } \underline{\lambda} \in E.$$

Using the same method of proof as in the case of R_h , we have the following results.

Theorem 3.4.1. For the procedure R_H , $\varphi(\underline{\lambda}; c, d, k)$ is Schur-convex in $\underline{\lambda} \in E$, provided that

$$(3.4.7) \quad \left(\frac{\partial}{\partial \lambda_i} - \frac{\partial}{\partial \lambda_j} \right) F_{\underline{\lambda}}(x) \cdot H'(x) f_{\underline{\lambda}}(H(x)) - f_{\underline{\lambda}}(x) \left(\frac{\partial}{\partial \lambda_i} - \frac{\partial}{\partial \lambda_j} \right) F_{\underline{\lambda}}(H(x)) \geq 0,$$

for $i < j, i, j=1, \dots, p$, where $H'(x) = \frac{d}{dx} H(x)$.

Let $\Omega' = \{\underline{\omega} = (\underline{\lambda}_1, \dots, \underline{\lambda}_k) : \underline{\lambda}_i \in E, i=1, \dots, k \text{ and there exist}$

$\lambda_{1m}, \dots, \lambda_{km}$ such that $\lambda_{1m} > \lambda_{2m} > \dots > \lambda_{km}\}$.

Theorem 3.4.2. $\sup_{\Omega'} E(S|R_H)$ takes place at \underline{V} where $\underline{\lambda}_1 = \dots = \underline{\lambda}_k = \underline{V}$ provided that, for $\underline{\lambda}_1 < \underline{\lambda}_2, \underline{\lambda}_1, \underline{\lambda}_2 \in E$,

$$(3.4.8) \quad H'(x) f_{\lambda_2}(H(x)) \left(\frac{\partial}{\partial \lambda_{1i}} - \frac{\partial}{\partial \lambda_{1j}} \right) F_{\lambda_1}(x) - f_{\lambda_2}(x) \left(\frac{\partial}{\partial \lambda_{1i}} - \frac{\partial}{\partial \lambda_{1j}} \right) F_{\lambda_1}(H(x)) \geq 0,$$

for $i < j$, $i, j=1, \dots, p$ where $\lambda_i = (\lambda_{i1}, \dots, \lambda_{ip})$, $i = 1, 2$.

3.5. Selection procedures for multivariate normal distributions in terms of majorization.

Let π_1, \dots, π_k be k populations. Let π_i be associated with $X_i^1 = (X_{i1}, \dots, X_{ip})$, $i = 1, \dots, k$ where X_i^1 is random vector with a p -variate normal distribution with unknown mean vectors $\mu_i = (\mu_{i1}, \dots, \mu_{ip})$ and positive definite covariance matrix Σ . We assume that $\mu_{i1} \geq \dots \geq \mu_{ip}$, $i = 1, \dots, k$.

It is assumed that among the k given populations, there always exists $\mu_{[k]}$ such that $\mu_{[k]} >_{\text{m}} \mu_{[i]}$ for all $i = 1, \dots, k$. This population is called the best population.

Let $X_{i1}^1, \dots, X_{in}^1$ denote n independent observation vectors, each with p components, from population π_i .

Let

$$(3.5.1) \quad X_{i\ell}^1 = (X_{i1\ell}, X_{i2\ell}, \dots, X_{ip\ell}), \quad i = 1, \dots, k, \quad \ell = 1, \dots, n,$$

$$(3.5.2) \quad \bar{X}_{ij} = \frac{1}{n} \sum_{\ell=1}^n X_{ij\ell}, \quad i = 1, \dots, k, \quad j = 1, \dots, p,$$

$$(3.5.3) \quad Y_{ij} = \sum_{\ell=1}^j \bar{X}_{i\ell}, \quad i = 1, \dots, k, \quad j = 1, \dots, p,$$

$$(3.5.4) \quad S_{ij}^2 = \frac{1}{n} \sum_{m=1}^n (X_{ijm} - \bar{X}_{ij})^2, \quad i = 1, \dots, k, \quad j = 1, \dots, p,$$

$$(3.5.5) \quad R_{ij}^2 = \sum_{\ell=1}^j S_{i\ell}^2, \quad i = 1, \dots, k, \quad j = 1, \dots, p,$$

$$(3.5.6) \quad V_j^i = \mu_{i1} + \dots + \mu_{ij}, \quad i = 1, \dots, k, \quad j = 1, \dots, p,$$

$$(3.5.7) \quad \underline{\mu}[i] = (\mu(i)_1, \dots, \mu(i)_p), \quad i = 1, \dots, k, \quad \text{and}$$

$$V_j^{(i)} = \mu(i)_1 + \dots + \mu(i)_j, \quad i = 1, \dots, k, \quad j = 1, \dots, p.$$

Let us denote by $\pi_{(i)}$ the population (unknown) associated with mean vector $\underline{\mu}[i]$ and let $Y_{(i)j}$ be the observation associated with $\pi_{(i)}$, $i = 1, \dots, k$, $j = 1, \dots, p$. Our goal is to select a subset (may be empty) of the k populations so as to include the population associated with $\underline{\mu}[k]$.

(A) Assume Σ is known.

We propose a selection rule as follows:

R_1 : Select population π_i if and only if

$$(3.5.8) \quad Y_{ij} \geq \max_{1 \leq v \leq k} Y_{vj} - d, \quad j = 1, \dots, p-1 \quad \text{and}$$

$$\min_{1 \leq v \leq k} Y_{vp} + d \geq Y_{ip} \geq \max_{1 \leq v \leq k} Y_{vp} - d.$$

Let

$$(3.5.9) \quad Q = D \Sigma^* D'$$

where

$$D_{((k-1)p) \times (kp)} = \begin{pmatrix} A & 0 & \dots & 0 & B \\ 0 & A & & & B \\ 0 & 0 & \ddots & & B \\ \vdots & \vdots & & A & B \\ 0 & 0 & \dots & 0 & A & B \end{pmatrix},$$

$$\Sigma_{(kp) \times (kp)}^* = \begin{pmatrix} T & 0 & 0 \\ 0 & T & \\ 0 & & \dots & T \end{pmatrix}, \quad T_{p \times p} = \frac{1}{n} C \Sigma C'$$

$$A_{p \times p} = \begin{pmatrix} 1 & & 0 \\ \cdot & \cdot & \\ 0 & & 1 \end{pmatrix}, \quad B_{p \times p} = \begin{pmatrix} -1 & & 0 \\ \cdot & \cdot & \\ 0 & & -1 \end{pmatrix} \text{ and } C_{p \times p} = \begin{pmatrix} 1 & & 0 \\ 1 & 1 & \\ \vdots & & \cdot & \cdot & 1 \\ 1 & \dots & 1 & 1 \end{pmatrix}.$$

Let $\Omega = \{\omega = (\underline{\mu}_1, \dots, \underline{\mu}_k) : \underline{\mu}_i \in E_i, i = 1, \dots, k \text{ and there exists some } i \text{ such that } \underline{\mu}_i > \underline{\mu}_j \forall j\}$, where

$$(3.5.10) \quad E_i = \{\underline{\lambda} = (\lambda_1, \dots, \lambda_p) : \lambda_1 \geq \dots \geq \lambda_p, -\infty < \lambda_i < \infty, i=1, \dots, p\}$$

The following theorem gives the infimum of the probability of a correct selection.

Theorem 3.5.1.

$$(3.5.11) \quad \inf_{\Omega} P[CS|R_1] = P[Z_i \leq d, |Z_j| \leq d, \text{ for } i \in \{1, 2, \dots, (k-1)p\} - \{p, 2p, \dots, (k-1)p\} \text{ and } j \in \{p, 2p, \dots, (k-1)p\}],$$

where $(Z_1, \dots, Z_{(k-1)p})' \sim N(\underline{0}, Q)$.

Proof.

$$(3.5.12) \quad P[CS|R_1] = P\{Y_{(k)j} \geq Y_{(v)j} - d, |Y_{(k)p} - Y_{(v)p}| < d, j=1, \dots, p-1, \\ v=1, \dots, k-1\}.$$

Note $Y_{ij} \sim N(v_j^i, \frac{1}{n} c_j' \Sigma c_j)$, $i = 1, \dots, k$, $j=1, \dots, p$, where

$$c_j^i = (\underbrace{1, \dots, 1}_{j \text{ times}}, 0, \dots, 0), \quad j = 1, \dots, p.$$

Since $\mu_{[k]} > \mu_{[i]}$, $i = 1, \dots, k-1$,

$$(3.5.13) \quad \inf_{\Omega} P[CS|R_1] = P\{Y_{kj}^* \geq Y_{vj}^* - d, |Y_{kp}^* - Y_{vp}^*| < d, j=1, \dots, p-1, \\ v=1, \dots, k-1\},$$

where

$$(3.5.14) \quad (Y_{v1}^*, \dots, Y_{vp}^*)' \sim N(\underline{0}, T), \quad v = 1, \dots, k.$$

Let $Y_{-v}^* = (Y_{v1}^*, \dots, Y_{vp}^*)'$, $v = 1, \dots, k$.

Hence from (3.5.14),

$$(3.5.15) \quad \underline{Y} = (Y_{-1}^*, \dots, Y_{-k}^*)' \sim N(\underline{0}, \Sigma^*), \\ k \times 1$$

Let $Z_{vi} = Y_{vi}^* - Y_{ki}^*$, $v = 1, \dots, k-1$, $i = 1, \dots, p$.

Let

$$Z_v^* = (Z_{v1}, \dots, Z_{vp})' \quad v = 1, \dots, k-1.$$

Since

$$\underline{Z} = (Z_1^*, \dots, Z_{k-1}^*)' = D \underline{Y} \text{ and from (3.5.15), then } \underline{Z} \sim N(\underline{0}, Q). \\ (k-1) \times 1$$

This completes the proof.

Theorem 3.5.2. If $Q = D\Sigma^*D' = (q_{ij})$ is positive definite with

$q_{11} = \dots = q_{(k-1)p, (k-1)p} = \sigma^2$ and $q_{ij} = \sigma^2 \rho$ when $i \neq j$, σ and ρ are

known, then

$$(3.5.16) \quad \sup_{\Omega} E(S|R_1) = KP^* \text{ provided that } \inf_{\Omega} P[CS|R_1] = P^*.$$

Proof.

$$(3.5.17) \quad E(S|R_1) = \sum_{i=1}^k P\{Y_{i\ell} \geq Y_{v\ell} - d, Y_{vp} - d \leq Y_{ip} \leq Y_{vp} + d, \\ v \neq i, v = 1, \dots, k, \ell = 1, \dots, p-1\} \\ = \sum_{i=1}^k P\{Z_{i\ell} \leq (V_{\ell}^i - V_{\ell}^v) + d, |Z_{i\ell}| \leq d, \\ v \neq i, v = 1, \dots, k, \ell = 1, \dots, p-1\},$$

where $Z_{i\ell} = (Y_{v\ell} - Y_{i\ell}) - (V_{\ell}^v - V_{\ell}^i)$, $v \neq i, v, i = 1, \dots, k, \ell = 1, \dots, p$.

Thus

$$(3.5.18) \quad \underline{Z}_i = \{Z_{i\ell}\} \sim N(\underline{0}, Q), \\ (k-1)p \times 1$$

Let

$$B_i = \{\underline{Z}_i' : Z_{i\ell} \leq d, |Z_{i\ell}| \leq d, \ell = 1, \dots, p-1, v \neq i, v = 1, \dots, k\}.$$

Let

$$\underline{T}_i = (T_{i\ell}) \text{ where } T_{i\ell} = V_{\ell}^i - V_{\ell}^v, v \neq i, v, i = 1, \dots, k, \ell = 1, \dots, p. \\ 1 \times (k-1)p$$

Hence

$$(3.5.19) \quad E(S|R_1) = \sum_{i=1}^k P[B_i + \underline{T}_i].$$

Since the joint density of \underline{Z} which is defined in (3.5.18) is

Schur-concave (see Marshall and Olkin [50]) and since $\underline{y} \in B_i$ and

$\underline{x} < \underline{y}$ implies $\underline{x} \in B_i$, then by Theorem 2.1 of Marshall and Olkin [50],

$P[B_i + \underline{T}_i]$ is Schur-concave function of \underline{T}_i , $i = 1, \dots, k$.

Since $T_i > (a_i, \dots, a_i)$, $i = 1, \dots, k$ where

$$a_i = \frac{1}{(k-1)p} \sum_{v,l} T_{i,v,l} = \frac{1}{(k-1)p} \{kA_i - \sum_{j=1}^k A_j\} \text{ and}$$

$$A_i = V_1^i + \dots + V_{p-1}^i, \text{ then from (3.5.19)}$$

$$(3.5.20) \quad E(S|R_1) \leq \sum_{i=1}^k P[B_i + (a_i, \dots, a_i)] \\ \leq \sum_{i=1}^k P[B_i + (a_{[k]}, \dots, a_{[k]})]$$

where $a_{[k]}$ is the largest value among $\{a_1, \dots, a_k\}$. That is, $\sup_{\Omega} E(S|R_1)$ is obtained when $a_1 = \dots = a_k$. If $a_1 = \dots = a_k$, then $A_1 = \dots = A_k$. Hence $a_1 = \dots = a_k = 0$.

Thus

$$\sup_{\Omega} E(S|R_1) = \sum_{i=1}^k P[B_i] = k P^* \text{ provided that } \inf_{\Omega} P[CS|R_1] = P^*.$$

This completes the proof.

(B) Let $\Sigma = \begin{pmatrix} \sigma^2 & & 0 \\ & \dots & \\ 0 & & \sigma^2 \end{pmatrix}$ where σ is known. Without loss of generality, we can assume $\sigma = 1$. We propose a subset selection rule as follows.

R_2 : Select π_j if, and only if,

$$(3.5.21) \quad Y_{ij} \geq \max_{1 \leq v \leq k} Y_{vj} - \frac{d\sqrt{2j}}{\sqrt{n}} \text{ for } j = 1, \dots, p-1 \text{ and}$$

$$\max_{1 \leq p < k} Y_{vp} - \frac{d\sqrt{2p}}{\sqrt{n}} \leq Y_{ip} \leq \min_{1 \leq v < k} Y_{vp} + \frac{d\sqrt{2p}}{\sqrt{n}}.$$

Theorem 3.5.3.

$$(3.5.22) \quad \inf_{\Omega} P[CS|R_2] = P[Z_{vi} \leq d, |Z_{vp}| \leq d \text{ for } i=1, \dots, p-1 \\ v=1, \dots, k-1]$$

where Z_{vi} ($v=1, \dots, k-1, i=1, \dots, p$) are standard normal random variables with

$$(3.5.23) \quad \text{Cov}(Z_{\mu i}, Z_{\nu j}) = \begin{cases} \left[\frac{\min(i, j)}{\max(i, j)} \right]^{\frac{1}{2}} & \text{for } \mu = \nu \\ \frac{\min(i, j)}{2\sqrt{ij}} & \text{for } \mu \neq \nu. \end{cases}$$

Proof.

$$(3.5.24) \quad P[CS|R_2] = P\{Y_{(k)i} \geq Y_{(v)i} - \frac{d\sqrt{2i}}{\sqrt{n}},$$

$$Y_{(v)p} - \frac{d\sqrt{2p}}{\sqrt{n}} \leq Y_{(k)p} \leq Y_{(v)p} + \frac{d\sqrt{2p}}{\sqrt{n}},$$

$$i=1, \dots, p-1, v=1, \dots, k-1\}.$$

Let

$$Z_{vi} = \frac{Y_{(v)i} - Y_{(k)i} - (V_i^{(v)} - V_i^{(k)})}{\sqrt{2i}/\sqrt{n}}, \quad v = 1, \dots, k-1, \\ i = 1, \dots, p.$$

Since $(Y_{(v)i} - Y_{(k)i}) \sim N(V_i^{(v)} - V_i^{(k)}, \frac{2i}{n})$,

then $Z_{vi} \sim N(0, 1)$.

Hence

$$(3.5.25) \quad P[CS|R_2] = P\{Z_{vi} \leq d + \frac{V_i^{(k)} - V_i^{(v)}}{\sqrt{2i}/\sqrt{n}}, |Z_{vp}| \leq d, i=1, \dots, p-1, \\ v = 1, \dots, k-1\}.$$

Since $V_i^{(k)} \geq V_i^{(v)}$, $i = 1, \dots, k-1$, then

$$\inf_{\Omega} P[CS|R_2] = P[Z_{vi} \leq d, |Z_{vp}| \leq d, \quad \begin{matrix} i=1, \dots, p-1 \\ v=1, \dots, k-1 \end{matrix}]$$

where Z_{vi} , $v = 1, \dots, k-1$, $i=1, \dots, p$ are $N(0,1)$ random variables with covariances as defined in (3.5.23). This completes the proof.

If $k = 2$ and $p = 2$, we can show that

$$(3.5.26) \quad \inf_{\Omega} P[CS|R_2] = \int_{-\infty}^{\infty} \phi\left(\frac{d - (\frac{1}{2})^{1/4}x}{(1 - \sqrt{\frac{1}{2}})^{1/2}}\right) \left[\phi\left(\frac{d - (\frac{1}{2})^{1/4}x}{(1 - \sqrt{\frac{1}{2}})^{1/2}}\right) - \phi\left(\frac{-d - (\frac{1}{2})^{1/4}x}{(1 - \sqrt{\frac{1}{2}})^{1/2}}\right) \right] d\phi(x)$$

Theorem 3.5.4.

$$(3.5.27) \quad P[CS|R_2] \geq \max\{[\phi(d) - \phi(-d)]^{p(k-1)}, 1 - (p+1)(k-1)\phi(-d)\}.$$

Proof. Let $\{Z_{vi}\}$ be defined as in Theorem 3.5.3, then

$$(3.5.28) \quad P[CS|R_2] \geq P[Z_{vi} \leq d, |Z_{vp}| \leq d, \quad i=1, \dots, p-1, v=1, \dots, k-1].$$

Note that

$$(3.5.29) \quad P\left[\bigcap_{i=1}^m A_i\right] \geq 1 - \sum_{i=1}^m \sum_{j=1}^m P[B_{ij}]$$

where A_1, \dots, A_m denotes a sequence of events, and A_i^c the event complementary to A_i , such that $A_i^c = \bigcup_{j=1}^n B_{ij}$, $i=1, \dots, m$.

Hence from (3.5.28) and (3.5.29),

$$\begin{aligned} (3.5.30) \quad P[CS|R_2] &\geq 1 - \sum_{v,i} P[Z_{vi} > d] - \sum_{v=1}^{k-1} P[|Z_{vp}| > d] \\ &= 1 - (p-1)(k-1)\phi(-d) - (k-1)2\phi(-d) \\ &= 1 - (p+1)(k-1)\phi(-d). \end{aligned}$$

By [68] and from (3.5.28)

$$(3.5.31) \quad P[CS|R_2] \geq P[|Z_{vi}| \leq d, v=1, \dots, k-1, i=1, \dots, p] \\ \geq \prod_{v,i} P[|Z_{vi}| \leq d] = [\Phi(d) - \Phi(-d)]^{p(k-1)}.$$

From (3.5.30) and (3.5.31), we prove the theorem.

For given $P^*(\frac{1}{k} < P^* < 1)$, a conservative value of d ($d > 0$) can be obtained by letting

$$\max\{[\Phi(d) - \Phi(-d)]^{p(k-1)}, 1 - (p+1)(k-1)\Phi(-d)\} = P^*.$$

Let $d_1 > 0$ be the value such that $[\Phi(d) - \Phi(-d)]^{p(k-1)} = P^*$ and let $d_2 > 0$ be the value such that $1 - (p+1)(k-1)\Phi(-d) = P^*$, then the minimum $d(> 0)$ satisfying the basic requirement is given by $d = \min\{d_1, d_2\}$.

Theorem 3.5.5.

$$(3.5.32) \quad E(S|R_2) \leq k[\Phi(d) - \Phi(-d)]$$

Proof.

$$(3.5.33) \quad E(S|R_2) = \sum_{j=1}^k P \left\{ \begin{array}{l} Y_{ji} \geq Y_{vi} - \frac{d\sqrt{2i}}{\sqrt{n}} \\ Y_{vp} - \frac{d\sqrt{2p}}{\sqrt{n}} \leq Y_{jp} \leq Y_{vp} + \frac{d\sqrt{2p}}{\sqrt{n}}, \end{array} \right.$$

for $i=1, \dots, p-1, v \neq j, v=1, \dots, k$ }.

$$\text{Let } Z_{vi} = \frac{Y_{vi} - Y_{ji} - (V_i^v - V_i^j)}{\sqrt{2i}/\sqrt{n}}, \quad i = 1, \dots, p, v \neq j, v=1, \dots, k.$$

Then

$$\begin{aligned}
 (3.5.34) \quad E(S|R_2) &= \sum_{j=1}^k P\{Z_{vi} \leq d + \frac{V_i^j - V_i^v}{\sqrt{2i}/\sqrt{n}}, |Z_{vp}| \leq d, \\
 &\quad i=1, \dots, p-1, v \neq j, v=1, \dots, k\} \\
 &\leq \sum_{j=1}^k P\{Z_{vi} \leq d + \frac{V_i^j - V_i^v}{\sqrt{2i}/\sqrt{n}} \text{ for } v \neq j, v=1, \dots, k\} = \Delta_i \text{ (say),} \\
 &\quad i=1, \dots, p-1.
 \end{aligned}$$

Also,

$$E(S|R_2) \leq \sum_{j=1}^k P[|Z_{vp}| \leq d, v \neq j, v=1, 2, \dots, k] = \Delta \text{ (say).}$$

$$\text{Since } P[Z_{vi} \leq d + \frac{V_i^j - V_i^v}{\sqrt{2i}/\sqrt{n}} \text{ } v \neq j, v=1, \dots, k]$$

$$\leq \frac{1}{k-1} \sum_{\substack{v=1 \\ v \neq j}}^k P[Z_{vi} \leq d + \frac{V_i^j - V_i^v}{\sqrt{2i}/\sqrt{n}}],$$

then

$$\begin{aligned}
 (3.5.35) \quad \Delta_i &\leq \frac{1}{k-1} \sum_{j=1}^k \sum_{\substack{v=1 \\ v \neq j}}^k (d + \frac{V_i^j - V_i^v}{\sqrt{2i}/\sqrt{n}}) \text{ (since } Z_{vi} \sim N(0,1)) \\
 &= \frac{1}{k-1} \sum_{j=1}^k \sum_{\substack{v=1 \\ v \neq j}}^k \phi(d + \frac{A_{[j]i} - A_{[v]i}}{\sqrt{2i}/\sqrt{n}}) \\
 &= \frac{1}{k-1} \cdot Q \text{ (say)}
 \end{aligned}$$

where $A_{[1]i} \leq \dots \leq A_{[k]i}$ are order values of $\{V_i^1, \dots, V_i^k\}$.

By the same argument in Gupta and Huang [34], we can show

that the sup of Q is obtained when $A_{[1]i} = \dots = A_{[k]i}$.

Thus from (3.5.35)

$$(3.5.36) \quad \Delta_i \leq \frac{1}{k-1} \sum_{j=1}^k \sum_{\substack{v=1 \\ v \neq j}}^k \phi(d) = k\phi(d), \quad i=1, \dots, p-1.$$

Also

$$(3.5.37) \quad \Delta \leq \sum_{j=1}^k \frac{1}{k-1} \sum_{\substack{v=1 \\ v \neq j}}^k P[|Z_{vp}| \leq d] \\ = k[\Phi(d) - \Phi(-d)].$$

Hence from (3.5.36) and (3.5.37),

$$(3.5.38) \quad E(S|R_2) \leq \min\{k\Phi(d), k[\Phi(d) - \Phi(-d)]\} \\ = k\{\Phi(d) - \Phi(-d)\}.$$

This completes the proof.

(C) Assume $\Sigma = \begin{pmatrix} \sigma^2 & & 0 \\ & \ddots & \\ 0 & & \sigma^2 \end{pmatrix}$ where σ^2 is unknown.

We propose a subset selection rule as follows.

R_3 : Select population π_j if and only if

$$(3.5.39) \quad Y_{ij} \geq \max_v Y_{vj} - \frac{\sqrt{2} d R_{ij}}{\sqrt{n-1}}, \quad j = 1, \dots, p-1 \text{ and} \\ \max_v Y_{vp} - \frac{\sqrt{2} d R_{ip}}{\sqrt{n-1}} \leq Y_{ip} \leq \min_v Y_{vp} + \frac{\sqrt{2} d R_{ip}}{\sqrt{n-1}}.$$

Theorem 3.5.6.

$$(3.5.40) \quad P[CS|R_3] \geq 1 - (k-1) \sum_{i=1}^{p-1} P[t_i < -d] - (k-1)P[|t_p| > d]$$

where t_i is a r.v. having a t-distribution with $i(n-1)$ d.f., $i=1, \dots, p$.

Proof.

$$(3.5.41) \quad P[CS|R_3] = P[Y_{(k)j} \geq Y_{(v)j} - \frac{\sqrt{2} d R_{(k)j}}{\sqrt{n-1}},$$

$$Y_{(v)p} - \frac{\sqrt{2} d R_{(k)p}}{\sqrt{n-T}} \leq Y_{(k)p} \leq Y_{(v)p} + \frac{\sqrt{2} d R_{(k)p}}{\sqrt{n-T}}, \quad j=1, \dots, p-1, \\ v=1, \dots, k-1]$$

$$= P[t_{vj} \leq d - \frac{v_j^{(k)} - v_j^{(v)}}{\sqrt{2} R_{(k)j}} \cdot \sqrt{n-T}, |t_{vp}| \leq d, j=1, \dots, p-1, v=1, \dots, k-1]$$

where

$$(3.5.42) \quad t_{vj} = \frac{Y_{(v)j} - Y_{(k)j} - (v_j^{(v)} - v_j^{(k)})}{\sqrt{2} R_{(k)j}} \cdot \sqrt{n-T}, \quad j=1, \dots, p, v=1, \dots, k-1.$$

Since $(Y_{(v)j} - Y_{(k)j}) \sim N(v_j^{(v)} - v_j^{(k)}, \frac{2j}{n} \sigma^2)$

and $\frac{nR_{(k)j}^2}{\sigma^2}$ is distributed as a r.v. of χ^2 with $j(n-1)$ d.f., then t_{vj} is a Student's t-distribution with $j(n-1)$ d.f.

Since $v_j^{(k)} \geq v_j^{(v)}, v=1, \dots, k-1, j=1, \dots, p$, then

$$(3.5.43) \quad \inf_{\Omega} P[CS|R_3] = P[t_{vj} \leq d, |t_{vp}| \leq d, j=1, \dots, p-1, v=1, \dots, k-1] \\ \geq 1 - \sum_{v,j} P[t_{vj} \leq d] - \sum_{v=1}^{k-1} P[|t_{vp}| \leq d] \text{ (by (3.5.29))} \\ = 1 - (k-1) \sum_{j=1}^{p-1} P[t_j \leq d] - (k-1) P[|t_p| \leq d]$$

where t_j is a r.v. of a t-distribution with $j(n-1)$ d.f., $j=1, \dots, p$.

This completes the proof.

Theorem 3.5.7.

$$(3.5.44) \quad E(S|R_3) \leq \min\{kG_1(d), k[G_p(d) - G_p(-d)]\}$$

where $G_j(x)$ is the c.d.f. of a r.v. of a t-distribution with $j(n-1)$ d.f., $j=1, \dots, p$.

Proof.

$$(3.4.45) \quad E(S|R_3) = \sum_{i=1}^k P \left\{ \begin{array}{l} Y_{ij} \geq Y_{vj} - \frac{d\sqrt{2} R_{ij}}{\sqrt{n-1}}, \\ Y_{vp} - \frac{d\sqrt{2} R_{ip}}{\sqrt{n-1}} \leq Y_{ip} \leq Y_{vp} + \frac{d\sqrt{2} R_{ip}}{\sqrt{n-1}}, \end{array} \right.$$

for $j = 1, \dots, p-1, v \neq i, v = 1, \dots, k$.

Let $t_{vj} = \frac{Y_{vj} - Y_{ij} - (V_j^v - V_j^i)}{\sqrt{2} R_{ij}} \cdot \sqrt{n-1}$, then t_{vj} is a Student's t -distribution with $j(n-1)$ d.f.

$$(3.5.46) \quad E(S|R_3) = \sum_{i=1}^k P \left\{ \begin{array}{l} t_{vj} \leq d - \frac{V_j^v - V_j^i}{\sqrt{2} R_{ij}} \sqrt{n-1}, |t_{vp}| \leq d, \\ j = 1, \dots, p-1, v \neq i, v = 1, \dots, k \end{array} \right\}.$$

Hence

$$(3.5.47) \quad E(S|R_3) \leq \sum_{i=1}^k P\{t_{vj} \leq d + \frac{V_j^i - V_j^v}{\sqrt{2} R_{ij}} \sqrt{n-1}, v \neq i, v = 1, \dots, k\}$$

$$= \Delta_j \text{ (say), } j = 1, \dots, p-1.$$

and

$$(3.5.48) \quad E(S|R_3) \leq \sum_{i=1}^k P\{|t_{vp}| \leq d, v \neq i, v = 1, \dots, k\}$$

$$= \Delta \text{ (say).}$$

As in Theorem 3.5.5, we can show that the sup of Δ_j is obtained when $V_j^i = \dots = V_j^k$, for $j = 1, \dots, p-1$. From (3.5.47) and (3.5.48), we get $\Delta_j \leq kG_j(d)$, $j=1, \dots, p-1$ and $\Delta \leq k[G_p(d) - G_p(-d)]$. Thus

$$E(S|R_3) \leq \min\{k[G_p(d) - G_p(-d)], kG_j(d), j=1, \dots, p-1\}$$

$$= \min\{k[G_p(d) - G_p(-d)], kG_1(d)\}$$

3.6. Selection procedures for multivariate normal distributions in terms of weak majorization

It is assumed that among the k populations, there always exists $\underline{\mu}[k]$ such that $\underline{\mu}[k] \gg_m \underline{\mu}[i]$, $i = 1, \dots, k$. This population is called the best population. We are using the same notation as in Section 3.5. Let $\Omega = \{\underline{\omega} \in (\underline{\mu}_1, \dots, \underline{\mu}_k) : \underline{\mu}_i \in E_1, i=1, \dots, k \text{ and there exists some } j \text{ such that } \underline{\mu}_j \gg_m \underline{\mu}_i \forall i\}$ where E_1 is defined in (3.5.10).

(A) Assume Σ is known.

We propose a selection rule as follows.

R_5 : Select population π_j if and only if

$$(3.6.1.) \quad Y_{ij} \geq \max_{1 \leq v \leq k} Y_{vj} - d, \quad j = 1, \dots, p.$$

Using the same argument as in Theorem 3.5.1, we obtain the following result.

Theorem 3.6.1.

$$(3.6.2) \quad \inf_{\Omega} P[CS|R_5] = P[Z_i \leq d, i = 1, \dots, (k-1)p]$$

where

$$(Z_1, \dots, Z_{(k-1)p})' \sim N\left(\begin{matrix} \underline{0} \\ (k-1)p \times 1 \end{matrix}, Q\right), \text{ where } Q \text{ is defined in (3.5.9).}$$

(B) Assume $\Sigma = \begin{pmatrix} \sigma^2 & & 0 \\ & \ddots & \\ 0 & & \sigma^2 \end{pmatrix}$ where σ is known and we assume $\sigma = 1$.

We propose a selection rule as follows.

R_6 : Select population π_j if and only if,

$$(3.6.3.) \quad Y_{ij} \geq \max_{1 \leq v \leq k} Y_{vj} - \frac{d\sqrt{2j}}{\sqrt{n}}, \quad j=1, \dots, p.$$

It is similar as in Theorem 3.5.3, we can show that

Theorem 3.6.2.

$$(3.6.4) \quad \inf_{\Omega} P[CS|R_6] = P[Z_{vj} \leq d, \quad v=1, \dots, k-1, j=1, \dots, p]$$

where Z_{vj} is defined in Theorem 3.5.3.

If $k = 2, p = 2$, we can show that

$$(3.6.5) \quad \inf_{\Omega} P[CS|R_6] = \int_{\Phi}^2 \left[\frac{d - (\frac{1}{2})^{1/4} x}{(1 - \sqrt{\frac{T}{2}})^{1/2}} \right] d\Phi(x).$$

In a same manner as in Theorem 3.5.4 and Theorem 3.5.5, we have the following results.

Theorem 3.6.3.

$$(3.6.6) \quad P[CS|R_6] \geq \max\{[\Phi(d) - \Phi(-d)]^{p(k-1)}, 1 - p(k-1)[1 - \Phi(d)]\}.$$

Theorem 3.6.4.

$$(3.6.7) \quad E(S|R_6) \leq k\Phi(d).$$

(C) Assume $\Sigma = \begin{pmatrix} \sigma^2 & & 0 \\ & \dots & \\ 0 & & \sigma^2 \end{pmatrix}$ where σ^2 is unknown.

We propose a subset selection rule as follows.

R_7 : Select population π_i if and only if

$$(3.6.8) \quad Y_{ij} \geq \max_v Y_{vj} - \frac{\sqrt{2} d R_{ij}}{\sqrt{n-1}}, \quad j = 1, \dots, p.$$

In a manner similar to that in the proof of Theorem 3.5.6, we have

Theorem 3.6.5.

$$(3.6.9) \quad P[CS|R_7] \geq 1 - (k-1) \sum_{i=1}^p P[t_i \leq -d]$$

where t_i is the r.v. of Student's t -distribution with $i(n-1)$ d.f., $i = 1, \dots, p$.

BIBLIOGRAPHY

- [1] Bahadur, R. R. (1950). On the problem in the theory of k populations. Ann. Math. Statist., 21, 362-375.
- [2] Barlow, R. E., Bartholomew, D. J., Bremner, J. M. and Brunk, H. D. (1972). Statistical Inference under Order Restrictions. John Wiley, New York.
- [3] Barlow, R. E. and Doksum, K. A. (1972). Isotonic tests for convex orderings. Proc. Sixth Berkeley Symp. on Math. Statist. and Prob. Vol. I, 293-323.
- [4] Barlow, R. E., Esary, J. D. and Marshall, A. W. (1966). A stochastic characterization of wear-out for components and systems. Ann. Math. Statist., 37, 816-825.
- [5] Barlow, R. E. and Gupta, S. S. (1969). Selection procedures for restricted families of distributions. Ann. Math. Statist., 40, 905-917.
- [6] Barlow, R. E., Gupta, S. S. and Panchapakesan, S. (1969). On the distribution of the maximum and minimum of ratios of order statistics. Ann. Math. Statist., 40, 918-934.
- [7] Barlow, R. E., Marshall, A. W. and Proschan, F. (1963). Properties of probability distributions with monotone hazard rate. Ann. Math. Statist., 34, 375-389.
- [8] Barlow, R. E. and Proschan, F. (1966). Inequalities for linear combinations of order statistics from restricted families. Ann. Math. Statist., 37, 1574-1592.
- [9] Barlow, R. E. and Proschan, F. (1966). Tolerance and confidence limits for classes of distributions based on failure rate. Ann. Math. Statist., 37, 1593-1601.
- [10] Barlow, R. E. and Proschan, F. (1969). A note on tests for monotone failure rate based on incomplete data. Ann. Math. Statist., 40, 595-600.


- [11] Barlow, R. E. and Van Zwet, W. R. (1970). Asymptotic properties of isotonic estimators for the generalized failure rate function. Part I: strong consistency. Nonparametric Techniques in Statistical Inference (ed. Puri, M. L.) 159-173. Cambridge University Press, London.
- [12] Barr, D. R. and Rizvi, M. H. (1966). An introduction to ranking and selection procedures. J. Amer. Statist. Assoc., 61, 640-646.
- [13] Bechhofer, R. E. (1954). A single-sample multiple decision procedure for ranking means of normal populations with known variances. Ann. Math. Statist., 25, 16-39.
- [14] Bechhofer, R. E., Dunnett, C. W., and Sobel, M. (1954). A two-sample multiple-decision procedure for ranking means of normal populations with a common unknown variance. Biometrika, 41, 170-176.
- [15] Bechhofer, R. E., Kiefer, J. and Sobel, M. (1968). Sequential Identification and Ranking Procedures. The University of Chicago Press, Chicago.
- [16] Bechhofer, R. E. and Sobel, M. (1954). A single-sample multiple-decision procedure for ranking variances of normal populations. Ann. Math. Statist., 25, 273-289.
- [17] Beckenbach, E. F. and Bellman, R. L. (1961). Inequalities. Springer-Verlag, Berlin.
- [18] Beckenbach, E. F. and Bellman, R. L. (1965). Inequalities. Springer-Verlag, Berlin.
- [19] Carroll, R. J., Gupta, S. S., and Huang, D. Y. (1975). On selection procedures for the t best populations and some related problems. Comm. Statist., 4, 987-1008.
- [20] Chen, H. J. and Dudewicz, E. J. (1973). Estimation of ordered parameters and subset selection. Tech. Report, Dept. of Statistics, Ohio State University, Columbus, Ohio 43210.
- [21] Desu, M. M. (1970). A selection problem. Ann. Math. Statist., 41, 1596-1603.
- [22] Desu, M. M. and Sobel, M. (1971). Nonparametric procedures for selecting fixed size subsets. Statistical Decision Theory and Related Topics (ed. Gupta, S. S. and Yackel, J.) 255-273. Academic Press, N.Y.

- [23] Doksum, K. (1969). Star-shaped transformations and the power of rank tests. Ann. Math. Statist., 40, 1167-1176.
- [24] Dudewicz, E. J. (1970). Confidence intervals for ranked means. Naval. Res. Logist. Quart., 17, 69-78.
- [25] Dudewicz, E. J. (1972). Two-sided confidence intervals for ranked means. J. Amer. Statist. Assoc., 67, 462-464.
- [26] Dudewicz, E. J. and Tong, Y. L. (1971). Optimum confidence intervals for the largest location parameter. Statistical Decision Theory and Related Topics (Ed. S. S. Gupta and J. Yackel), Academic Press, New York, pp. 363-376.
- [27] Gnanadesikan, M. (1966). Some Selection and Ranking Procedures for Multivariate Normal Populations. Ph.D. Thesis. Dept. of Statist., Purdue Univ., West Lafayette, Indiana 47907.
- [28] Gnanadesikan, M. and Gupta, S. S. (1970). Selection procedures for multivariate normal distributions in terms of measures of dispersion. Technometrics, 12, 103-130.
- [29] Gupta, S. S. (1956). On a decision rule for a problem in ranking means. Mimeo. Ser. No. 150. Inst. of Statist., Univ. of North Carolina, Chapel Hill, North Carolina.
- [30] Gupta, S. S. (1963). On a selection and ranking procedure for gamma populations. Ann. Inst. Statist. Math., 14, 199-216.
- [31] Gupta, S. S. (1965). On some multiple decision (selection and ranking) rules. Technometrics, 7, 225-245.
- [32] Gupta, S. S. (1966). On some selection and ranking procedures for multivariate normal populations using distance functions. Multivariate Analysis (ed. Krishnaiah, P. R.) 457-475. Academic Press, N.Y.
- [33] Gupta, S. S. and Huang, D. Y. (1975). Subset selection procedures in analysis of variance. Mimeo. Ser. No. 436. Dept. of Statist., Purdue Univ., West Lafayette, Indiana 47907.
- [34] Gupta, S. S. and Huang, D. Y. (1976). Selection procedures for the means and variances of normal populations when the sample sizes are unequal. Sankhyā.
- [35] Gupta, S. S. and McDonald, G. C. (1970). On some classes of selection procedures based on ranks. Nonparametric Techniques in Statistical Inference (ed. Puri, M. L.) 419-514. Cambridge University Press, London.

- [36] Gupta, S. S. and Panchapakesan, S. (1969). Selection and ranking procedures. The Design of Computer Simulation Experiments (ed. Naylor, T. H.) 132-160. Duke University Press, Durham, North Carolina.
- [37] Gupta, S. S. and Panchapakesan, S. (1972). On a class of subset selection procedures. Ann. Math. Statist., 43, 814-822.
- [38] Gupta, S. S. and Panchapakesan, S. (1972). On multiple decision (subset selection) procedures. J. Mathematical and Physical Sci., 6, 1-72.
- [39] Gupta, S. S. and Panchapakesan, S. (1974). Inference for restricted families: (a) multiple decision procedures (b) order statistics inequalities. Reliability and Biometry (Ed. F. Proschan and R. J. Serfling), SIAM, Philadelphia, pp. 503-596.
- [40] Gupta, S. S. and Panchapakesan, S. (1975). On a quantile selection procedure and associated distribution of ratios of order statistics from a restricted family of probability distributions. Reliability and Fault Tree Analysis (Ed. R. E. Barlow, J. B. Fussell, and N. D. Singpurwalla), SIAM, Philadelphia, pp. 557-576.
- [41] Gupta, S. S. and Santner, T. J. (1973). On selection and ranking procedures - a restricted subset selection rule. Contributed Papers, Vol. I (International Statistical Institute, 39th Session) pp. 409-417.
- [42] Gupta, S. S. and Sobel, M. (1958). On the distribution of a statistic based on ordered uniform chance variables. Ann. Math. Statist., 29, 274-281.
- [43] Gupta, S. S. and Studden, W. J. (1970). On some selection and ranking procedures with applications to multivariate populations. Essays in Probability and Statistics (Ed. R. C. Bose et al.), University of North Carolina Press, Chapel Hill, North Carolina, pp. 327-338.
- [44] Gupta, S. S. and Wong, W. Y. (1975). Subset selection procedures for finite schemes in information theory. Mimeo. Ser. No. 422. Dept. of Statist., Purdue Univ., West Lafayette, Indiana 47907.
- [45] Hardy, G. H., Littlewood, J. E. and Pólya, G. (1959). Inequalities. Cambridge University Press, London.
- [46] Huang, D. Y. (1974). Some Contributions to Fixed Sample and Sequential Multiple Decision (Selection and Ranking) Theory. Ph.D. Thesis (Mimeo. Ser. No. 363). Dept. of Statist., Purdue Univ., West Lafayette, Indiana 47907.

- [47] Lawrence, M. J. (1966). Inequalities and tolerance limits for s-ordered distributions. Rep. ORC 66-37, Operations Research Center, Univ. of California, Berkeley, California.
- [48] Lawrence, M. J. (1975). Inequalities of s-ordered Distributions. Ann. Statist., 3, 413-428.
- [49] Lehmann, E. L. [1959]. Testing Statistical Hypotheses. John Wiley, New York.
- [50] Marshall, A. W. and Olkin, I. (1974). Majorization in multivariate distributions. Ann. Statist., 2, 1189-1200.
- [51] Marshall, A. W., Olkin, I. and Proschan, F. (1967). Monotonicity of ratios of means and other applications of majorization. Inequalities (ed. Shisha, O.) 177-190, Academic Press, N.Y.
- [52] Marshall, A. W. and Proschan, F. (1965). An inequality for convex functions involving majorization. J. Math. Anal. Appl., 12, 87-90.
- [53] Moore, D. S. (1968). An elementary proof of asymptotic normality of linear functions of order statistics. Ann. Math. Statist., 39, 263-265.
- [54] Mosteller, F. (1948). A k-sample slippage test for an extreme population. Ann. Math. Statist., 19, 58-65.
- [55] Nevius, S. E., Proschan, F. and Sethuraman, J. [1974]. Schur functions in statistics. II: Stochastic majorization. Tech. Report M306. Dept. of Statistics, Florida State Univ., Tallahassee, Florida.
- [56] Nevius, S. E., Proschan, F. and Sethuraman, J. [1974]. Schur functions in statistics. III: A stochastic version of weak majorization, with applications. To appear in Statistical Decision Theory and Related Topics-II (ed. Gupta, S. S. and Moore, D. S.), Academic Press, 1976.
- [57] Ostrowski, A. (1952). Sur quelques applications des fonctions convexes et concaves au sens de I. Schur. J. Math. Pure Appl., 31, 253-292.
- [58] Patel, J. K. (1976). Ranking and selection of IFR populations based on means. J. Amer. Statist. Assoc., 71, 143-146.
- [59] Paulson, E. (1949). A multiple decision procedure for certain problem in analysis of variance. Ann. Math. Statist., 20, 95-98.

- [60] Pledger, G. and Proschan, F. (1973). Stochastic comparisons of random processes, with applications in reliability. J. Appl. Prob. 10, 572-585.
- [61] Proschan, F. and Sethuraman, J. [1974]. Schur functions in statistics. I: The preservation theorem. Tech. Report M254-RR. Dept. of Statist., Florida State Univ., Tallahassee, Florida.
- [62] Rinott, Y. (1973). Multivariate majorization and rearrangement inequalities with some applications to probabilities and statistics. Israel J. Math., 15, 60-77.
- [63] Rizvi, M. H. and Sobel, M. (1967). Nonparametric procedures for selecting a subset containing the population with the largest α -quantile. Ann. Math. Statist., 38, 1788-1803.
- [64] Santner, T. J. (1975). A restricted subset selection approach to ranking and selection problems. Ann. Statist., 3, 334-349.
- [65] Saxena, K. M. Lal and Tong, Y. L. (1969). Interval estimation of the largest mean of k normal populations with known variances. J. Amer. Statist. Assoc., 64, 296-299.
- [66] Saxena, K. M. Lal and Tong, Y. L. (1971). Optimum interval estimation for the largest scale parameter (Abstract). Optimizing Methods in Statistics (ed. Rustagi, J. S.) 477-477. Academic Press, New York and London.
- [67] Schur, I. (1923). Über eine Klasse von Mittelbildungen mit Anwendungen auf die Determinantentheorie. Sitzber. Berl. Math. Ges., 22, 9-20.
- [68] Sidák, Z. (1967). Rectangular confidence regions for the means of multivariate normal distributions. J. Amer. Statist. Assoc., 62, 626-633.
- [69] Slepian, D. (1962). On the one-sided barrier problem for gaussian noise. Bell System Tech. J., 41, 463-501.
- [70] Sobel, M. and Huyett, M. J. (1957). Selecting the best one of several binomial populations. Bell System Tech. J., 36, 537-576.
- [71] Van Zwet, W. R. (1964). Convex Transformations of Random Variables. Mathematical Center, Amsterdam.
- [72] Wong, W. Y. (1976). Some contributions to subset selection procedures. Ph.D. Thesis, Dept. of Statistics, Purdue University, West Lafayette, Indiana 47907.



Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Mimeograph Series #464	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) On Some Multiple Decision Problems.		5. TYPE OF REPORT & PERIOD COVERED Technical
		6. PERFORMING ORG. REPORT NUMBER Mimeo. Series #464
7. AUTHOR(s) Ming-Wei Lu	8. CONTRACT OR GRANT NUMBER(s) N00014-67-A-0226-0014	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Purdue University Department of Statistics, W. Lafayette, IN 47907		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 042-243
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Washington, DC		12. REPORT DATE August 1976
		13. NUMBER OF PAGES 119
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		18. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release, distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Selection procedures, IFR distributions, estimations, good populations, majorization, Schur functions, multivariate normal.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Chapter I of this thesis deals with some selection and ranking procedures for restricted families of probability distributions. A selection rule is proposed for distributions which are convex-ordered with respect to a specified distribution G. Some properties of this selection rule are derived. The asymptotic relative efficiencies of this rule with respect to other selection rules are evaluated. A selection rule is also proposed and studied for distributions which are s-ordered with respect to G. In Chapter II, some interval estimation		

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

problems for the unknown parameters of the k populations are studied. The infimum of the probability that a given confidence interval (based on suitably chosen order statistics) contains at least one "good" population is obtained. Different modifications and variations of this problem are also studied. The last section of Chapter II illustrates some results by means of two examples. Chapter III discusses the selection procedures in terms of majorization and weak majorization. The parameter is partially ordered by means of majorization or weak majorization. A class of procedures R_n for selecting the best population is defined. A sufficient condition is obtained for the infimum of the probability of a correct selection to be Schur-convex in λ . Also another sufficient condition for the same infimum of the probability of a correct selection to be nondecreasing and Schur-convex in λ is obtained. We also propose and study the selection procedures for multivariate normal distributions in terms of majorization and weak majorization. Various cases corresponding to the known or unknown common covariance matrix Σ are studied.