

ON $(n-1)$ -WISE AND JOINT INDEPENDENCE AND
NORMALITY OF n RANDOM VARIABLES: AN EXAMPLE

by

Wolfgang J. Bühler and Klaus J. Miescke
Mainz University

Department of Statistics
Division of Mathematical Sciences
Mimeograph Series #80-27

September 1980

*This research was supported by the Office of Naval Research under Contract
N00014-75-C-0455 at Purdue University. Reproduction in whole or in part is
permitted for any purpose of the United States Government.

ON $(n-1)$ -WISE AND JOINT INDEPENDENCE AND
NORMALITY OF n RANDOM VARIABLES: AN EXAMPLE

Wolfgang J. Bühler and Klaus J. Miescke

Mainz University

ABSTRACT

An example is given of a vector of n random variables such that any $(n-1)$ -dimensional subvector consists of $n-1$ independent standard normal variables. The whole vector however is neither independent nor normal.

1. INTRODUCTION

When discussing stochastic independence in a course on probability theory, it is customary to give an example of three identically distributed random variables X , Y and Z which are pairwise independent but not mutually independent. As Driscoll (1978) has pointed out, the standard examples (Feller (1957); Gnedenko (1963); DeGroot (1975); Hogg and Craig (1970)) can be reduced to consideration of a random triple (X, Y, Z) which takes the values $(0,0,0)$, $(0,1,1)$, $(1,0,1)$ and $(1,1,0)$ each with probability one-fourth.

Driscoll gave a more interesting example: \tilde{X} , \tilde{Y} independent each with the rectangular distribution on the unit interval and

$\tilde{Z} = \tilde{X} + \tilde{Y} \text{ mod } 1$. This example also yielded a characterization of the rectangular distribution.

Our example shares with Driscoll's the fact of being more interesting than the standard ones and at the same time illustrates a point concerning the multi-dimensional normal distribution.

It is well known that the whole distribution of an n -dimensional normal vector (X_1, X_2, \dots, X_n) is determined if the distribution of each pair (X_i, X_j) is known. In a different context one of the authors (KJM) raised the question whether (X_1, X_2, \dots, X_n) is necessarily normal if all the pairs (X_i, X_j) are two-dimensional normal vectors. The following example shows that even joint normality of all $(n-1)$ -tuples does not suffice.

2. THE EXAMPLE

Let $n \geq 3$ and let (Y_1, Y_2, \dots, Y_n) be a random vector of signs, i.e. with components $+1$ or -1 such that any particular sign vector (y_1, y_2, \dots, y_n) is taken with probability a if $\prod_{i=1}^n y_i = +1$ and with probability $b = 2^{-(n-1)} - a$ if $\prod_{i=1}^n y_i = -1$. Here $0 \leq a \leq 2^{-(n-1)}$.

Proposition: The random variables Y_1, Y_2, \dots, Y_n are $(n-1)$ -wise independent. If $a \neq 2^{-n}$ they are not mutually independent.

Proof: Let $1 \leq k \leq n-1$. Any vector $(y_{i_1}, y_{i_2}, \dots, y_{i_k})$ can then be extended in 2^{n-k-1} ways to a vector (y_1, y_2, \dots, y_n) with $\prod_{i=1}^n y_i = +1$ and in as many ways to one for which the product of its components is -1 . Thus $P(Y_{i_1} = y_{i_1}, Y_{i_2} = y_{i_2}, \dots, Y_{i_k} = y_{i_k}) = 2^{n-k-1} a + 2^{n-k-1} b = 2^{-k}$ for all $k \leq n-1$. This is the $(n-1)$ -wise independence. However the relation $P(Y_1 = +1, Y_2 = +1, \dots, Y_n = +1) = a$ contradicts the total independence unless $a = 2^{-n}$.

Now let Z_1, Z_2, \dots, Z_n be standard normal variables mutually independent and independent of the random vector (Y_1, Y_2, \dots, Y_n)

and define $X_i = Y_i/|Z_i|$, $i = 1, \dots, n$. Then clearly the X_i are again standard normal. Also the independence of the Z_i together with the proposition imply that X_1, X_2, \dots, X_n are $(n-1)$ -wise independent. Thus any $(n-1)$ -tuple out of X_1, X_2, \dots, X_n is also $(n-1)$ -dimensional normal. However $P(X_1 > 0, X_2 > 0, \dots, X_n > 0) = P(Y_1 = Y_2 = \dots = Y_n = +1) = a$ which, if $a \neq 2^{-n}$, contradicts the mutual independence of X_1, X_2, \dots, X_n and thus also their joint normality, where mutual independence would be equivalent to all covariances being zero.

3. REMARKS

The example does in no way characterize the normal distribution. In fact we can replace the normal distribution of the Z_i by any other distribution symmetric around zero to obtain a similar example where all subvectors of (Z_1, \dots, Z_n) except (Z_1, \dots, Z_n) itself consist of mutually independent identically distributed random variables. With $n = 3$ and $a = 0$ the vector $2^{-1}(Y_1 + 1, Y_2 + 1, Y_3 + 1)$ is the random triple (X, Y, Z) mentioned in the introduction.

ACKNOWLEDGEMENT

This research was partly supported by the Office of Naval Research Contract N00014-75-C-0455 at Purdue University.

BIBLIOGRAPHY

- DeGroot, Morris H. (1975). *Probability and Statistics*. Addison-Wesley Publishing Co., Reading, Massachusetts.
- Driscoll, Michael F. (1978). On pairwise and mutual independence: Characterizations of rectangular distributions. *Journal of the American Statistical Association*, 73, 432-3.
- Feller, William (1957). *Introduction to Probability Theory and Its Applications*. 2nd ed., John Wiley & Sons, New York.
- Gnedenko, B. V. (1963). *The Theory of Probability*. 2nd ed., Chelsea Publishing Co., New York.
- Hogg, Robert V. and Craig, Allen T. (1970). *Introduction to Mathematical Statistics*. 3rd ed., The MacMillan Co., New York.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Mimeograph Seres #80- 27	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ON (n-1)-WISE AND JOINT INDEPENDENCE AND NORMALITY OF n RANDOM VARIABLES: AN EXAMPLE		5. TYPE OF REPORT & PERIOD COVERED Technical
		6. PERFORMING ORG. REPORT NUMBER Mimeo. Series #80-27
7. AUTHOR(s) Wolfgang J. Bühler and Klaus J. Miescke		8. CONTRACT OR GRANT NUMBER(s) ONR N00014-75-C-0455
9. PERFORMING ORGANIZATION NAME AND ADDRESS Purdue University Department of Statistics West Lafayette, IN 47907		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Washington, DC		12. REPORT DATE September, 1980
		13. NUMBER OF PAGES 3
14. MONITORING AGENCY NAME & ADDRESS (If different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release, distribution unlimited.		
17. DISTRIBUTION STATEMENT (of this abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Independence; (n-1)-wise independence; joint normality.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) An example is given of a vector of n random variables such that any (n-1)- dimensional subvector consists of n-1 independent standard normal variables. The whole vector however is neither independent nor normal.		