

QUADRATIC COVARIATION
AND AN EXTENSION OF ITO'S FORMULA

by

H. Föllmer P. Protter
Universität Bonn Purdue University

and

A.N. Shiryaev
Steklov Mathematical Institute

Technical Report #94-34

Department of Statistics
Purdue University

November 1994

**Quadratic Covariation
and an Extension of Itô's Formula**

H. Föllmer, Ph. Protter, A.N. Shiryaev

Abstract. Let X be a standard Brownian motion. We show that for any locally square integrable function f the quadratic covariation $[f(X), X]$ exists as the usual limit of sums converging in probability. For an absolutely continuous function F with derivative f , Itô's formula takes the form $F(X_t) = F(X_0) + \int_0^t f(X_s) dX_s + \frac{1}{2}[f(X), X]_t$. This is extended to the time dependent case. As an example, we introduce the local time of Brownian motion at a continuous curve.

To appear in: *Bernoulli*, Journal of the Bernoulli Society

*Institut für angewandte Mathematik, Universität Bonn, Wegelerstrasse 6,
53115 Bonn, Germany*

*Mathematics and Statistics Departments, Purdue University,
West Lafayette, IN 47907-1395, USA*

*Steklov Mathematical Institute, Vavilova str. 42
Moscow 117966 GSP-1, Russia*

1. Introduction.

Let $X = (X_t)_{0 \leq t \leq 1}$ be a standard Brownian motion, and let F be an absolutely continuous function with locally square integrable derivative f . Our purpose is to prove the following extension of Itô's formula:

$$(1.1) \quad F(X_t) = F(X_0) + \int_0^t f(X_s) dX_s + \frac{1}{2} [f(X), X]_t.$$

where

$$(1.2) \quad [f(X), X]_t = \lim_{n \rightarrow \infty} \sum_{t_i \in D_n, t_i < t} \{f(X_{t_{i+1}}) - f(X_{t_i})\} (X_{t_{i+1}} - X_{t_i}).$$

denotes the *quadratic covariation* of the processes $f(X)$ and X . In particular we are going to show that the quadratic covariation exists for any locally square integrable function f , as a limit in probability along a sequence of partitions D_n of the time interval $[0,1]$.

If f is absolutely continuous with derivative f' then the quadratic covariation is given by

$$(1.3) \quad [f(X), X]_t = \int_0^t f'(X_s) ds,$$

and so equation (1.1) reduces to Itô's formula in its classical form. If f is locally bounded then we have

$$(1.4) \quad [f(X), X]_t = - \int f(a) d_a L_t^a$$

where L_t^a denotes the local time of Brownian motion at level a , and so (1.1) reduces to the formula of Bouleau and Yor [1].

The quadratic covariation in (1.2) admits the representation

$$(1.5) \quad [f(X), X]_t = \int_0^t f(X_s) d^* X_s - \int_0^t f(X_s) dX_s.$$

in terms of a backward and a forward stochastic integral, and this is the key to our existence result. In fact, the existence of quadratic covariation will follow from an approximation of these integrals by forward and backward sums. In section 2 we motivate our approach

by looking at the discrete version of (1.1) along a fixed partition. In section 3 we prove the basic existence result for the quadratic covariation $[f(X, \cdot), X]$ in the general time dependent case where $f(x, t)$ is locally square integrable in x and satisfies a mild continuity condition in t . In general, the quadratic covariation will not be of bounded variation. But we show that it is always a process of *zero energy*, i.e., a process with continuous paths of zero quadratic variation. In section 4 we derive our Itô formula (1.1). Thus, the process of zero energy appearing in Fukushima's decomposition of the Dirichlet process $F(X)$ ([4] Ch. 5) is identified as a quadratic covariation. Our formula also shows that the Stratonovich integral can be defined on the same level of generality as the Itô integral, without additional restrictions on the function f . In section 5 we extend these results to the time dependent case where $F(x, t)$ is absolutely continuous in x and where the derivative $f(x, t)$ satisfies our conditions for the existence of $[f(X, \cdot), X]$. In this case the Itô formula takes the form

$$(1.6) \quad F(X_t, t) = F(X_0, 0) + \int_0^t f(X_s, s) dX_s + \frac{1}{2} [f(X, \cdot), X]_t + \int_0^t F(X_s, ds)$$

where

$$(1.7) \quad \int_0^t F(X_s, ds) \equiv \lim_{n \rightarrow \infty} \sum_{t_i \in D_n, t_i < t} F(X_{t_{i+1}}, t_{i+1}) - F(X_{t_i}, t_i)$$

exists as a limit in probability. If $F(x, \cdot)$ is absolutely continuous in t with derivative $F_t(x, \cdot)$ then the last term in (1.6) takes the usual form

$$(1.8) \quad \int_0^t F(X_s, ds) = \int_0^t F_t(X_s, s) ds.$$

In the case where $f(x, t) = I_{[a(t), \infty)}(x)$ for some continuous function $a(\cdot)$, the existence of the quadratic covariation $[f(X, \cdot), X]$ amounts to a construction of the local time of Brownian motion at a continuous curve, and (1.6) may be viewed as a time dependent version of the Tanaka formula.

The idea of using time reversal and the duality (1.5) for extended versions of the Stratonovich integral and of Itô's formula also appears in recent work of Lyons and Zhang

[7] and of Russo and Vallois [9]. Section 5 shows that, in contrast to the Dirichlet space techniques used in [7] and [4], our approach admits a straightforward extension to the time dependent case. Extensions to higher dimensions and applications to the existence of solutions of Stratonovich stochastic differential equations will be discussed elsewhere.

2. Preliminaries

Let F be an absolutely continuous function of the form

$$(2.1) \quad F(x) = F(0) + \int_0^x f(y)dy$$

where f is locally square integrable. In order to motivate our approach, let us first see how the structure of our Itô formula (1.1) appears in discrete time. Let D_n denote a partition of the form $0 = t_0 < \dots < t_k = 1$ and let us write $|D_n| = \max |t_{i+1} - t_i|$. An increment

$$(2.2) \quad F(X_{t_{i+1}}) - F(X_{t_i}) = \int_{X_{t_i}}^{X_{t_{i+1}}} f(y)dy$$

can be written as

$$(2.3) \quad f(X_{t_i})(X_{t_{i+1}} - X_{t_i}) + \int_{X_{t_i}}^{X_{t_{i+1}}} (f(y) - f(X_{t_i}))dy$$

and also as

$$(2.4) \quad f(X_{t_i})(X_{t_{i+1}} - X_{t_i}) + \{f(X_{t_{i+1}}) - f(X_{t_i})\}(X_{t_{i+1}} - X_{t_i}) + \int_{X_{t_i}}^{X_{t_{i+1}}} (f(y) - f(X_{t_{i+1}}))dy.$$

Averaging both expressions and summing the increments up to time $t \in D_n$, we obtain

$$(2.5) \quad F(X_t) - F(X_0) = \sum_{\substack{t_i \in D_n \\ t_i < t}} f(X_{t_i})(X_{t_{i+1}} - X_{t_i}) + \frac{1}{2}Q_t^n + R_t^n$$

with discrete quadratic covariation

$$(2.6) \quad Q_t^n \equiv \sum_{\substack{t_i \in D_n \\ t_i < t}} \{f(X_{t_{i+1}}) - f(X_{t_i})\}(X_{t_{i+1}} - X_{t_i})$$

and remainder terms

$$(2.7) \quad R_t^n \equiv \sum_{\substack{t_i \in D_n \\ t_i < t}} \int_{X_{t_i}}^{X_{t_{i+1}}} \left\{ f(y) - \frac{1}{2}(f(X_{t_i}) + f(X_{t_{i+1}})) \right\} dy$$

The first sum in (2.5) is the discrete analogue of the stochastic integral in (1.1), and Q_t^n is the discrete version of the quadratic covariation in (1.1). Let us now pass to the limit

along a sequence of partitions D_n such that $|D_n|$ converges to 0. Let us say that a sequence of processes $Y^n = (Y_t^n)_{0 \leq t \leq 1}$ converges *uniformly in probability (in u.p.)* to some process $Y = (Y_t)_{0 \leq t \leq 1}$ if the supremum norm of the difference converges to 0 in probability. The proof of our Itô formula (1.1) will involve convergence in u.p. of the following terms:

$$(2.8) \quad \lim_{n \rightarrow \infty} \sum_{\substack{t_i \in D_n \\ t_i < t}} f(X_{t_i})(X_{t_{i+1}} - X_{t_i}) = \int_0^t f(X_s) dX_s,$$

$$(2.9) \quad \lim_{n \rightarrow \infty} Q_t^n = [f(X), X]_t$$

$$(2.10) \quad \lim_{n \rightarrow \infty} R_t^n = 0.$$

(2.11) **Remark.** In each case, convergence in u.p. is known to hold under certain regularity conditions on the function f :

- 1) For $f \in \mathcal{C}$ the convergence in (2.8) holds in u.p.; cf. [8, p.57]
- 2) For $f \in \mathcal{C}^1$ one can proceed in a strictly pathwise manner; cf. [3]. In fact, almost all Brownian paths have the property that the discrete measures

$$\sum_{t_i \in D_n} (X_{t_{i+1}} - X_{t_i})^2 \delta_{t_i}$$

converge weakly to the uniform distribution on $[0, 1]$. But for any such path one can verify the following facts, without any further use of probabilistic arguments: Writing

$$(2.12) \quad Q_t^n = \sum_{\substack{t_i \in D_n \\ t_i < t}} f'(\xi_i)(X_{t_{i+1}} - X_{t_i})^2$$

for some ξ_i between $X_{t_{i+1}}$ and X_{t_i} , we get convergence of (2.12) to the covariation

$$(2.13) \quad [f(X), X]_t = \int_0^t f'(X_s) ds.$$

Moreover, the stochastic integral in (2.8) can be defined pathwise as the limit of the sums in (2.8), and the remainder terms in (2.10) converge to 0; see [3] for the proofs.

3) For a function $f \in C^2$ the convergence of the remainder terms R_t^n to 0 can also be seen from the trapezoidal rule

$$(2.14) \quad \int_a^b \left\{ f(y) - \frac{1}{2}(f(a) + f(b)) \right\} dy = -\frac{1}{12}(b-a)^3 f''(\xi) :$$

This implies

$$(2.15) \quad |R_t^n| \leq \frac{1}{12} \max |f''(X_t)| \sum_{\substack{t_i \in D_n \\ t_i < t}} |X_{t_{i+1}} - X_{t_i}|^3$$

hence convergence to 0 for any continuous path with bounded quadratic variation.

The main point of this paper is to show that no such regularity conditions are needed and that each of the three convergence statements (2.8) - (2.10) holds in u.p. for any locally square integrable function f . Since we can write

$$(2.16) \quad Q_t^n = \sum_{\substack{t_i \in D_n \\ t_i < t}} f(X_{t_{i+1}})(X_{t_{i+1}} - X_{t_i}) - \sum_{\substack{t_i \in D_n \\ t_i < t}} f(X_{t_i})(X_{t_{i+1}} - X_{t_i})$$

the existence of the quadratic covariation in (2.9) and its representation (1.5) as the difference of a backward and a forward stochastic integral will both follow from the discrete approximation (2.8) of the forward integral and from a corresponding approximation of the backward integral by the first sum in (2.16). Let us now recall the definition of the stochastic backward integral.

We assume without loss of generality that our Brownian motion $X = (X_t)_{0 \leq t \leq 1}$ is defined as the coordinate process on the canonical path space $\Omega = C[0, 1]$ with Wiener measure P . We will have need of the time reversal operator $(RX)_t = X_{1-t}$ on Ω . Note that the law P^* of RX is the law of the Brownian bridge process with initial distribution $N(0, 1)$ and terminal value 0. Thus, the coordinate process X is a semimartingale under P^* with decomposition

$$(2.17) \quad X_t = X_0 + W_t + \int_0^t \frac{-X_s}{1-s} ds$$

where W is a Brownian motion with respect to P^* . For a measurable function $f(x, t)$ on $R \times [0, 1]$ such that

$$(2.18) \quad \begin{aligned} E^* \left[\int_0^1 \left\{ f^2(X_s, 1-s) + |f(X_s, 1-s)| \frac{|X_s|}{1-s} \right\} ds \right] \\ = E \left[\int_0^1 \left\{ f^2(X_s, s) + |f(X_s, s)| \frac{|X_s|}{s} \right\} ds \right] < \infty \end{aligned}$$

the stochastic integral

$$(2.19) \quad \int_0^t f(X_s, 1-s) dX_s = \int_0^t f(X_s, 1-s) dW_s + \int_0^t f(X_s, 1-s) \frac{-X_s}{1-s} ds$$

is well defined for any $t \in [0, 1]$ with respect to P^* . Thus, the following *backward integral*

$$(2.20) \quad \int_0^t f(X_s, s) d^* X_s \equiv - \left(\int_{1-t}^1 f(X_s, 1-s) dX_s \right) \circ R.$$

is well defined in terms of Wiener measure P .

3. Existence of quadratic covariation

Our purpose is to show that the quadratic variation

$$(3.1) \quad [f(X), X]_t = \lim_{n \rightarrow \infty} \sum_{t_i \in D_n, t_i \leq t} (f(X_{t_{i+1}}) - f(X_{t_i}))(X_{t_{i+1}} - X_{t_i})$$

exists as a limit in u.p., beyond the usual cases where $f(X)$ is a semimartingale. In order to illustrate the basic idea we begin with the following preliminary result:

(3.2) **Proposition.** *Let f be a continuous function. Then the quadratic covariation $[f(X), X]$ exists as a continuous process, and it satisfies*

$$(3.3) \quad [f(X), X]_t = \int_0^t f(X_s) d^* X_s - \int_0^t f(X_s) dX_s.$$

Proof. Since X is a semimartingale and $f(X)$ is an adapted continuous process, we know that

$$(3.4) \quad \lim_{n \rightarrow \infty} \sum_{t_i \in D_n, t_i \leq t} f(X_{t_i})(X_{t_{i+1}} - X_{t_i}) = \int_0^t f(X_s) dX_s$$

with convergence in u.p.; cf. [7, p. 57]). On the other hand we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \sum_{\substack{t_i \in D_n \\ t_i \leq t}} f(X_{t_{i+1}})(X_{t_{i+1}} - X_{t_i}) &= \lim_{n \rightarrow \infty} \left\{ \sum_{\substack{t_i \in D_n \\ t_i \leq t}} f(X_{1-t_{i+1}})(X_{1-t_{i+1}} - X_{1-t_i}) \right\} \circ R \\ &= \lim_{n \rightarrow \infty} - \left\{ \sum_{\substack{s_i \in D_n \\ s_{i+1} \geq 1-t}} f(X_{s_i})(X_{s_{i+1}} - X_{s_i}) \right\} \circ R. \end{aligned}$$

Since the time reversed process $X \circ R$ is also a semimartingale (as shown by (2.17)), we have convergence in u.p. of the above sums (again by [8, p. 57]) to

$$- \left\{ \int_{1-t}^1 f(X_s) dX_s \right\} \circ R = \int_0^t f(X_s) d^* X_s,$$

hence

$$(3.5) \quad \lim_{n \rightarrow \infty} \sum_{t_i \in D_n, t_i \leq t} f(X_{t_{i+1}})(X_{t_{i+1}} - X_{t_i}) = \int_0^t f(X_s) d^* X_s.$$

If we now subtract (3.4) from (3.5) we obtain the existence of $[f(X), X]$ as a limit of sums in u.p., as well as its identification as the difference of two stochastic integrals. Since both stochastic integral processes are continuous, so also is $[f(X), X]$.

(3.6) **Remark.** The process $[f(X), X]$, while continuous a.s., need not be of bounded variation as it is in the usual theory when f is taken to be \mathcal{C}^1 . Indeed if f is continuous but not of bounded variation, then its primitive F is not the difference of two convex functions, and therefore $F(X)$ is *not* a semimartingale. Theorem (4.3) will show that $F(X_t) - \frac{1}{2}[f(X), X]_t$ is a local martingale. Thus, $[f(X), X]$ cannot have path of bounded variation if f is not a function of bounded variation. However, theorem (3.45) will show that the quadratic covariation is always a process of "zero energy".

Let us now show that the quadratic covariation $[f(X), X]$ exists for any f which is locally square integrable. The idea is quite simple: As in the proof of (3.2), it is enough to show that the forward and the backward stochastic integral can both be approximated by the simple sums in (3.4) and (3.5), without any additional smoothing. The existence of quadratic covariation then follows by subtracting these sums.

(3.7) **Proposition.** 1) Let f be a square-integrable function on R^1 . Then

$$(3.8) \quad \lim_{n \rightarrow \infty} \sum_{0 < t_i \in D_n, t_i \leq t} f(X_{t_i})(X_{t_{i+1}} - X_{t_i}) = \int_0^t f(X_s) dX_s$$

in $L^2(P)$ and

$$(3.9) \quad \lim_{n \rightarrow \infty} \sum_{t_i \in D_n, t_i \leq t} f(X_{t_{i+1}})(X_{t_{i+1}} - X_{t_i}) = \int_0^t f(X_s) d^* X_s$$

in $L^1(P)$.

2) If f is locally square integrable then convergence in (3.8) and (3.9) holds in u.p., and so the quadratic covariation exists in u.p. and satisfies

$$(3.10) \quad [f(X), X]_t = \int_0^t f(X_s) d^* X_s - \int_0^t f(X_s) dX_s.$$

Instead of proving proposition (3.7), we pass directly to the *time-dependent* case and prove the corresponding general version. Let $f(x, t)$ be a measurable function on $R^1 \times [0, 1]$. The following proof shows that the analogue of part 1) holds if $f(\cdot, t)$ is square integrable and weakly continuous in t as a map from $[0, 1]$ to $L^2(R^1)$. The analogue of part 2) involves weak continuity of $f(\cdot, t)$ as a map from $[0, 1]$ to $L^2_{loc}(R^1)$ in the sense that the integral $\int f(x, t)g(x)dx$ is continuous in t for any square integrable function g with compact support.

(3.11) **Theorem.** *Suppose that $f(\cdot, t)$ is locally square integrable and that*

$$(3.12) \quad f(\cdot, t) \text{ is weakly continuous in } t$$

as a map from $[0, 1]$ to $L^2_{loc}(R^1)$. Then the quadratic covariation

$$(3.13) \quad [f(X, \cdot), X]_t = \lim_{n \rightarrow \infty} \sum_{t_i \in D_n, 0 < t_i \leq t} \{f(X_{t_{i+1}}, t_{i+1}) - f(X_{t_i}, t_i)\}(X_{t_{i+1}} - X_{t_i})$$

exists as a limit in u.p., and

$$(3.14) \quad [f(X, \cdot), X]_t = \int_0^t f(X_s, s) d^* X_s - \int_0^t f(X_s, s) dX_s.$$

Proof. By the usual localization argument, it is enough to consider the case where $f(\cdot, t)$ is square integrable and weakly continuous in t as a map from $[0, 1]$ to $L^2(R^1)$. It is also enough to consider the case $t = 1$.

1) Consider the processes ϕ and ϕ_n defined by

$$(3.15) \quad \phi(\omega, t) = f(X_t(\omega), t)$$

and

$$(3.16) \quad \phi_n(\omega, t) = \sum_{0 < t_i \in D_n} f(X_{t_i}(\omega), t_i) I_{(t_i, t_{i+1}]}(t).$$

In order to prove

$$(3.17) \quad \lim_{n \rightarrow \infty} \sum_{0 < t_i \in D_n} f(X_{t_i}, t_i) (X_{t_{i+1}} - X_{t_i}) = \int_0^1 f(X_s, s) dX_s,$$

in $L^2(P)$, we have to show that

$$\|\phi - \phi_n\|_2 \equiv E \left[\int_0^1 (\phi(\omega, t) - \phi_n(\omega, t))^2 dt \right]^{\frac{1}{2}}$$

converges to 0. Let us use the notation

$$p_t(x) \equiv \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{x^2}{2t}\right).$$

Then

$$(3.18) \quad \begin{aligned} \|\phi\|_2^2 &= E \left[\int_0^1 f^2(X_t, t) dt \right] = \int_0^1 \int f^2(x, t) p_t(x) dx dt \\ &\leq \frac{1}{\sqrt{2\pi}} \int_0^1 \int f^2(x, t) dx \frac{1}{\sqrt{t}} dt \equiv \|f\|_2^2 \end{aligned}$$

and

$$\begin{aligned} \|\phi_n\|_2^2 &= \sum_{0 < t_i \in D_n} \int f^2(x, t_i) p_{t_i}(x) dx (t_{i+1} - t_i) \\ &\leq \frac{1}{\sqrt{2\pi}} \sum_{0 < t_i \in D_n} \int f^2(x, t_i) dx \frac{1}{\sqrt{t_i}} (t_{i+1} - t_i), \end{aligned}$$

hence

$$(3.19) \quad \overline{\lim}_n \|\phi_n\|_2^2 \leq \|f\|_2^2$$

due to the continuity of $\int f(x, t)^2 dx$ in t . Now take a continuous function g on $R \times [0, 1]$ with compact support such that $\|g - f\|_2 \leq \epsilon$ and define the corresponding processes ψ and ψ_n as in (3.15) and (3.16). Then

$$\|\phi - \phi_n\|_2 \leq \|\phi - \psi\|_2 + \|\psi - \psi_n\|_2 + \|\psi_n - \phi_n\|_2.$$

Since $\lim \|\psi - \psi_n\|_2 = 0$ by continuity of g and by Lebesgue's theorem, the two preceding estimates (3.18) and (3.19) applied to $f - g$ imply

$$\overline{\lim}_n \|\phi - \phi_n\|_2 \leq 2\|f - g\|_2 \leq 2\epsilon,$$

hence

$$(3.20) \quad \lim_n \|\phi - \phi_n\|_2 = 0.$$

2) The convergence

$$(3.21) \quad \lim_{n \rightarrow \infty} \sum_{t_i \in D_n, t_i \leq t} f(X_{t_{i+1}}, t_{i+1})(X_{t_{i+1}} - X_{t_i}) = \int_0^t f(X_s, s) d^* X_s$$

in $L^1(P)$ for $t = 1$ is equivalent to the convergence

$$(3.22) \quad \lim_n \sum_{0 < t_i \in D_n} f(X_{t_i}, 1 - t_i)(X_{t_{i+1}} - X_{t_i}) = \int_0^1 f(X_t, 1 - t) dX_t$$

in $L^1(P^*)$, where P^* is the distribution of the time reversed process $X \circ R$. Recall the decomposition (2.17) of X under P^* , and let us first check that

$$(3.23) \quad \lim_n \sum_{0 < t_i \in D_n} f(X_{t_i}, 1 - t_i)(W_{t_{i+1}} - W_{t_i}) = \int_0^1 f(X_t, 1 - t) dW_t$$

in $L^2(P^*)$. But (3.23) is equivalent to the convergence of

$$\begin{aligned} & E^* \left[\int_0^1 [f(X_t, 1 - t) - \sum_{0 < t_i \in D_n} f(X_{t_i}, 1 - t_i) I_{(t_i, t_{i+1})}(t)]^2 dt \right] \\ &= E \left[\int_0^1 [f(X_s, s) - \sum_{1 > s_{i+1} \in D_n} f(X_{s_{i+1}}, s_{i+1}) I_{[s_i, s_{i+1})}(s)]^2 ds \right]. \end{aligned}$$

to 0, and this follows as in part 1). In order to prove (3.22) it remains to show, due to (2.17), that

$$\lim_n \sum_{0 < t_i \in D_n} f(X_{t_i}, 1 - t_i) \int_{t_i}^{t_{i+1}} \frac{X_s}{1-s} ds = \int_0^1 f(X_t, 1-t) \frac{X_t}{1-t} dt$$

in $L^1(P^*)$ or, equivalently,

$$(3.24) \quad \lim_n \sum_{1 > s_{i+1} \in D_n} f(X_{s_{i+1}}, s_{i+1}) \int_{s_i}^{s_{i+1}} \frac{X_s}{s} ds = \int_0^1 f(X_s, s) \frac{X_s}{s} ds$$

in $L^1(P)$. For a measurable function $\psi(\omega, t)$ on $\Omega \times [0, 1]$ we introduce the norm

$$(3.25) \quad \|\psi\|_1 \equiv E \left[\int_0^1 |\psi(\omega, s)| \frac{|X_s|}{s} ds \right].$$

For the function ϕ defined in (3.15) we get

$$\begin{aligned} \|\phi\|_1 &\equiv E \left[\int_0^1 |\psi(\omega, s)| \frac{|X_s|}{s} ds \right] \\ &\leq \int_0^1 \frac{1}{s} E[f^2(X_s, s)]^{\frac{1}{2}} E[X_s^2]^{\frac{1}{2}} ds \\ &= \int_0^1 \frac{1}{s} \left(\int f^2(x, s) p_s(x) dx \right)^{\frac{1}{2}} s^{\frac{1}{2}} ds \\ &\leq \left(\frac{1}{2\pi} \right)^{\frac{1}{4}} \int_0^1 \left(\int f^2(x, s) dx \right)^{\frac{1}{2}} s^{-\frac{3}{4}} ds \equiv \|f\|_1. \end{aligned}$$

Similarly, we see that

$$\phi_n^*(\omega, s) \equiv \sum_{1 > s_{i+1} \in D_n} f(X_{s_{i+1}}, s_{i+1}) I_{[s_i, s_{i+1})}(s)$$

satisfies

$$\begin{aligned} \|\phi_n^*\|_1 &= \sum_{1 > s_{i+1} \in D_n} E \left[|f(X_{s_{i+1}}, s_{i+1})| \int_{s_i}^{s_{i+1}} \frac{|X_s|}{s} ds \right] \\ &\leq \left(\frac{1}{2\pi} \right)^{\frac{1}{4}} \sum_{1 > s_i \in D_n} \left(\int f^2(x, s_{i+1}) dx \right)^{\frac{1}{2}} s_{i+1}^{-\frac{1}{4}} \int_{s_i}^{s_{i+1}} s^{-\frac{1}{2}} ds \\ &\leq \left(\frac{1}{2\pi} \right)^{\frac{1}{4}} \sum_{1 > s_i \in D_n} \left(\int f^2(x, s_{i+1}) dx \right)^{\frac{1}{2}} s_{i+1}^{-\frac{1}{4}} s_i^{-\frac{1}{2}} (s_{i+1} - s_i), \end{aligned}$$

hence

$$(3.26) \quad \overline{\lim} \|\phi_n^*\|_1 \leq \left(\frac{1}{2\pi}\right)^{\frac{1}{2}} \int_0^1 \left(\int f^2(x, s) dx\right)^{\frac{1}{2}} s^{-\frac{3}{2}} ds = \|f\|_1.$$

Now we choose a function $g \in C(R^1 \times [0, 1])$ with compact support such that $\|g - f\|_1 \leq \epsilon$ and conclude as in part 1) that

$$(3.27) \quad \lim_n \|\phi - \phi_n^*\|_1 = 0.$$

This implies (3.24), and so we have shown (3.21).

3) Subtracting (3.17) from (3.21) we obtain (3.13).

Even though $f(X, \cdot)$ is in general not a semimartingale, the existence of quadratic covariation $[f(X, \cdot), X]$ implies the *existence* of the stochastic integral

$$(3.28) \quad \int_0^t X_s df(X_s, s) = \lim_{n \rightarrow \infty} \sum_{t_i \in D_n, 0 < t_i \leq t} X_{t_i} \{f(X_{t_{i+1}}, t_{i+1}) - f(X_{t_i}, t_i)\}$$

and the following *integration by parts formula*:

(3.29) **Corollary.** *Under the assumptions of theorem (3.11) we have*

$$(3.30) \quad f(X_t, t)X_t = f(X_0, 0)X_0 + \int_0^t X_s df(X_s, s) + \int_0^t f(X_s, s) dX_s + [f(X, \cdot), X]_t.$$

Proof. This follows from the relation

$$\begin{aligned} f(X_{t_{i+1}}, t_{i+1})X_{t_{i+1}} - f(X_{t_i}, t_i)X_{t_i} &= X_{t_i} (f(X_{t_{i+1}}, t_{i+1}) - f(X_{t_i}, t_i)) \\ &\quad + f(X_{t_i}, t_i)(X_{t_{i+1}} - X_{t_i}) \\ &\quad + (f(X_{t_{i+1}}, t_{i+1}) - f(X_{t_i}, t_i))(X_{t_{i+1}} - X_{t_i}) \end{aligned}$$

and the convergence in (3.17) and (3.13).

(3.31) **Remarks.** 1) If we average (3.17) and (3.21) and use (3.14) then we obtain

$$(3.32) \quad \begin{aligned} \lim_{n \rightarrow \infty} \sum_{t_i \in D_n, t_i \leq t} \frac{1}{2} \{f(X_{t_{i+1}}, t_{i+1}) + f(X_{t_i}, t_i)\} (X_{t_{i+1}} - X_{t_i}) \\ = \int_0^t f(X_s, s) dX_s + \frac{1}{2} [f(X, \cdot), X]_t. \end{aligned}$$

in u.p.. This will be used in our definition (4.18) and (5.12) of the Stratonovich integral.
 2) Suppose that f_n converges to f in the norm $\|\cdot\|_2$ of (3.18). The proof of (3.11) shows that the corresponding forward and backward stochastic integrals converge, and so (3.14) implies

$$(3.33) \quad [f(X, \cdot), X]_t = \lim_{n \rightarrow \infty} [f_n(X, \cdot), X]_t.$$

3) If $f(\cdot, t)$ is absolutely continuous with derivative $f_x(\cdot, t)$ then

$$(3.34) \quad [f(X, \cdot), X] = \int_0^t f_x(X_s, s) ds.$$

For a smooth function $f \in C^1(R^1 \times [0, 1])$ this is well known (cf. [8, p. 75]) and can be checked directly from definition (3.13). The extension of (3.34) to the absolutely continuous case follows by approximating f with smooth functions f_n in the norm $\|\cdot\|_2$ and applying the preceding remark; see the proof of (4.3).

(3.35) **Example** Let $a(t)$ denote a continuous function on $[0, 1]$. The function

$$(3.36) \quad f(x, t) = I_{[a(t), \infty)}(x)$$

satisfies the conditions of (3.11), and so the quadratic covariation in (3.13) exists. Let us define the *local time of Brownian motion at the continuous curve $a(\cdot)$* as the continuous process $L^{a(\cdot)}$ given by

$$(3.37) \quad L_t^{a(\cdot)} = [f(X, \cdot), X]_t.$$

In order to motivate this definition, we take a sequence (ε_n) decreasing to 0. By (3.34) the function

$$f_n(x, t) = \frac{1}{2\varepsilon_n} \int_0^x I_{(a(s)-\varepsilon_n, a(s)+\varepsilon_n)}(y) dy$$

has quadratic covariation

$$[f_n(X, \cdot), X]_t = \frac{1}{2\varepsilon_n} \int_0^t I_{(a(s)-\varepsilon_n, a(s)+\varepsilon_n)}(X_s) ds.$$

But f_n converges to f in the norm $\|\cdot\|_2$ used in (3.18), and so (3.33) implies

$$(3.38) \quad L_t^{a(\cdot)} = \lim_{n \rightarrow \infty} \frac{1}{2\varepsilon_n} \int_0^t I_{(a(s)-\varepsilon_n, a(s)+\varepsilon_n)}(X_s) ds$$

in u.p. In particular we can conclude that the continuous process $L^{a(\cdot)}$ has increasing paths, and that the corresponding random measure on $[0, 1]$ is a.s. concentrated on times where the Brownian path intersects the given function $a(\cdot)$:

$$(3.39) \quad \int_0^1 I_{\{X_t \neq a(t)\}} d_t L_t^{a(\cdot)} = 0.$$

If the function $a(\cdot)$ assumes a constant level a then

$$(f(X_{t_{i+1}}, t_{i+1}) - f(X_{t_i}, t_i))(X_{t_{i+1}} - X_{t_i}) = |X_{t_{i+1}} - X_{t_i}| I_{C_{n,i}}$$

where

$$C_{n,i} = \{\text{sign}(X_{t_i} - a) \neq \text{sign}(X_{t_{i+1}} - a)\}$$

with $t_i \in D_n$ denotes the set of paths which exhibit a crossing of the level a if checked at times t_i and t_{i+1} . Thus the existence of the quadratic covariation (3.13) implies the identity

$$(3.40) \quad L_t^a = \lim_{n \rightarrow \infty} \sum_{t_i \in D_n, t_i \leq t} |X_{t_{i+1}} - X_{t_i}| I_{C_{n,i}}$$

for the local time of Brownian motion at a constant level a .

Let us now return to the general situation of theorem (3.11) and let us show that quadratic covariation is a continuous process of zero energy:

(3.41) **Definition.** For a process $Y = (Y_t)_{0 \leq t \leq 1}$ with continuous paths we define the *quadratic variation*

$$(3.42) \quad [Y]_t = \lim_{n \rightarrow \infty} \sum_{t_i \in D_n, t_i \leq t} (Y_{t_{i+1}} - Y_{t_i})^2$$

whenever this limit exists uniformly in probability. If $[Y]_1 = 0$ P -a.s. then Y is called a process of *zero energy*.

(3.43) **Remarks.** 1) Any process with continuous paths of bounded variation has zero energy.

2) A local martingale of the form

$$Y_t = \int_0^t f(X_s, s) dX_s$$

has quadratic variation

$$(3.44) \quad [Y]_t = \int_0^t f^2(X_s, s) ds.$$

(3.45) **Theorem.** *The quadratic covariation $[f(X, \cdot), X]$ is a continuous process of zero energy.*

Proof. 1) The quadratic covariation $Y_t = [f(X, \cdot), X]_t$ is of the form

$$(3.46) \quad Y_t = Y_t^{(1)} + Y_t^{(2)}$$

with

$$Y_t^{(1)} = \int_0^t f(X_s, s) d^* X_s, \quad Y_t^{(2)} = - \int_0^t f(X_s, s) dX_s.$$

The process $Y^{(2)}$ has quadratic variation

$$(3.47) \quad [Y^{(2)}]_t = \int_0^t f^2(X_s, s) ds.$$

Since

$$Y_t^* = \int_{1-t}^1 f(X_s, 1-s) dX_s$$

has quadratic variation

$$[Y^*]_t = \int_{1-t}^1 f^2(X_s, 1-s) ds$$

under P^* , the process $Y^{(1)} = Y^* \circ R$ has quadratic variation

$$(3.48) \quad [Y^{(1)}]_t = \left\{ \int_{1-t}^1 f^2(X_s, 1-s) ds \right\} \circ R = \int_0^t f^2(X_s, s) ds$$

under P . Thus, the decomposition (3.46) implies the a priori estimate

$$(3.49) \quad [Y]_t \leq 2([Y^{(1)}]_t + [Y^{(2)}]_t) \leq 4 \cdot \int_0^t f^2(X_s, s) ds$$

if, for the moment, we define $[Y]_t$ as the lim sup of the sums in (3.42).

2) Take a sequence of functions $f_n \in C^1(\mathbb{R}^1 \times [0, 1])$ such that f_n converges to f in the norm $\|\cdot\|_2$ used in (3.8). The process

$$Y_t^{(n)} = [f_n(X, \cdot), X]_t = \int_0^t (f_n)_x(X_s, s) ds$$

has continuous paths of bounded variation and therefore has zero energy. Evaluating the squares in the definition (3.42) for $[Y - Y^{(n)}]_1$ and using Cauchy-Schwartz, we see that

$$(3.50) \quad [Y - Y^{(n)}]_1 = [Y]_1.$$

Applying our a priori estimate (3.49) to the difference $f - f_n$, we get

$$(3.51) \quad [Y]_1 = [Y - Y^{(n)}]_1 \leq 4 \int_0^t (f - f_n)^2(X_s, s) ds.$$

But the expectation of the last term converges to 0 due to our estimate (3.18). Thus we have $E[[Y]_1] = 0$, hence $[Y]_1 = 0$ $P - a.s.$

4. An Extension of Itô's Formula

In this section we prove the time independent extension (1.1) of Itô's formula. For $f \in \mathcal{C}^1$ the quadratic covariation takes the well-known form

$$(4.1) \quad [f(X), X]_t = \int_0^t f'(X_s) ds$$

which appears in Itô's formula

$$(4.2) \quad F(X_t) = F(X_0) + \int_0^t f(X_s) dX_s + \frac{1}{2} \int_0^t f'(X_s) ds$$

for $F \in \mathcal{C}^2$ and $f = F'$. Thus, the following formula is an extension of Itô's formula to the case where f is locally square integrable:

(4.3) **Theorem.** *Let F be absolutely continuous with locally square integrable derivative f . Then*

$$(4.4) \quad F(X_t) = F(X_0) + \int_0^t f(X_s) dX_s + \frac{1}{2}[f(X), X]_t.$$

Proof. The quadratic covariation process $[f(X), X]$ exists as shown in Proposition (3.7). By the usual localization argument we may assume that f is square integrable. Take a sequence $f_n \in \mathcal{C}^1$ such that f_n converges to f in L^2 . The estimates in the proof of (3.11) show that

$$(4.5) \quad \lim_{n \rightarrow \infty} \int_0^t f_n(X_s) dX_s = \int_0^t f(X_s) dX_s$$

in $L^2(P)$ and

$$(4.6) \quad \lim_{n \rightarrow \infty} \int_0^t f_n(X_s) d^* X_s = \int_0^t f(X_s) d^* X_s.$$

in $L^1(P)$. By (3.10) and the usual Itô formula (4.2) applied to the functions

$$(4.7) \quad F_n(x) \equiv F(0) + \int_0^x f_n(y) dy$$

we obtain

$$\begin{aligned} \frac{1}{2}[f(X), X]_t &= \lim_{n \rightarrow \infty} \frac{1}{2}[f^n(X), X]_t \\ &= \lim_{n \rightarrow \infty} \{F_n(X_t) - F_n(X_0) - \int_0^t f_n(X_s) dX_s\}, \end{aligned}$$

and this is equal to

$$F(X_t) - F(X_0) - \int_0^t f(X_s) dX_s,$$

again by (4.5).

(4.8) **Remark.** For $t \in D_n$ we can write

$$(4.9) \quad F(X_t) - F(X_0) = \sum_{\substack{t_i \in D_n \\ t_i \leq t}} \{f(X_{t_i})(X_{t_{i+1}} - X_{t_i}) + \int_{X_{t_i}}^{X_{t_{i+1}}} (f(y) - f(X_{t_i})) dy\}$$

and also

$$(4.10) \quad F(X_t) - F(X_0) = \sum_{\substack{t_i \in D_n \\ t_i \leq t}} \{f(X_{t_{i+1}})(X_{t_{i+1}} - X_{t_i}) + \int_{X_{t_i}}^{X_{t_{i+1}}} (f(y) - f(X_{t_{i+1}}))dy\}.$$

Equation (4.9) together with (4.4) and (3.8) implies

$$(4.11) \quad \frac{1}{2}[f(X), X]_t = \lim_{n \rightarrow \infty} \sum_{\substack{t_i \in D_n \\ t_i \leq t}} \int_{X_{t_i}}^{X_{t_{i+1}}} (f(y) - f(X_{t_i}))dy.$$

On the other hand, equation (4.10) together with (4.4), (3.9) and (3.10) implies

$$(4.12) \quad \frac{1}{2}[f(X), X]_t = \lim_{n \rightarrow \infty} \sum_{\substack{t_i \in D_n \\ t_i \leq t}} \int_{X_{t_i}}^{X_{t_{i+1}}} (f(X_{t_{i+1}}) - f(y))dy.$$

If we subtract (4.12) from (4.11) we obtain the remarkable relation

$$(4.13) \quad \lim_{n \rightarrow \infty} \sum_{\substack{t_i \in D_n \\ t_i \leq t}} \int_{X_{t_i}}^{X_{t_{i+1}}} \{f(y) - \frac{1}{2}(f(X_{t_i}) + f(X_{t_{i+1}}))\}dy = 0$$

uniformly in probability for any locally square integrable function f .

(4.14) **Example.** Let us return to example (3.35) in the case where the function $a(\cdot)$ assumes a constant level a . The function $f(x) = I_{[a, \infty)}(x)$ is the derivative of the absolutely continuous function $F(x) = (x - a)^+$. Due to (3.40), our Itô formula (4.4) takes the form

$$(4.15) \quad (X_t - a)^+ = (X_0 - a)^+ + \int_0^t I_{[a, \infty)}(X_s) dX_s + \frac{1}{2}L_t^a$$

where

$$(4.16) \quad L_t^a = \lim_{n \rightarrow \infty} \sum_{t_i \in D_n, t_i \leq t} |X_{t_{i+1}} - X_{t_i}| I_{C_{n,i}}.$$

Comparing (4.15) with the well known Tanaka formula (see, e.g., [8] p.169), we see that L_t^a coincides with Lévy's local time at level a . Note that (4.10) and (4.12) imply the alternative descriptions

$$(4.17) \quad \begin{aligned} \frac{1}{2}L_t^a &= \lim_{n \rightarrow \infty} \sum_{t_i \in D_n, t_i \leq t} |X_{t_i} - a| I_{C_{n,i}} \\ &= \lim_{n \rightarrow \infty} \sum_{t_i \in D_n, t_i \leq t} |X_{t_{i+1}} - a| I_{C_{n,i}}. \end{aligned}$$

K. Itô ([5], [6]) proposed defining the Stratonovich integral by the following formula (4.19) when the quadratic covariation $[f(X), X]$ exists (see also [8, p. 216]). Proposition (3.7) shows that this definition makes sense for any locally square integrable function f , and (3.31) shows that the Stratonovich integral can be described as well as a limit of sums:

(4.18) **Definition.** For any locally square integrable function f the *Stratonovich integral* is well defined by

$$(4.19) \quad \int_0^t f(X_s) \circ dX_s \equiv \int_0^t f(X_s) dX_s + \frac{1}{2} [f(X), X]_t,$$

and it can be computed as

$$(4.20) \quad \int_0^t f(X_s) \circ dX_s = \lim_{n \rightarrow \infty} \sum_{\substack{t_i \in \mathcal{D}_n \\ t_i \leq t}} \frac{1}{2} \{f(X_{t_i}) + f(X_{t_{i+1}})\} (X_{t_{i+1}} - X_{t_i}).$$

With this definition, the Itô formula (4.4) can be reformulated as:

(4.21) **Corollary.** For an absolutely continuous function F with locally square integrable derivative $F' = f$,

$$(4.22) \quad F(X_t) = F(X_0) + \int_0^t f(X_s) \circ dX_s.$$

Note that this is an improvement upon [5] and [8, pp. 222–224], where F is required to be \mathcal{C}^2 . Note also that $\int_0^t f(X_s) \circ dX_s$ need not be a semimartingale, but that it is a *Dirichlet process* in the sense of the following remark.

(4.23) **Remarks.** 1) In order to compare (4.3) with the extension of Itô's formula obtained by Bouleau and Yor (see [1] or alternatively [7, p. 179]), let L_t^a denote the local time of our standard Brownian motion X at level a . Let F be absolutely continuous with a locally bounded Borel measurable derivative f . Using a vector valued measure approach Bouleau and Yor show

$$(4.24) \quad F(X_t) = F(X_0) + \int_0^t f(X_s) dX_s - \frac{1}{2} \int_{\mathbb{R}} f(a) d_a L_t^a.$$

Comparing the Bouleau-Yor formula (4.24) with our formula (4.4), we see that if f is Borel and locally bounded, then

$$(4.25) \quad [f(X), X]_t = - \int_{\mathbb{R}} f(a) d_{\alpha} L_t^{\alpha}$$

2) A general result of Fukushima in the context of Dirichlet spaces implies that, for an absolutely continuous function F on R^1 with locally square integrable derivative f , the process $F(X)$ is a "Dirichlet process", i.e., $F(X)$ can be represented as the sum

$$(4.26) \quad F(X_t) = F(X_0) + \int_0^t f(X_s) dX_s + A_t$$

of a local martingale and a continuous process A of "zero energy"; cf. [4, Ch. 5]. Thus, our Itô formula (4.4) together with theorem (3.45) identifies the process of zero energy A in the Fukushima decomposition (4.26) as a quadratic covariation:

$$(4.27) \quad A_t = \frac{1}{2} [f(X), X]_t.$$

Note that in our approach the process A is computed directly on the paths of Brownian motion, without any smoothing of the function f as it is usually done in the theory of Dirichlet spaces.

The following section shows that, in contrast to the Bouleau-Yor formula (4.24) and to the Fukushima formula (4.26), our version (4.4) of the Itô formula for locally square integrable functions f admits a straightforward extension to the time dependent case.

5. The time dependent case

Let $f(x, t)$ be a measurable function on $R^1 \times [0, 1]$ such that $f(\cdot, t)$ is locally square integrable and assume that

$$(5.1) \quad f(\cdot, t) \text{ is weakly continuous in } t$$

as a map from $[0, 1]$ to L_{loc}^2 . Under this assumption, the existence of quadratic covariation

$$(5.2) \quad [f(X, \cdot), X]_t = \lim_{n \rightarrow \infty} \sum_{t_i \in D_n, t_i \leq t} \{f(X_{t_{i+1}}, t_{i+1}) - f(X_{t_i}, t_i)\} (X_{t_{i+1}} - X_{t_i})$$

was already shown in the proof of (3.11). Let us now establish the corresponding time dependent version of Itô's formula:

(5.3) **Theorem.** *Suppose that $F(x, t)$ is absolutely continuous in x and that the partial derivative $f(\cdot, t) \equiv F_x(\cdot, t)$ satisfies the preceding assumptions. Then Itô's formula holds in the form*

$$(5.4) \quad F(X_t, t) = F(X_0, 0) + \int_0^t f(X_s, s) dX_s + \frac{1}{2} [f(X, \cdot), X]_t + \int_0^t F(X_s, ds)$$

where

$$(5.5) \quad \int_0^t F(X_s, ds) \equiv \lim_{n \rightarrow \infty} \sum_{t_i \in D_n, t_i \leq t} F(X_{t_{i+1}}, t_{i+1}) - F(X_{t_{i+1}}, t_i)$$

exists uniformly in probability.

(5.6) **Remark.** If $F(x, \cdot)$ is absolutely continuous in t with derivative $F_t(x, \cdot)$ and if

(5.7) $F_t(\cdot, t)$ is weakly continuous in t

as a map from $[0, 1]$ to L^1_{loc} , then the last term in our Itô formula (5.4) takes the usual form

$$(5.8) \quad \int_0^t F(X_s, ds) = \int_0^t F_t(X_s, s) ds.$$

Proof. 1) Let us write

$$F(X_t, t) - F(X_0, 0) = A_t^n + B_t^n$$

where

$$A_t^n = \sum_{t_i \in D_n, t_i \leq t} F(X_{t_{i+1}}, t_{i+1}) - F(X_{t_{i+1}}, t_i)$$

and

$$B_t^n = \sum_{t_i \in D_n, t_i \leq t} F(X_{t_{i+1}}, t_i) - F(X_{t_i}, t_i).$$

By the time-homogeneous Itô formula (4.4),

$$\begin{aligned} F(X_{t_{i+1}}, t_i) - F(X_{t_i}, t_i) &= \int_{t_i}^{t_{i+1}} f(X_t, t_i) dX_t + \frac{1}{2} ([f(X, t_i), X]_{t_{i+1}} - [f(X, t_i), X]_{t_i}) \\ &= \frac{1}{2} \left\{ \int_{t_i}^{t_{i+1}} f(X_t, t_i) d^* X_t + \int_{t_i}^{t_{i+1}} f(X_t, t_i) dX_t \right\} \end{aligned}$$

due to (3.10). But under our continuity assumption (5.1), the arguments in the proof of (3.11) show that the sum B_t^n of these terms converges to

$$\begin{aligned} \lim_{n \rightarrow \infty} B_t^n &= \frac{1}{2} \left\{ \int_0^t f(X_s, s) d^* X_s + \int_0^t f(X_s, s) dX_s \right\} \\ &= \int_0^t f(X_s, s) dX_s + \frac{1}{2} [f(X, \cdot), X]_t \end{aligned}$$

uniformly in probability. Thus, also the sum A_t^n converges uniformly in probability, and

$$\begin{aligned} \int_0^t F(X_s, ds) &\equiv \lim_{n \rightarrow \infty} A_t^n \\ &= F(X_t, 0) - F(X_0, 0) - \int_0^t f(X_s, s) dX_s - \frac{1}{2} [f(X, \cdot), X]_t. \end{aligned}$$

This shows that the limit in (5.5) exists and that (5.4) holds.

2) In view of (5.8) suppose that $F(x, \cdot)$ is absolutely continuous in t with derivative $F_t(x, \cdot)$.

For $t \in D_n$ we have

$$A_t^n = \sum_{t_i \in D_n, t_i \leq t} \int_{t_i}^{t_{i+1}} F_t(X_{t_{i+1}}, s) ds = \int_0^t F_t(X_s^{(n)}, s) ds$$

where $X_s^{(n)} \equiv X_{t_{i+1}}$ for $s \in (t_i, t_{i+1}]$. Under an additional continuity assumption on F_t we see that

$$\lim_{n \rightarrow \infty} A_t^n = \int_0^t F_t(X_s, s) ds$$

uniformly in probability, and this implies (5.8). For example, the arguments in the proof of (3.11) show that it is enough to assume condition (5.7).

(5.9) **Remarks.** 1) Our Itô formula (5.4) shows that the process

$$(5.10) \quad F(X_t, t) - \int_0^t F(X_s, ds) \quad (0 \leq t \leq 1)$$

is a *Dirichlet process* in the sense of (4.26). Under the additional assumptions in (5.6) the second term has continuous paths of bounded variation. In this case the process $F(X_t, t)$ is itself a Dirichlet process.

2) In analogy to the previous section we see that, under the assumptions on f in (5.3), the *Stratonovich integral* is well defined by

$$(5.11) \quad \int_0^t f(X_s, s) \circ dX_s \equiv \int_0^t f(X_s, s) dX_s + \frac{1}{2} [f(X, \cdot), X]_t,$$

and that it can be computed as

$$(5.12) \quad \int_0^t f(X_s, s) \circ dX_s = \lim_{n \rightarrow \infty} \sum_{\substack{t_i \in D_n \\ t_i \leq t}} \frac{1}{2} \{f(X_{t_i}, t_i) + f(X_{t_{i+1}}, t_{i+1})\} (X_{t_{i+1}} - X_{t_i}).$$

Thus, the Itô formula (5.4) takes the form

$$(5.13) \quad F(X_t, t) = F(X_0, 0) + \int_0^t f(X_s, s) \circ dX_s + \int_0^t F(X_s, ds).$$

3) The time dependent version of the argument for (4.12) shows that

$$(5.14) \quad \frac{1}{2} [f(X, \cdot), X]_t = \lim_{n \rightarrow \infty} \sum_{\substack{t_i \in D_n \\ t_i \leq t}} \int_{X_{t_i}}^{X_{t_{i+1}}} (f(X_{t_{i+1}}, t_i) - f(y, t_i)) dy.$$

(5.15) **Example.** Let us return to the situation in (3.35) where $f(x, t) = I_{[a(t), \infty)}(x)$ is the partial derivative F_x of the function $F(x, t) = (x - a(t))^+$. Itô's formula takes the form

$$(5.16) \quad (X_t - a(t))^+ = (X_0 - a(0))^+ + \int_0^t I_{[a(s), \infty)}(X_s) dX_s + \frac{1}{2} L_t^{a(\cdot)} + \int_0^t F(X_s, ds).$$

In the special case where $a(\cdot)$ is a function of bounded variation we get

$$(5.17) \quad \int_0^t F(X_s, ds) = - \int_0^t I_{[a(s), \infty)}(X_s) da(t),$$

and so (5.16) reduces to

$$(5.18) \quad (X_t - a(t))^+ = (X_0 - a(0))^+ + \int_0^t I_{[a(s), \infty)}(X_s) d(X - a(\cdot)) + \frac{1}{2} L_t^{a(\cdot)}.$$

In this special case, the process $X - a(\cdot)$ is a semimartingale, $L^{a(\cdot)}$ may be viewed as the local time of this semimartingale at level 0, and (5.18) is the corresponding Tanaka formula. Thus (5.16) is an extension of the Tanaka formula to the general case of a continuous function $a(\cdot)$.

As a special case of (5.14) we obtain the following identity for the local time at a continuous curve, in analogy to (4.17):

$$(5.19) \quad \frac{1}{2} L_t^{a(\cdot)} = \lim_{n \rightarrow \infty} \sum_{t_i \in D_n, t_i \leq t} |X_{t_{i+1}} - a(t_i)| I_{C_{n,i}}$$

where

$$C_{n,i} = \{\text{sign}(X_{t_i} - a(t_i)) \neq \text{sign}(X_{t_{i+1}} - a(t_i))\}$$

denotes the set of paths which exhibit a crossing of the level $a(t_i)$ if checked at times t_i and t_{i+1} .

References

- [1] Bouleau, N. and Yor (1981) Sur la variation quadratique des temps locaux de certaines semimartingales. In: *C.R. Acad. Sc. Paris*, vol 292, 491-494.
- [2] Dellacherie, C. and Meyer, P.A. (1980) Probabilités et Potentiel, Ch. V - VIII, Théorie des Martingales. *Hermann, Paris*.
- [3] Föllmer, H. (1981) Calcul d'Itô sans Probabilités. In: *Séminaire Probabilités XV. Lecture Notes in Math.* vol 850, 144-150. Springer, Berlin, New York.

- [4] Fukushima, M. (1980) Dirichlet Forms and Markov Processes. *North-Holland*.
- [5] Itô, K. (1974) Stochastic Differentials. *Applied Math. and Optimization*, vol 1, 374–381.
- [6] Itô, K. (1975) Stochastic Calculus. *Springer Lect. Notes in Physics*, vol. 39, 218–223.
- [7] Lyons, T.J. and Zhang, T.S. (1994) Decomposition of Dirichlet processes and its application. *Annals of Probability*, vo. 22, No.1, 494 – 524.
- [8] Protter, P. (1990) Stochastic Integration and Differential Equations: A New Approach. *Springer, Berlin, New York*.
- [9] Russo, F. and Vallois, P. (1994) Itô formula for C^1 -functions of semimartingales. *Preprint*.