

OPTIMAL RATE OF EMPIRICAL BAYES TESTS FOR  
LOWER TRUNCATION PARAMETERS

by

Shanti S. Gupta      and      Jianjun Li  
Purdue University      Purdue University

Technical Report # 00-07

Department of Statistics  
Purdue University  
West Lafayette, IN USA

October 2000

# OPTIMAL RATE OF EMPIRICAL BAYES TESTS FOR LOWER TRUNCATION PARAMETERS <sup>1</sup>

Shanti S. Gupta

Jianjun Li

Department of Statistics    Department of Statistics

Purdue University

Purdue University

W. Lafayette, IN 47907    W. Lafayette, IN 47907

**Abstract:** The distributions with lower truncation parameters are important models in statistics and have been studied in recent years. In this paper, we consider the one-sided testing problem for lower truncation parameters through the empirical Bayes approach. The optimal rate of the monotone empirical Bayes tests is obtained and a monotone empirical Bayes test  $\delta_n$  achieving the optimal rate is constructed. It is shown that  $\delta_n$  has good performance for both small samples and large samples.

MS Classification: 62C12.

Keywords: Empirical Bayes, regret Bayes risk, optimal rate of convergence, minimax.

---

<sup>1</sup>This research was supported in part by US Army Research Office, Grant DAAD19-00-1-0502 at Purdue University.

**1. Introduction.** Let  $X$  denote a random variable having density function

$$f(x|\theta) = a(x)/A(\theta), \quad \theta \leq x < b \leq \infty, \quad (1.1)$$

where  $a(x)$  is a positive, continuous function on  $(0, b)$ ,  $A(\theta) = \int_{\theta}^b a(x)dx < \infty$  for every  $\theta > 0$ ,  $\theta$  is the parameter, which is distributed according to an unknown prior distribution  $G$  on  $(0, b)$ . Two typical examples of (1.1) are: (I) the exponential distribution with a location parameter:  $f(x|\theta) = e^{-(x-\theta)}$ ,  $x \geq \theta$ , and (II) the Pareto distribution:  $f(x|\theta) = \alpha\theta^\alpha/x^{\alpha+1}$ ,  $x \geq \theta$ .

We consider the problem of testing the hypotheses  $H_0 : \theta \leq \theta_0$  versus  $H_1 : \theta > \theta_0$ , where  $\theta_0$  is known and  $0 < \theta_0 < b$ . The loss function is  $l(\theta, 0) = \max\{\theta - \theta_0, 0\}$  for accepting  $H_0$  and  $l(\theta, 1) = \max\{\theta_0 - \theta, 0\}$  for accepting  $H_1$ . A test  $\delta(x)$  is defined to be a measurable mapping from  $(0, \infty)$  into  $[0, 1]$  so that  $\delta(x) = P\{\text{accepting } H_1 | X = x\}$ , i.e.,  $\delta(x)$  is the probability of accepting  $H_1$  when  $X = x$  is observed. Let  $R(G, \delta)$  denote the Bayes risk of the test  $\delta$  when  $G$  is the prior distribution. Given that  $\int_0^\infty \theta dG(\theta) < \infty$ , a Bayes test  $\delta_G$  is found as

$$\delta_G(x) = 1 \quad \text{if} \quad E[\theta|X = x] \geq \theta_0; \quad \delta_G(x) = 0 \quad \text{if} \quad E[\theta|X = x] < \theta_0. \quad (1.2)$$

Because  $E[\theta|X = x]$  involves  $G$ , the above solution works only if the prior  $G$  is given. If  $G$  is unknown, this testing problem is formed as a compound decision problem and the empirical Bayes approach is used. Let  $X_1, X_2, \dots, X_n$  be the observations from  $n$  independent past experiences. Based on  $\widetilde{X}_n = (X_1, X_2, \dots, X_n)$  and  $X$ , an empirical Bayes rule  $\delta_n(X, \widetilde{X}_n)$  can be constructed. The performance of  $\delta_n$  is measured by  $R(G, \delta_n) - R(G, \delta_G)$ , where  $R(G, \delta_n) = E[R(G, \delta_n | \widetilde{X}_n)]$ . The quantity  $R(G, \delta_n) - R(G, \delta_G)$  is referred as the regret Bayes risk (or regret) in the literature.

This empirical Bayes approach was introduced by Robbins (1956, 1964). Since then, it has been widely used in statistics. For the family (1.1), some problems of statistical

inference based on the empirical Bayes method have been considered by Prasad and Singh (1990), Liang (1993), Datta (1994), Huang (1995), Balakrishnan and Ma (1997), Huang and Liang (1997), Ma and Balakrishnan (2000), among others. In this paper, we consider the testing problem and study the empirical Bayes tests for the family (1.1). The optimal rate of convergence of monotone empirical Bayes tests is obtained and a test with the optimal rate is constructed.

The paper is organized as follows. In Section 2 we provide a few preliminary results. In Section 3 we construct a monotone empirical Bayes test  $\delta_n$  and obtain an upper bound of its regret. In Section 4, a minimax lower bound of the regrets of monotone empirical Bayes tests is obtained. Since the rates in the upper bound of Section 3 and lower bound of Section 4 coincide, the optimal rate is identified. As a byproduct, we see that  $\delta_n$  achieves the optimal rate of convergence. The proofs of main results in Section 3 and 4 are given in Section 5.

**2. Preliminary.** We assume that  $P(\theta > \theta_0) \cdot P(\theta < \theta_0) > 0$  in this paper. If  $P(\theta > \theta_0) = 0$  or  $P(\theta < \theta_0) = 0$ , we know which action we should take regardless of the value of  $x$ . For example, if  $P(\theta < \theta_0) = 0$ , we accept  $H_1$  always. So both two cases are excluded in the testing problem. We also assume  $\mu_G \equiv \int_0^\infty \theta dG(\theta) < \infty$  so that the Bayes analysis can be carried out.

Let  $f_G(x) = \int f(x|\theta)dG(\theta)$  be the marginal pdf of  $X$ , and  $\phi_G(x) = E[\theta|X = x]$  be the posterior mean of  $\theta$  given  $X = x$ . Note that  $\phi_G(x)$  is increasing and  $\phi_G(\theta_0) < \theta_0$ . Then the Bayes rule stated in Section 1 can be represented as

$$\delta_G(x) = 1 \text{ if } \phi_G(x) \geq \theta_0 \iff x \geq c_G; \quad \delta_G(x) = 0 \text{ if } \phi_G(x) < \theta_0 \iff x < c_G.$$

where  $c_G = \inf\{x \in (\theta_0, b) : \phi_G(x) \geq \theta_0\}$ .  $c_G$  is called the critical point corresponding to  $G$ .

Since the Bayes rule  $\delta_G$  is characterized by a single number  $c_G$ , a monotone empirical

Bayes test (MEBT) can be constructed through estimating  $c_G$  by  $c_n(X_1, X_2, \dots, X_n)$ , say, and defining

$$\delta_n = \begin{cases} 1 & \text{if } x \geq c_n, \\ 0 & \text{if } x < c_n. \end{cases} \quad (2.1)$$

Then the regret of  $\delta_n$  is

$$R(G, \delta_n) - R(G, \delta_G) = E \int_{c_n}^{c_G} w(x) a(x) dx, \quad (2.2)$$

where  $w(x) = \alpha_G(x)[\theta_0 - \phi_G(x)]$  and  $\alpha_G(x) = \int_{(0,x]} dG(\theta)/A(\theta)$ .

To consider the rate of convergence of  $R(G, \delta_n) - R(G, \delta_G)$ , we assume that for some  $r \geq 1$ ,  $\alpha_G(x)$  is  $r$ -times continuously differentiable and for  $i = 0, 1, \dots, r$ ,

$$\sup_{\theta_0/2 < x < b} |\alpha_G^{(i)}(x)| \leq B_r < \infty. \quad (2.3)$$

Furthermore, we assume that

$$g(c_G) = G'(c_G) \neq 0. \quad (2.4)$$

From (2.3), we know that  $\phi_G(x)$  is continuous. Then  $c_G > \theta_0$  since  $\phi_G(\theta_0) < \theta_0$ . Also, from (2.3) and (2.4),  $\theta_0 < b$  since  $A(b-) = \lim_{x \uparrow b} A(x) = 0$ .

**3. An upper bound.** We shall construct a MEBT and find an upper bound of its regret. The kernel method is used in the construction. Let  $K_0(y)$  be a Borel-measurable, bounded function vanishing outside the interval  $[0, 1]$  such that

$$\int_0^1 y^j K_0(y) dy = \begin{cases} 1 & \text{if } j = 0, \\ 0 & \text{if } j = 1, 2, \dots, r. \end{cases} \quad (3.1)$$

Suppose  $|K_0(y)| \leq B$ . Denote  $K_1(y) = \int_0^y K_0(s) ds$  and  $u_n = n^{-1/(2r+1)}$ . For any  $x \in (0, b)$ ,

define

$$W_n(x) = \frac{\theta_0 - x}{nu_n} \sum_{j=1}^n \frac{K_0\left(\frac{x-X_j}{u_n}\right)}{a(X_j)} - \frac{1}{n} \sum_{j=1}^n \frac{K_1\left(\frac{x-X_j}{u_n}\right)}{a(X_j)}. \quad (3.2)$$

It is shown later that  $W_n(x)$  is a consistent estimator of  $w(x)$ . Since  $P(\theta < \theta_0) > 0$ ,  $\alpha_G(x) > 0$  for  $x > \theta_0$ . Thus  $c_G = \int_{\theta_0}^b I_{[w(x)>0]} dx + \theta_0$ . Let

$$c_n = \int_{\theta_0}^{d_n} I_{[W_n(x)>0]} dx + \theta_0, \quad (3.3)$$

where

$$d_n = \begin{cases} (\theta_0 + \ln n) \wedge b & \text{if } a(b-) > 0, \\ \inf\{x \geq \theta_0 : a(x) < u_n^{1/3}\} \wedge (\theta_0 + \ln n) \wedge b & \text{if } a(b-) = 0. \end{cases}$$

Then we propose a monotone empirical Bayes test  $\delta_n(x)$  by

$$\delta_n = \begin{cases} 1 & \text{if } x \geq c_n, \\ 0 & \text{if } x < c_n. \end{cases} \quad (3.4)$$

Note that  $d_n \rightarrow b$ . As  $d_n > c_G$ ,

$$c_n - c_G = - \int_{\theta_0}^{c_G} I_{[W_n(x) \leq 0]} dx + \int_{c_G}^{d_n} I_{[W_n(x) > 0]} dx. \quad (3.5)$$

Note that  $\delta_n$  is a monotone rule. It has good performance for small samples (See Van Houwelingen (1976)). Next we show that  $\delta_n$  is a good procedure not only for small samples but also for large samples.

From (2.3),  $w'(x)$  is continuous and  $w'(x) = g(x)(\theta_0 - x)/A(x)$ . Since  $g(c_G) \neq 0$  and  $c_G > \theta_0$ ,  $w'(c_G) < 0$ . Then  $w'(x) < 0$  in a neighbourhood of  $c_G$ . For  $\epsilon > 0$ , define  $A_\epsilon \equiv \min\{-w'(x) : x \in [c_G - \epsilon, c_G + \epsilon]\}$ . Suppose  $\epsilon_G > 0$  such that  $\theta_0 < c_G - \epsilon_G < c_G + \epsilon_G < b$  and  $A_{\epsilon_G} > 0$ . Then for  $0 < \epsilon < \epsilon_G$ ,  $A_\epsilon \geq A_{\epsilon_G} > 0$ .

**Lemma 3.1.** *Let  $\bar{a}$  and  $\bar{w}$  be the supremum values of  $a(x)$  and  $-w'(x)$  on  $[c_G - \epsilon_G, c_G + \epsilon_G]$  respectively. Then*

$$R(G, \delta_n) - R(G, \delta_G) \leq (\theta_0 + \mu_G) \epsilon_G^{-4} E(c_n - c_G)^4 + 1/2 \bar{a} \bar{w} E(c_n - c_G)^2. \quad (3.6)$$

**Lemma 3.2.** *Let  $M = B^2 c_G^2 \{3 + 16B_r [a(c_G)]^2\}$ . Then*

$$(3.7.1) \quad \lim_{n \rightarrow \infty} n^{\frac{2r}{2r+1}} E(c_n - c_G)^2 \leq M/[a(c_G)w'(c_G)]^2; \quad (3.7.2) \quad \lim_{n \rightarrow \infty} n^{\frac{2r}{2r+1}} E(c_n - c_G)^4 = 0.$$

The proofs of Lemma 3.1 and Lemma 3.2 are given in Section 5. Note that, as  $\epsilon_G \rightarrow 0$ ,  $\bar{a}\bar{w} \rightarrow a(c_G)|w'(c_G)|$ . Therefore the previous two lemmas give the following theorem.

**Theorem 3.1.** *Let  $M$  be the number defined in Lemma 3.2. Then*

$$\lim_{n \rightarrow \infty} n^{\frac{2r}{2r+1}} [R(G, \delta_n) - R(G, \delta_G)] \leq M/[2a(c_G)|w'(c_G)|]. \quad (3.8)$$

To consider the uniform convergence rate of  $\delta_n$ , we define a class of prior distributions.

Denote

$$\mathcal{G} = \{G : G \text{ satisfies } \mu_G \leq \mu_0, (2.3), c_0 \leq c_G \leq \rho_0, \min_{x \in [\bar{c}_0, \bar{\rho}_0]} |w'(x)| \geq L\}, \quad (3.9)$$

where  $\mu_0 < \infty$ ,  $\theta_0 < c_0 < \rho_0 < b$ ,  $\bar{c}_0 = (c_0 + \theta_0)/2$ ,  $\bar{\rho}_0 = (2\rho_0) \wedge ((\rho_0 + b)/2)$  and  $L > 0$ .

Assume that  $\mathcal{G}$  is not empty in the following.

**Theorem 3.2.** *For some  $l > 0$ ,*

$$\sup_{G \in \mathcal{G}} [R(G, \delta_n) - R(G, \delta_G)] \leq l \cdot n^{-\frac{2r}{2r+1}} \quad (3.10)$$

**4. A lower bound.** We shall obtain a minimax lower bound for the regrets of all monotone empirical Bayes tests first. In the following parts of this paper,  $l_1, l_2, \dots$  stand for the positive constants, which may have different values on different occasions.

Let  $\mathcal{C}$  be the set of all estimators  $c_n^*$  with  $c_n^* \geq 0$  and let  $\mathcal{D}$  be the set of all empirical Bayes rules of type (2.1) with  $c_n = c_n^* \in \mathcal{C}$ . Let  $\mathcal{F} = \{f_G(x) = \int f(x|\theta)dG(\theta) : G \in \mathcal{G}\}$  and

$c_f$  be the critical points corresponding to  $f \in \mathcal{F}$ .

**Lemma 4.1.**

$$\begin{aligned} & \inf_{\delta_n^* \in \mathcal{D}} \sup_{G \in \mathcal{G}} [R(G, \delta_n^*) - R(G, \delta_G)] \\ & \geq l_1 \sup \{(c_{f_1} - c_{f_2})^2 : \int [\sqrt{f_1(x)} - \sqrt{f_2(x)}]^2 dx \leq l_2/n, f_1, f_2 \in \mathcal{F}\}. \end{aligned}$$

Suppose that  $G_1 \in \mathcal{G}$  with density  $g_1(\theta)$  and  $c_{f_1} \in (c_0, \rho_0)$ . Let  $g_2(\theta) = (1 + u_n^r \mu_n)^{-1} [g_1(\theta) + u_n^{r-1} A(\theta) H(\frac{\theta - c_{f_1}}{u_n})] I_{[\theta > 0]}$ , where  $\mu_n = \int_{-1}^1 A(c_{f_1} + t u_n) H(t) dt$  and  $H(t)$  is a function such that (1) it has support  $[-1, 1]$ , (2)  $\int_{-1}^1 H(t) dt = 0$  and  $\int_{-1}^0 H(t) dt \neq 0$ , and (3) it has bounded derivatives upto order  $r$ . Let  $f_i(x) = a(x) \int_0^x \frac{g_i(\theta)}{A(\theta)} d\theta$  for  $i = 1, 2$ .

**Lemma 4.2.** *As  $n$  is large,  $f_2 \in \mathcal{F}$ ,*

$$\int [\sqrt{f_1(x)} - \sqrt{f_2(x)}]^2 dx \leq l_2/n \quad \text{and} \quad (c_{f_1} - c_{f_2})^2 \geq l_3 n^{-\frac{2r}{2r+1}}.$$

**Theorem 4.1.** *For some  $l > 0$ ,*

$$\inf_{\delta_n^* \in \mathcal{D}} \sup_{G \in \mathcal{G}} [R(G, \delta_n^*) - R(G, \delta_G)] \geq l \cdot n^{-\frac{2r}{2r+1}}.$$

Theorem 4.1 says that  $n^{-\frac{2r}{2r+1}}$  is the best possible rate of convergence. With the result in (3.10), we conclude that  $n^{-\frac{2r}{2r+1}}$  is the optimal rate of monotone empirical Bayes tests and  $\delta_n$  defined by (3.4) achieves this rate. So  $\delta_n$  has good performance not only for small samples but also for large samples.

## 5. Proofs.

**5.1. Proof of Lemma 3.1.** From (2.2),

$$\begin{aligned} R(G, \delta_n) - R(G, \delta_G) &\leq E[I_{\{|c_n - c_G| > \epsilon_G\}} \int_{c_n}^{c_G} w(x)a(x)dx] + \bar{a}E[I_{\{|c_n - c_G| \leq \epsilon_G\}} \int_{c_n}^{c_G} w(x)a(x)dx] \\ &\leq (\theta_0 + \mu_G)\epsilon_G^{-4}E(c_n - c_G)^4 + 1/2\bar{a}\bar{w}E(c_n - c_G)^2, \end{aligned}$$

where  $\int_{c_n}^{c_G} w(x)a(x)dx \leq (\theta_0 + \mu_G)$  and by Taylor expansion

$$I_{\{|c_n - c_G| \leq \epsilon_G\}} \int_{c_n}^{c_G} w(x)dx = -1/2 \times w'(\hat{c}_n)(c_n - c_G)^2 I_{\{|c_n - c_G| \leq \epsilon_G\}} \leq 1/2\bar{w}(c_n - c_G)^2.$$

**5.2. Proof of Lemma 3.2.** Recall  $A_\epsilon > 0$  for  $\epsilon < \epsilon_G$ . For  $\epsilon < \epsilon_G$ , Let  $\eta_1 = c_G - \epsilon$  and

$\eta_2 = c_G + \epsilon$ . Rewrite  $W_n(x) = \frac{1}{n} \sum_{j=1}^n V_n(X_j, x)$ , where

$$V_n(X_j, x) = \frac{\theta_0 - x}{u_n} \times \frac{K_0(\frac{x-X_j}{u_n})}{a(X_j)} - \frac{K_1(\frac{x-X_j}{u_n})}{a(X_j)}.$$

Note that  $V_n(X_j, x)$  are i.i.d. with for fixed  $x$  and  $n$ . Let  $w_n(x) = E[V_n(X_j, x)]$ ,  $Z_{jn} = V_n(X_j, x) - w_n(x)$ ,  $\sigma_n^2 = EZ_{jn}^2$  and  $\gamma_n = E[|Z_{jn}|^3]$ . Denote  $p_n = \min\{a(x) : x \in [\eta_1 - u_n, \eta_2]\}$ .

Then we have the following lemma. Its proof is given in the subsection 5.5.

**Lemma 5.1.** *The following statements hold for large  $n$ .*

(i) For  $x \in [\theta_0, \eta_1) \cup (\eta_2, b)$ ,  $|w(x)| \geq \epsilon A_\epsilon$  as  $\epsilon < \epsilon_G$ ;

For  $x \in [\theta_0, d_n]$ ,  $|w(x)| \leq (2\theta_0 + \ln n)B_r$ .

(ii) For all  $x \in [\theta_0, d_n]$ ,  $|w_n(x) - w(x)| \leq B_r B u_n^r x \equiv 1/2\beta(x)$ .

(iii) For  $x \in [\eta_1, \eta_2]$ ,  $\sigma_n^2 \leq m(p_n u_n)^{-1}$ ,  $m = (\eta_2 - \theta_0 + u_n)^2 B^2 B_r$ ;

For  $x \in [\theta_0, d_n]$ ,  $\sigma_n^2 \leq l_2(\ln n)^2 u_n^{-4/3}$ .

(iv) For  $x \in [\eta_1, \eta_2]$ ,  $\gamma_n \leq l_3(p_n u_n)^{-2}$ ; For  $x \in [\theta_0, d_n]$ ,  $\gamma_n \leq l_4(\ln n)^3 u_n^{-8/3}$ .

(v) For  $x \in [\theta_0, d_n]$ ,  $w(x) > \beta(x) \implies w_n(x) \geq 1/2w(x)$ .

(vi) For  $x \in [\theta_0, d_n]$ ,  $w(x) < -\beta(x) \implies w_n(x) \leq 1/2w(x)$ .

Since  $c_G < b$ , assume that  $c_G < d_n$  for all  $n$  without loss of generality. Based on (3.5), we

decompose  $c_n - c_G$  as follows:

$$c_n - c_G = -I_1 - I_3 - I_5 + I_2 + I_4 + I_6, \quad (5.1)$$

where

$$\begin{aligned} I_1 &= \int_{-\theta_0}^{\eta_1} I_{[W_n(x) \leq 0]} dx, & I_2 &= \int_{\eta_2}^{d_n} I_{[W_n(x) > 0]} dx, \\ I_3 &= \int_{\eta_1}^{c_G} I_{[W_n(x) \leq 0, w(x) \leq \beta(\eta_1)]} dx, & I_4 &= \int_{c_G}^{\eta_2} I_{[W_n(x) > 0, w(x) \geq -\beta(\eta_2)]} dx, \\ I_5 &= \int_{\eta_1}^{c_G} I_{[W_n(x) \leq 0, w(x) > \beta(\eta_1)]} dx, & I_6 &= \int_{c_G}^{\eta_2} I_{[W_n(x) > 0, w(x) < -\beta(\eta_2)]} dx. \end{aligned}$$

Note that  $E(c_n - c_G)^2 \leq 2\{E[d_n I_1 + I_3^2 + I_5^2 + d_n I_2 + I_4^2 + I_6^2]\}$ . To prove (3.7.1), we want to show that

$$(5.2.1) \quad \lim_{\epsilon \rightarrow 0} \lim_{n \rightarrow \infty} E[d_n I_1 + I_3^2 + I_5^2] \leq \frac{M}{4[a(c_G)w'(c_G)]^2},$$

and

$$(5.2.2) \quad \lim_{\epsilon \rightarrow 0} \lim_{n \rightarrow \infty} E[d_n I_2 + I_4^2 + I_6^2] \leq \frac{M}{4[a(c_G)w'(c_G)]^2}.$$

We only prove (5.2.2). The proof for (5.2.1) is similar. For  $w(x) < -\beta(x)$ ,  $w_n(x) < 1/2w(x) < 0$  from (vi) of Lemma 5.1. Then we have

$$P(W_n(x) > 0) = P\left(\frac{1}{\sqrt{n\sigma_n^2}} \sum_{j=1}^n Z_{jn} > \frac{-\sqrt{n}w_n(x)}{\sigma_n}\right) \leq P\left(\frac{1}{\sqrt{n\sigma_n^2}} \sum_{j=1}^n Z_{jn} > \frac{-\sqrt{n}w(x)}{2\sigma_n}\right).$$

Applying Theorem 5.16 on page 168 in Petrov (1995) to the left-hand-side of the above inequality,  $P(W_n(x) > 0) \leq [1 - \Phi(\frac{\sqrt{n}|w(x)|}{2\sigma_n})] + \frac{8A\gamma_n}{\sqrt{n}[2\sigma_n + \sqrt{n}|w(x)|]^3} \equiv S_n(x) + T_n(x)$ , where  $A$  is a constant and  $\Phi(\cdot)$  is the cdf of  $N(0, 1)$ . For  $x \in [\eta_2, d_n]$ ,  $w(x) \leq -\epsilon A_\epsilon$ . Then  $w(x) < -\beta(x)$  as  $n$  is large since  $u_n^r d_n \rightarrow 0$ . Then  $P(W_n(x) > 0) \leq S_n(x) + T_n(x)$ . Note that  $\sigma_n^2 \leq l_2(\ln n)^2 u_n^{-4/3}$  and  $\gamma_n \leq l_4(\ln n)^3 u_n^{-8/3}$ . Then  $S_n(x) \leq 1 - \Phi(n^{1/4})$  and  $T_n(x) \leq n^{-1}$  as  $n$  is large. Thus

$$d_n E[I_2] = d_n \int_{\eta_2}^{d_n} P(W_n(x) > 0) dx \leq d_n^2 [1 - \Phi(n^{1/4}) + n^{-1}] = o(n^{-2r/(2r+1)}). \quad (5.3)$$

For  $x \in [c_G, \eta_2]$ ,  $|w'(x)| \geq A_\epsilon$ . Then  $I_4^2 \leq A_\epsilon^{-2} [\int_{c_G}^{\eta_2} I_{[w(x) \geq -\beta(\eta_2)]} w'(x) dx]^2 \leq A_\epsilon^{-2} [\beta(\eta_2)]^2$  by

letting  $y = w(x)/\beta(\eta_2)$ . Therefore

$$\lim_{\epsilon \rightarrow 0} \lim_{n \rightarrow \infty} n^{\frac{2r}{2r+1}} I_4^2 \leq [2B_r B_{c_G}/w'(c_G)]^2. \quad (5.4)$$

By Holder inequality,

$$E[I_6^2] \leq \int_{c_G}^{\eta_2} P(W_n(x) > 0) |w(x)|^3 I_{[w(x) < -\beta(\eta_2)]} dx \times \int_{c_G}^{\eta_2} |w(x)|^{-3} I_{[w(x) < -\beta(\eta_2)]} dx.$$

Similar to  $I_4^2$ ,  $\int_{c_G}^{\eta_2} |w(x)|^{-3} I_{[w(x) > -\beta(\eta_2)]} dx \leq 1/\{2A_\epsilon[\beta(\eta_2)]^2\}$ . Since  $w(x) \leq -\beta(\eta_2) \leq -\beta(x)$  for all  $x \in [c_G, \eta_2]$ ,  $P(W_n(x) > 0) \leq S_n(x) + T_n(x)$  and

$$E[I_6^2] \leq 1/\{2A_\epsilon[\beta(\eta_2)]^2\} \cdot [\int_{c_G}^{\eta_2} S_n(x) |w(x)|^3 dx + \int_{c_G}^{\eta_2} T_n(x) |w(x)|^3 dx]. \quad (5.5)$$

For  $x \in [c_G, \eta_2]$ ,  $\sigma_n^2 \leq m(p_n u_n)^{-1}$ ,  $\gamma_n \leq l_3(p_n u_n)^{-2}$ . Therefore

$$\int_{c_G}^{\eta_2} S_n(x) |w(x)|^3 dx \leq \frac{1}{A_\epsilon} \int_{c_G}^{\eta_2} [1 - \Phi(\frac{\sqrt{nu_n p_n} |w(x)|}{2\sqrt{m}})] |w(x)|^3 w'(x) dx \leq \frac{6m^2}{A_\epsilon (nu_n p_n)^2}, \quad (5.6)$$

and

$$\int_{c_G}^{\eta_2} T_n(x) |w(x)|^3 dx \leq 8Al_3\epsilon/(n^2 u_n^2 p_n^2). \quad (5.7)$$

Combining (5.5), (5.6) and (5.7), we have

$$\lim_{\epsilon \rightarrow 0} \lim_{n \rightarrow \infty} n^{\frac{2r}{2r+1}} E[I_6^2] \leq 3B^2 c_G^2 / [2a(c_G)w'(c_G)]^2, \quad (5.8)$$

Then (5.2.2) follows (5.3), (5.4) and (5.8). Thus (3.7.1) is proved. (3.7.2) can be proved similarly. The details are omitted here. Now we complete the proof of Lemma 3.2.

**5.3. Proof of Lemma 4.1.** Denote  $\bar{\mathcal{C}} = \{\bar{c}_n = c_n^* \vee c_0 \wedge \rho_0 : c_n^* \in \mathcal{C}\}$ . For  $c_n^* \in \mathcal{C}$ ,  $\bar{c}_n = c_n^* \vee c_0 \wedge \rho_0 \in \bar{\mathcal{C}}$ . Define  $\underline{a} = \{a(x) : x \in [c_0, \rho_0]\}$ . Then  $\underline{a} > 0$  and

$$\int_{c_n^*}^{c_G} w(x) a(x) dx \geq \int_{\bar{c}_n}^{c_G} w(x) a(x) dx \geq \underline{a} \int_{\bar{c}_n}^{c_G} w(x) dx = -\frac{\underline{a}}{2} w'(\hat{c}_n) (\bar{c}_n - c_G)^2,$$

where  $\hat{c}_n$  is an intermediate value between  $\bar{c}_n$  and  $c_G$ . Clearly,  $\hat{c}_n \in [c_0, \rho_0]$ . Therefore

$|w'(\hat{c}_n)| \geq L$ . Then  $\int_{c_n^*}^{c_G} w(x) a(x) dx \geq l_1 (\bar{c}_n - c_G)^2$  and

$$\inf_{c_n^* \in \mathcal{C}} \sup_{G \in \mathcal{G}} E[\int_{c_n^*}^{c_G} w(x) a(x) dx] \geq l_1 \inf_{\bar{c}_n \in \bar{\mathcal{C}}} \sup_{G \in \mathcal{G}} E[(\bar{c}_n - c_G)^2].$$

Note that  $\bar{\mathcal{C}} \subset \mathcal{C}$ . Using (2.2),

$$\inf_{\delta_n^* \in \mathcal{D}} \sup_{G \in \mathcal{G}} [R(G, \delta_n^*) - R(G, \delta_G)] \geq l_1 \inf_{\bar{c}_n \in \bar{\mathcal{C}}} \sup_{G \in \mathcal{G}} E(\bar{c}_n - c_G)^2 \geq l_1 \inf_{c_n^* \in \mathcal{C}} \sup_{G \in \mathcal{G}} E(c_n^* - c_G)^2.$$

From the results in Donoho and Liu (1991) (Theorem 3.1 and the remark after Lemma 3.3),

$$\inf_{c_n^* \in \mathcal{C}} \sup_{G \in \mathcal{G}} E(c_n^* - c_G)^2 \geq l_1 \sup \{(c_{f_1} - c_{f_2})^2 : \int [\sqrt{f_1(x)} - \sqrt{f_2(x)}]^2 dx \leq l_2/n, f_1, f_2 \in \mathcal{F}\}.$$

Then Lemma 4.1 is proved.

**5.4. Proof of Lemma 4.2.** Clearly, as  $n$  is large,  $c_0 < c_{f_1} - u_n < c_{f_1} + u_n < \rho_0$ ,  $g_2(\theta) \geq 0$  and  $c_{f_2} \in (c_0, \rho_0)$ . Then  $f_2 \in \mathcal{F}$ . Note that  $\int_0^x H((\theta - c_{f_1})/u_n^{-1})d\theta = 0$  for  $x \leq c_0$  or  $x \geq \rho_0$ , and  $f_1(x) = 0 \implies x < \theta_0 \implies f_2(x) = 0$ . Also for  $G \in \mathcal{G}$ ,  $\int_0^{c_0} dG(\theta)/A(\theta) > 0$ .

Then

$$\begin{aligned} [\sqrt{f_1(x)} - \sqrt{f_2(x)}]^2 &\leq I_{[f_1(x) > 0]} [f_1(x) - f_2(x)]^2 / f_1(x) \\ &\leq l_1 \{u_n^{2r-2} [\int_0^{c_0} \frac{g_1(\theta)}{A(\theta)} d\theta]^{-1} a(x) [\int_{c_0}^x H(\frac{\theta - c_{f_1}}{u_n}) d\theta]^2 + u_n^{2r} \mu_n^2 f_1(x)\}. \end{aligned}$$

Note that  $\mu_n = \int_{-1}^1 A(c_{f_1} + tu_n)H(t)dt = O(u_n)$  and

$$\int_0^\infty a(x) [\int_{c_0}^x H(\frac{\theta - c_{f_1}}{u_n}) d\theta]^2 dx \leq l_3 u_n^3 \int_{-1}^1 a(c_{f_1} + yu_n) [\int_{-1}^y H(t)dt]^2 dy = O(u_n^3).$$

Then we have

$$\int_0^\infty [\sqrt{f_1(x)} - \sqrt{f_2(x)}]^2 dx \leq l_1 [O(u_n^{2r+1}) + O(u_n^{2r+2})] \leq l_2/n.$$

On the other hand, we have  $[w_2(c_{f_1})]^2 = [w_2(c_{f_2}) - w_2(c_{f_1})]^2 = [w_2'(\hat{c}_{f_1})]^2 (c_{f_2} - c_{f_1})^2$ , where  $\hat{c}_{f_1}$  is an intermediate value between  $c_{f_1}$  and  $c_{f_2}$ . It is easy to see that  $[w_2'(\hat{c}_{f_1})]^2 \leq 1/l_1$ .

Then  $(c_{f_2} - c_{f_1})^2 \geq l_1 [w_2(c_{f_1})]^2$ . Note that

$$[w_2(c_{f_1})]^2 \geq l_2 u_n^{2r-2} [\int_0^{c_{f_1}} (\theta_0 - \theta) H(\frac{\theta - c_{f_1}}{u_n}) d\theta]^2 \geq l_2 u_n^{2r} [\int_{-1}^0 (\theta_0 - c_{f_1} + tu_n) H(t) dt]^2$$

and  $\int_{-1}^0 H(t) dt \neq 0$ . Therefore  $(c_{f_2} - c_{f_1})^2 \geq l_3 n^{-\frac{2r}{2r+1}}$ . The proof of Lemma 4.2 is complete

now.

**5.5. Proof of Lemma 5.1.** Noting that  $w(x)$  is decreasing on  $(\theta_0, b)$  and  $w(c_G) = 0$ ,  $|w(x)| \geq |w(c_G - \epsilon)| \wedge |w(c_G + \epsilon)|$  for  $x \in (\theta_0, \eta_1) \cup (\eta_2, b)$ . Since  $w'(x) > A_\epsilon$  for  $x \in (\eta_1, \eta_2)$ ,  $|w(c_G - \epsilon)| \geq A_\epsilon \epsilon$  and  $|w(c_G + \epsilon)| \geq A_\epsilon \epsilon$ . Then  $|w(x)| \geq A_\epsilon \epsilon$ . On the other hand, since  $\phi_G(x) \leq x$  and  $\alpha_G(x) \leq B_r$ ,  $|w(x)| \leq (2\theta_0 + \ln n)B_r$  for  $x \in [\theta_0, d_n]$ . Then (i) is obtained.

With loss of generality, assume that  $u_n \leq \theta_0/2$  for all  $n$ . It is easy to verify that  $w(x) = (\theta_0 - x)\alpha_G(x) + \int_0^x \alpha_G(s)ds$ . A straight forward computation shows that for  $x \in [\theta_0, d_n]$ ,  $|E[V_n(X_j, x)] - w(x)| \leq u_n^r(x - \theta_0 + u_n)B_r B$ . Then (ii) is proved. Note that

$$\sigma_n^2 \leq E[V(X_j, x, n)]^2 = \int_0^1 \frac{1}{u_n a(x - u_n t)} [(\theta_0 - x)K_0(t) - u_n K_1(t)]^2 \alpha_G(x - u_n t) dt.$$

Therefore  $\sigma_n^2 \leq m(p_n u_n)^{-1}$  for  $x \in [\eta_1, \eta_2]$  and  $\sigma_n^2 \leq l_2(\ln n)^2 u_n^{-4/3}$  for  $x \in [\theta_0, d_n]$ . The results for  $\gamma_n$  can be proved similarly. This completes the proofs of (iii) and (iv).

For  $x \in [\theta_0, d_n]$ , if  $w(x) > \beta(x)$ ,

$$\frac{w_n(x)}{w(x)} = \frac{w(x) + [w_n(x) - w(x)]}{w(x)} \geq \frac{w(x) - \beta(x) + 1/2\beta(x)}{w(x) - \beta(x) + \beta(x)} \geq \frac{1}{2}.$$

Then (v) is proved. (vi) can be proved in a similar way.

## References.

- [1] Balakrishnan, N. and Ma, Y. (1997). Convergence rates of empirical Bayes estimation and selection for exponential populations with location parameters. *Advances in Statistical Decision Theory and Applications* (Ed. S. Panchapakesan and N. Balakrishnan) Birkhauser.
- [2] Datta, S. (1994). Empirical Bayes estimation in a threshold model. *Sankhya Ser. A* **56**, 106-117.
- [3] Donoho, D. L. and Liu, R. C. (1991). Geometrizing rates of convergence. II. *Ann. Statist.* **19**, 633-667.
- [4] Huang, S. Y. (1995). Empirical Bayes testing procedures in some nonexponential families using asymmetric Linex loss function. *J. Statist. Plann. Inference* **46**, 293-309.

- [5] Huang, S. Y. and Liang T. (1997). Empirical Bayes estimation of the truncation parameter with linex loss. *Statistica Sinica*. **7**, 755-769.
- [6] Liang, T. (1993). Convergence rates for empirical Bayes estimation of the scale parameter in a Pareto distribution. *Comput. Statist. Data Anal.* **16**, 35-46.
- [7] Ma, Y. and Balakrishnan, N. (2000). Empirical Bayes estimation for truncation parameters. *J. Statist. Plann. Inference* **84**, 111-120.
- [8] Prasad, B. and Singh, R. S. (1990). Estimation of prior distribution and empirical Bayes estimation in a nonexponential family. *J. Statist. Plann. Inference* **24**, 81-86.
- [9] Petrov, V. V. (1995). *Limits Theorems of Probability Theory*. Clarendon Press · Oxford.
- [10] Robbins, H. (1964). The empirical Bayes approach to statistical decision problems. *Ann. Math. Statist.* **35**, 1-20.
- [11] Van Houwelingen, J. C. (1976). Monotone empirical Bayes tests for the continuous one-parameter exponential family. *Ann. Statist.* **4**, 981-989.

REPORT DOCUMENTATION PAGE			Form Approved OMB NO. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comment regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE October 2000	3. REPORT TYPE AND DATES COVERED Technical Reports; October 2000		
4. TITLE AND SUBTITLE Optimal rate of empirical Bayes tests for lower truncation parameters			5. FUNDING NUMBERS DAAD 19-00-1-0502	
6. AUTHOR(S) Shanti S. Gupta and Jianjun Li				
7. PERFORMING ORGANIZATION NAMES(S) AND ADDRESS(ES) Purdue University Department of Statistics West Lafayette, IN 47907-1399			8. PERFORMING ORGANIZATION REPORT NUMBER Technical Report #00-07	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709-2211			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision, unless so designated by other documentation.				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited.			12 b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The distributions with lower truncation parameters are important models in statistics and have been studied in recent years. In this paper, we consider the one-sided testing problem for lower truncation parameters through the empirical Bayes approach. The optimal rate of the monotone empirical Bayes tests is obtained and a monotone empirical Bayes test $\delta_n$ achieving the optimal rate is constructed. It is shown that $\delta_n$ has good performance for both small samples and large samples.				
14. SUBJECT TERMS Empirical Bayes, regret Bayes risk, optimal rate of convergence, minimax.			15. NUMBER OF PAGES 13	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OR REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL	

## GENERAL INSTRUCTIONS FOR COMPLETING SF 298

The Report Documentation Page (RDP) is used in announcing and cataloging reports. It is important that this information be consistent with the rest of the report, particularly the cover and title page. Instructions for filling in each block of the form follow. It is important to ***stay within the lines*** to meet ***optical scanning requirements***.

**Block 1.** Agency Use Only (*Leave blank*)

**Block 2.** Report Date. Full publication date including day, month, and year, if available (e.g. 1 Jan 88). Must cite at least year.

**Block 3.** Type of Report and Dates Covered. State whether report is interim, final, etc. If applicable, enter inclusive report dates (e.g. 10 Jun 87 - 30 Jun 88).

**Block 4.** Title and Subtitle. A title is taken from the part of the report that provides the most meaningful and complete information. When a report is prepared in more than one volume, repeat the primary title, add volume number, and include subtitle for the specific volume. On classified documents enter the title classification in parentheses.

**Block 5.** Funding Numbers. To include contract and grant numbers; may include program element number(s), project number(s), task number(s), and work unit number(s). Use the following labels:

<b>C</b> - Contract	<b>PR</b> - Project
<b>G</b> - Grant	<b>TA</b> - Task
<b>PE</b> - Program Element	<b>WU</b> - Work Unit Accession No.

**Block 6.** Author(s). Name(s) of person(s) responsible for writing the report, performing the research, or credited with the content of the report. If editor or compiler, this should follow the name(s).

**Block 7.** Performing Organization Name(s) and Address(es). Self-explanatory.

**Block 8.** Performing Organization Report Number. Enter the unique alphanumeric report number(s) assigned by the organization performing the report.

**Block 9.** Sponsoring/Monitoring Agency Name(s) and Address(es). Self-explanatory.

**Block 10.** Sponsoring/Monitoring Agency Report Number. (*If known*)

**Block 11.** Supplementary Notes. Enter information not included elsewhere such as; prepared in cooperation with...; Trans. of...; To be published in.... When a report is revised, include a statement whether the new report supersedes or supplements the older report.

**Block 12a.** Distribution/Availability Statement. Denotes public availability or limitations. Cite any availability to the public. Enter additional limitations or special markings in all capitals (e.g. NORFORN, REL, ITAR).

**DOD** - See DoDD 4230.25, "Distribution Statements on Technical Documents."

**DOE** - See authorities.

**NASA** - See Handbook NHB 2200.2.

**NTIS** - Leave blank.

**Block 12b.** Distribution Code.

**DOD** - Leave blank

**DOE** - Enter DOE distribution categories from the Standard Distribution for Unclassified Scientific and Technical Reports

**NASA** - Leave blank.

**NTIS** - Leave blank.

**Block 13.** Abstract. Include a brief (*Maximum 200 words*) factual summary of the most significant information contained in the report.

**Block 14.** Subject Terms. Keywords or phrases identifying major subjects in the report.

**Block 15.** Number of Pages. Enter the total number of pages.

**Block 16.** Price Code. Enter appropriate price code (*NTIS only*).

**Block 17. - 19.** Security Classifications. Self-explanatory. Enter U.S. Security Classification in accordance with U.S. Security Regulations (i.e., UNCLASSIFIED). If form contains classified information, stamp classification on the top and bottom of the page.

**Block 20.** Limitation of Abstract. This block must be completed to assign a limitation to the abstract. Enter either UL (unlimited) or SAR (same as report). An entry in this block is necessary if the abstract is to be limited. If blank, the abstract is assumed to be unlimited.