

Lecture 9: Balanced Incomplete Block Design

Montgomery Section 4.4

Catalyst Experiment

Four catalysts are being investigated in an experiment. The experimental procedure consists of selecting a batch raw material, loading the pilot plant, applying each catalyst in a separate run and observing the reaction time. The batches of raw material are considered as blocks, however each batch is only large enough to permit three catalysts to be run.

Block(raw material)					
Catalyst	1	2	3	4	$y_{i.}$
1	73	74	-	71	218
2	-	75	67	72	214
3	73	75	68	-	216
4	75	-	72	75	222
$y_{.j}$	221	224	207	218	870= $y_{..}$

Balanced Incomplete Block Design (BIBD)

Example 1.

treatment	block					
	1	2	3			
A	A	-	A	1	0	1
B	B	B	-	1	1	0
C	-	C	C	0	1	1

$$a = 3, b = 3, k = 2, r = 2, \lambda = 1$$

Incidence Matrix: $\mathcal{N} = (n_{ij})_{a \times b}$ where $n_{ij} = 1$, if treatment i is run in block j ; $=0$ otherwise.

Example 2.

	block											
treatment	1	2	3	4	5	6						
A	A	A	A	-	-	-	1	1	1	0	0	0
B	B	-	-	B	B	-	1	0	0	1	1	0
C	-	C	-	C	-	C	0	1	0	1	0	1
D	-	-	D	-	D	D	0	0	1	0	1	1

$$a = 4, b = 6, k = 2, r = 3, \lambda = 1, \mathcal{N} = (n_{ij})_{4 \times 6}$$

BIBD: Design Properties

- there are a treatments and b blocks.
- each block contains k (different) treatments.
- each treatment appears in r blocks.
- each pair of treatments appears together in λ blocks.

$a, b, k, r,$ and λ are not independent

- $N = ar = bk$, where N is the total number of runs;

- $\lambda(a - 1) = r(k - 1)$:
 1. for any fixed treatment i_0
 2. two different ways to count the total number of pairs including treatment i_0 in the experiment.
 - I. $a - 1$ possible pairs, each appears in λ blocks, so $\lambda(a - 1)$;
 - II. treatment i_0 appears in r blocks. Within each block, there are $k - 1$ pairs including i_0 , so $r(k - 1)$
- $\lambda < r < b$
- $b \geq a$ (a brainteaser for math/stat students).
- Nonorthogonal design

Extensive list of BIBDs can be found in Fisher and Yates (1963) and Cochran and Cox (1957).

Generating BIBDs by SAS

```

DATA candidates;
  DO treatment = 1 to 4; OUTPUT; END;
PROC OPTEX DATA=candidates SEED=5140514 CODING=ORTH;
  CLASS treatment;
  MODEL treatment;
  BLOCKS STRUCTURE=(6)2; /* (b)k: b=6, k=2 */
  OUTPUT OUT=bibd BLOCKNAME=block;
PROC PRINT DATA=bibd; RUN; QUIT;

```

Obs	BLOCK	treatment	Obs	BLOCK	treatment
1	1	1	7	4	3
2	1	2	8	4	1
3	2	3	9	5	1
4	2	4	10	5	4
5	3	2	11	6	3
6	3	4	12	6	2

BIBD: Statistical Model

- Statistical Model

$$y_{ij} = \mu + \tau_i + \beta_j + \epsilon_{ij} \quad \begin{cases} i = 1, 2, \dots, a \\ j = 1, 2, \dots, b \end{cases}$$

- additive model (without interaction)
- Not all y_{ij} exist because of incompleteness
- Usual treatment and block restrictions : $\sum \tau_i = 0$; $\sum \beta_j = 0$
- Nonorthogonality of treatments and blocks

Use Type III Sums of Squares and lsmeans

Model Estimates

- Least squares estimates for μ , etc.

$$\hat{\mu} = \frac{y_{..}}{N}; \quad \hat{\tau}_i = \frac{kQ_i}{\lambda a}; \quad \hat{\beta}_j = \frac{rQ'_j}{\lambda b}$$

where

$$Q_i = y_{i.} - \frac{1}{k} \sum n_{ij} y_{.j}; \quad Q'_j = y_{.j} - \frac{1}{r} \sum n_{ij} y_{i.}$$

$$\begin{aligned} \text{Var}(Q_i) &= \text{Var}(y_{i.}) + \text{Var}\left(\frac{1}{k} \sum n_{ij} y_{.j}\right) - 2\text{Cov}\left(y_{i.}, \frac{1}{k} \sum n_{ij} y_{.j}\right) \\ &= r\sigma^2 + \frac{r}{k^2} k\sigma^2 - \frac{2}{k} r\sigma^2 \\ &= \frac{(k-1)r}{k} \sigma^2 \end{aligned}$$

- $\text{Var}(\hat{\tau}_i) = \left(\frac{k}{\lambda a}\right)^2 \text{Var}(Q_i) = \left(\frac{k}{\lambda a}\right)^2 \frac{(k-1)r}{k} \sigma^2 = \frac{k(a-1)}{\lambda a^2} \sigma^2$; S.E. $_{\hat{\tau}_i} = ?$
- $\text{Var}(\hat{\tau}_i - \hat{\tau}_j) = \frac{2k\sigma^2}{\lambda a}$; S.E. $_{\hat{\tau}_i - \hat{\tau}_j} = ?$

Analysis of Variance Table (Type I)

Source of Variation	Sum of Squares	Degrees of Freedom	Mean Square	F_0
Blocks	SS_{Block}	$b - 1$	MS_{Block}	
Treatment	$SS_{\text{Treatment}_{\text{Adj}}}$	$a - 1$	$MS_{\text{Treatment}_{\text{Adj}}}$	F_0
Error	SS_E	$N - a - b + 1$	MS_E	
Total	SS_T	$N - 1$		

- $SS_T = \sum \sum y_{ij}^2 - y_{..}^2 / N$
- $SS_{\text{Block}} = \frac{1}{k} \sum y_{.j}^2 - y_{..}^2 / N$

- $SS_{\text{Treatments}_{\text{Adj}}}$ needs adjustment for incompleteness

$$Q_i = y_{i.} - \frac{1}{k} \sum_{j=1}^b n_{ij} y_{.j} \quad \text{where} \quad n_{ij} = \begin{cases} 1 & \text{if trt } i \text{ in blk } j \\ 0 & \text{otherwise} \end{cases}$$

– trt i 's **total** minus trt i 's block averages

– $\sum Q_i = 0$

$$SS_{\text{Treatment}(\text{adjusted})} = k \sum Q_i^2 / \lambda a = \frac{\lambda a}{k} \sum \hat{\tau}_i^2$$

- SS_E by subtraction: $SS_E = SS_{\text{Total}} - SS_{\text{Block}} - SS_{\text{Trt}_{\text{Adj}}}$
- If $F_0 > F_{\alpha, a-1, N-a-b+1}$ then reject H_0

Why don't we adjust the SS_{Block}

- Type I SS:

model reps = block trt;

$$y_{ij} = \mu + \epsilon_{ij} \xleftrightarrow{SS_{\text{Block}}} y_{ij} = \mu + b_j + \epsilon_{ij} \xleftrightarrow{SS_{\text{TrtAdj}}} y_{ij} = \mu + \tau_i + b_j + \epsilon_{ij};$$

- Type I SS:

model reps = trt block;

$$y_{ij} = \mu + \epsilon_{ij} \xleftrightarrow{SS_{\text{Trt}}} y_{ij} = \mu + \tau_i + \epsilon_{ij} \xleftrightarrow{SS_{\text{BlkAdj}}} y_{ij} = \mu + \tau_i + b_j + \epsilon_{ij};$$

- Sequential Sum of Square, guarantee that

$$\begin{aligned} SS_{\text{Model}} &= SS_{\text{Block}} + SS_{\text{TrtAdj}} \\ &= SS_{\text{BlkAdj}} + SS_{\text{Trt}} \end{aligned}$$

- Type III SS

model reps = trt block;

model reps = block trt;

$$\begin{array}{c}
 y_{ij} = \mu + \tau_i + \epsilon_{ij} \quad \overset{SS_{\text{BlkAdj}}}{\longleftrightarrow} \quad y_{ij} = \mu + \tau_i + b_j + \epsilon_{ij} \quad \overset{SS_{\text{TrtAdj}}}{\longleftrightarrow} \quad y_{ij} = \mu + b_j + \epsilon_{ij}; \\
 \Downarrow SS_{\text{Model}} \\
 y_{ij} = \mu + \epsilon_{ij}
 \end{array}$$

- Partial Sum of Square, not guarantee:

$$SS_{\text{Model}} = SS_{\text{BlkAdj}} + SS_{\text{TrtAdj}}$$

- If we want to test blocking effect, we need to use MS_{BlkAdj}/MS_E .
- Just use Type III SS for test.

Mean Tests and Contrasts

- Must compute adjusted means (lsmeans)
- Adjusted mean is $\hat{\mu} + \hat{\tau}_i$
- Standard error of adjusted mean is $\sqrt{\text{MSE} \left(\frac{k(a-1)}{\lambda a^2} + \frac{1}{N} \right)}$
- Contrasts based on adjusted treatment totals

For a contrast: $\sum c_i \mu_i$

Its estimate: $\sum c_i \hat{\tau}_i = \frac{k}{\lambda a} \sum c_i Q_i$

Contrast sum of squares:

$$SS_C = \frac{k \left(\sum_{i=1}^a c_i Q_i \right)^2}{\lambda a \sum_{i=1}^a c_i^2}$$

Pairwise Comparison

- Pairwise comparison $\tau_i - \tau_j$:

1. Bonferroni:

$$CD = t_{\alpha/2m, ar-a-b+1} \sqrt{MS_E \frac{2k}{\lambda a}}$$

2. Tukey:

$$CD = \frac{q_\alpha(a, ar - a - b + 1)}{\sqrt{2}} \sqrt{MS_E \frac{2k}{\lambda a}}$$

SAS Code and output

```
options nocenter ps=60 ls=75;
data example;
  input trt block resp @@;
  datalines;
1 1 73 1 2 74 1 4 71 2 2 75 2 3 67 2 4 72
3 1 73 3 2 75 3 3 68 4 1 75 4 3 72 4 4 75
;

proc glm;
class block trt;
model resp = block trt;
lsmeans trt / tdiff pdiff adjust=bon stderr;
lsmeans trt / pdiff adjust=tukey;
contrast 'a' trt 1 -1 0 0;
estimate 'b' trt 0 0 1 -1;
run;
```

SAS output

Dependent Variable: resp

Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	6	77.75000000	12.95833333	19.94	0.0024
Error	5	3.25000000	0.65000000		
Corrected Total	11	81.00000000			

Source	DF	Type I SS	Mean Square	F Value	Pr > F
block	3	55.00000000	18.33333333	28.21	0.0015
trt	3	22.75000000	7.58333333	11.67	0.0107

Source	DF	Type III SS	Mean Square	F Value	Pr > F
block	3	66.08333333	22.02777778	33.89	0.0010
trt	3	22.75000000	7.58333333	11.67	0.0107

Least Squares Means

Adjustment for Multiple Comparisons: Bonferroni

trt	resp LSMEAN	Standard Error	Pr > t	LSMEAN Number
1	71.3750000	0.4868051	<.0001	1
2	71.6250000	0.4868051	<.0001	2
3	72.0000000	0.4868051	<.0001	3
4	75.0000000	0.4868051	<.0001	4

Bonferroni Method:

i/j	1	2	3	4
1		-0.35806	-0.89514	-5.19183
		1.0000	1.0000	0.0209
2	0.358057		-0.53709	-4.83378
	1.0000		1.0000	0.0284
3	0.895144	0.537086		-4.29669
	1.0000	1.0000		0.0464
4	5.191833	4.833775	4.296689	
	0.0209	0.0284	0.0464	

 Tukey's Method:

i/j	1	2	3	4
1		0.9825	0.8085	0.0130
2	0.9825		0.9462	0.0175
3	0.8085	0.9462		0.0281
4	0.0130	0.0175	0.0281	

Dependent Variable: resp

Contrast	DF	Contrast SS	Mean Square	F Value	Pr > F
c1	1	0.08333333	0.08333333	0.13	0.7349

Parameter	Estimate	Standard		t Value	Pr > t
		Error			
b	-3.00000000	0.69821200		-4.30	0.0077

Other Incomplete Designs

- Youden Square
- Partially Balanced Incomplete Block Design
- Cyclic Designs
- Square, Cubic, and Rectangular Lattices