Statistics 512: Applied Linear Models Topic 4

Topic Overview

This topic will cover

- General Linear Tests (Chapter 7)
- Extra Sums of Squares (Chapter 7)
- Partial Correlations (Chapter 7)
- Multicollinearity (Chapter 7)
- Model Selection (Chapter 9)

General Linear Tests

These are a different way to look at the comparison of models. So far we have looked at comparing/selecting models based on ...

- model significance test and R^2 values
- *t*-tests for variables added last

These are good things to look at, but they are ineffective in cases where ...

- explanatory variables work together in groups
- we want to test some hypotheses for some $\beta_i = b$ rather than $\beta_i = 0$ (for example, maybe we want to test $H_0: \beta_1 = 3, \beta_4 = 7$ against the alternative hypothesis that at least one of those is false)

General Linear Tests look at the difference between models

- in terms of SSE (unexplained SS)
- in terms of SSM (explained SS)

Because SSM + SSE = SST, these two comparisons are equivalent.

The models we compare are *hierarchical* in the sense that one (the full model) includes all of the explanatory variables of the other (the reduced model).

We can compare models with different explanatory variables. For example:

 X_1, X_2 vs X_1 X_1, X_2, X_3, X_4, X_5 vs X_1, X_2, X_3 Note that the first model includes all Xs of the second model.

We will get an *F*-test that compares the two models.

We are testing a null hypothesis that the regression coefficients for the *extra* variables are all zero.

For X_1, X_2, X_3, X_4, X_5 vs X_1, X_2, X_3

$$\begin{aligned} \mathbf{H}_0: & \beta_4 = \beta_5 = 0 \\ \mathbf{H}_a: & \beta_4 \text{ and } \beta_5 \text{ are not both } 0 \end{aligned}$$

F-test

The test statistic in general is

$$F = \frac{\left(SSE(R) - SSE(F)\right) / \left(df_E(R) - df_E(F)\right)}{SSE(F)/df_E(F)}$$

Under the null hypothesis (reduced model) this statistic has an F distribution where the degrees of freedom are the number of *extra* variables and the df_E for the larger model. So we reject if the *p*-value for this test is $p \leq 0.05$, and in that case conclude that at least one of the extra variables is useful for predicting Y in the linear model that already contains the variables in the reduced model.

Example

Suppose n = 100 and we are testing X_1 , X_2 , X_3 , X_4 , X_5 (full) vs X_1 , X_2 , X_3 (reduced). Our hypotheses are:

$$\begin{aligned} \mathbf{H}_0: & \beta_4 = \beta_5 = 0 \\ \mathbf{H}_a: & \beta_4 \text{ and } \beta_5 \text{ are not both } 0 \end{aligned}$$

Since we are considering removing 2 variables $(X_4 \text{ and } X_5)$, the numerator df is 2. The denominator df is n-6 = 94 (since p = 6 for the full model). We reject if the *p*-value ≤ 0.05 and in that case would conclude that either X_4 or X_5 or both contain additional information that is useful for predicting Y in a linear model that also includes X_1 , X_2 , and X_3 .

Extra Sums of Squares

Notation for Extra SS

Example using 5 variables:

- $SSE(X_1, X_2, X_3, X_4, X_5)$ is the SSE for the full model, SSE(F)
- $SSE(X_1, X_2, X_3)$ is the SSE for a reduced model, SSE(R).

- The "extra sum of squares" for this comparison is denoted $SSM(X_4, X_5|X_1, X_2, X_3)$.
- This is the difference in the SSE's:

$$SSM(X_4, X_5 | X_1, X_2, X_3) = SSE(R) - SSE(F)$$

= $SSE(X_1, X_2, X_3) - SSE(X_1, X_2, X_3, X_4, X_5)$

• We call this an "extra sum of squares" because it is the extra amount of SS explained by the full model that is not explained by the reduced model. In this case it is the extra sums of squares explained by X_4 and X_5 , given that X_1 , X_2 , and X_3 are already in the model.

For each model, SST = SSM + SSE, so SSE = SST - SSM, and SST is the same for both models. It follows that

$$SSM(X_4, X_5 | X_1, X_2, X_3)$$

$$= SSE(X_1, X_2, X_3) - SSE(X_1, X_2, X_3, X_4, X_5)$$

$$= [SST - SSM(X_1, X_2, X_3)] - [SST - SSM(X_1, X_2, X_3, X_4, X_5)]$$

$$= SSM(X_1, X_2, X_3, X_4, X_5) - SSM(X_1, X_2, X_3)$$

$$= SSM(F) - SSM(R)$$

Thus, the extra sums of squares in the numerator of the test may also be written as the difference in the SSM's (full - reduced).

Similarly, since $df_M + df_E = df_T$ for each model, we have

$$df_E(R) - df_E(F) = [df_T - df_M(R)] - [df_T - df_M(F)] = df_M(F) - df_M(R)$$

Thus the df in the numerator of the test may also be written as the difference in the df_M 's (full - reduced).

It doesn't matter whether you look at as the difference in the models or the difference in the errors, as long as you get the signs right.

F-test (General Linear Test)

Numerator: $SSM(X_4, X_5|X_1, X_2, X_3)/2$ (where 2 = # variables before "|") Denominator: $MSE(X_1, X_2, X_3, X_4, X_5)$ (full model MSE) $F \sim F_{2,n-6}$ Numerator df = 2 since two parameters fixed, i.e. difference in parameters = full - reduced. Denominator df = n - 6 since p = 6 for full model.

Reject if the *p*-value ≤ 0.05 , and in that case would conclude that either X_4 or X_5 or both contain additional information that is useful for predicting Y in a linear model that also includes X_1 , X_2 , and X_3 .

Example

Predict bone density using age, weight and height; does diet add any useful information? Predict GPA using 3 HS grade variables; do SAT scores add any useful information? Predict yield of an industrial process using temperature and pH; does the supplier of the raw material (categorical) add any useful information?

Extra SS Special Cases (SAS)

Type II SS: Each variable given all others are in the model

- Compare models that differ by one explanatory variable, $F_{1,n-p} = t_{n-p}^2$ (i.e. 'variable added last' tests).
- SAS's individual parameter t-tests are equivalent to the general linear test based on $SSM(X_i|X_1, \ldots, X_{i-1}, X_{i+1}, \ldots, X_n)$
- These are called Type II SS (not related to Type II Error).
- Type II SS are the extra sums of squares for each variable, given that all other variables are in the model.
- Option SS2 to proc reg.

Type I SS: Add one variable at a time in order

Consider 4 explanatory variables *in a particular order* and the extra sums of squares we get by adding one variable at a time, starting with the first one (SAS takes them in the order listed in the model statement), and testing each successive model with the one before it...

- $SSM(X_1)$
- $SSM(X_2|X_1)$
- $SSM(X_3|X_1, X_2)$
- $SSM(X_4|X_1, X_2, X_3)$

It is easy to work out from the definitions that these sum to the SSM for the full model:

$$SSM(X_1) + SSM(X_2|X_1) + SSM(X_3|X_1, X_2) + SSM(X_4|X_1, X_2, X_3)$$

= SSM(X_1, X_2, X_3, X_4)

For the tests,

• Numerator df is 1 for all of them (one parameter before "|").

- Denominator df depends on which model acts as the full one.
- $F = (SSR/1)/MSE(F) \sim F_{1,n-p}$
- These are called Type I SS (not related to Type I Error). Notice that the Type I SS depend on the order in which the variables are listed.
- Type I SS are the extra SS for each variable, given that *all previous* variables are in the model. ("previous" meaning having a lower subscript number)

KNNL Example page 256

- SAS code in nknw260.sas
- 20 healthy female subjects ages 25-34
- Y is fraction body fat
- X_1 is triceps skin fold thickness
- X_2 is thigh circumference
- X_3 is midarm circumference
- Goal is to find a good model based on these three easy measurements. Otherwise we will have to use underwater weight/density measurements (difficult and more expensive).

Step 1: Check the data and run proc reg on full model

```
data bodyfat;
    infile 'H:\System\Desktop\Ch07ta01.dat';
    input skinfold thigh midarm fat;
proc print data=bodyfat;
proc reg data=bodyfat;
    model fat=skinfold thigh midarm;
```

			Sum of	Mean		
Source		DF	Squares	Square	F Value	Pr > F
Model		3	396.98461	132.32820	21.52	<.0001
Error		16	98.40489 6.150			
Corrected To	tal	19	495.38950			
Root MSE		2.47998	R-Square	0.8014		
Something is	s usefi	ıl.				
		Parameter	Standard			
Variable	DF	Estimate	Error	t Value	Pr > t	
Intercept	1	117.08469	99.78240	1.17	0.2578	
skinfold	1	4.33409	3.01551	1.44	0.1699	
thigh	1	-2.85685	2.58202	-1.11	0.2849	
midarm	1	-2.18606	1.59550	-1.37	0.1896	

But what? None of the *p*-values are significant.

Dilemma

The *p*-value for $F_{3,16}$ is 7×10^{-6} , but the *p*-values for the individual regression coefficients are 0.1699, 0.2849, and 0.1896. None of these are near our standard of 0.05. We have a seeming contradiction. What is the reason?

Look at the Extra SS. Here's how to do it in SAS.

proc reg data=bodyfat; model fat=skinfold thigh midarm /ss1 ss2; Pr > |t|Variable Type I SS Type II SS skinfold 0.1699 352.26980 12.70489 0.2849 thigh 33.16891 7.52928 midarm 0.1896 11.54590 11.54590

Compare these values to the model SS

Model	396.98461
Error	98.40489
Corrected Total	495.38950

Interpretation

Type I:

- SSM(skinfold) = 352.26
- SSM(thigh|skinfold) = 33.16
- SSM(midarm|skinfold, thigh) = 11.54

Type II:

- SSM(skinfold|thigh, midarm) = 12.70
- SSM(thigh|skinfold, midarm) = 7.52

Skinfold accounts for a lot of the SS by itself, but if thigh and midarm are included first then skinfold is redundant.

Notes

- Type I and Type II SS are very different for this example (except for the last one).
- The order in which we list the variables in the model statement affects the Type I SS, but does not affect the Type II SS.
- If we reorder the variables in the model statement we will get different Type I SS but the same Type II SS

Could variables be explaining the same SS and "canceling each other out"? We suspect so because the Type I and II SS are so different.

We also observe parameter estimates that do not "make sense": the regression coefficients for thigh and midarm are negative. (Large thighs = low body fat?)

Run additional models to clarify the situation (more on this later).

Also look at pairwise relationships, keeping in mind that more complicated relationships may exist.

proc corr data=bodyfat noprob;

Pearson Correlation Coefficients, $N = 20$					
	skinfold	thigh	midarm	fat	
skinfold	1.00000	0.92384	0.45778	0.84327	
thigh	0.92384	1.00000	0.08467	0.87809	
midarm	0.45778	0.08467	1.00000	0.14244	
fat	0.84327	0.87809	0.14244	1.00000	

The correlation between skinfold and thigh is extremely high, so we most likely do not want to have both of these in our model. Both of these are also highly correlated with fat, which means they will be good predictors.



Rerun with single explanatory variables

proc reg data=bodyfat;

model	fat =	skinfold;			
model	fat =	thigh;			
model	fat =	midarm;			
		Parameter	Standard		
Variable	DF	Estimate	Error	t Value	Pr > t
Intercept	1	-1.49610	3.31923	-0.45	0.6576
skinfold	1	0.85719	0.12878	6.66	<.0001
Root MSE		2.81977	R-Square	0.7111	
Intercept	1	-23.63449	5.65741	-4.18	0.0006
thigh	1	0.85655	0.11002	7.79	<.0001
Root MSE		2.51024	R-Square	0.7710	
Intercept	1	14.68678	9.09593	1.61	0.1238
midarm	1	0.19943	0.32663	0.61	0.5491
Root MSE		5.19261	R-Square	0.0203	

test statement in proc reg

This does any general linear test you specify. Note that the labels before the ":" are just names picked to identify the tests; they are useful, since you can do more than one test here and need to distinguish the output. For example, the command test thigh, midarm; tests $H_0: \beta_2 = \beta_3 = 0.$

```
proc reg data=bodyfat;
model fat=skinfold thigh midarm;
skinonly: test thigh, midarm;
thighonly: test skinfold, midarm;
skinmid: test thigh;
```

Test	skinonly	Results	for	Dependent Mean	Variable	fat
Source		DF		Square	F Value	Pr > F
Numerator		2	22	2.35741	3.64	0.0500
Denominat	or	16	6	6.15031		

This is really borderline – probably should not remove both thigh and midarm.

Test thighonly Results for Dependent Variable fat

		Mean		
Source	DF	Square	F Value	Pr > F
Numerator	2	7.50940	1.22	0.3210
Denominator	16	6.15031		

This says a model with thigh only would probably be okay.

Test	skinmid	Results	for	Dependent	Variable	fat
				Mean		
Source		DF		Square	F Value	Pr > F
Numerator		1	-	7.52928	1.22	0.2849
Denominato	or	16	(6.15031		

This says a model with skinfold and midarm only, but not thigh, would probably be okay.

How to pick the very best model depends on your objectives. If your goal is only "blackbox" prediction, with no interpretation on the coefficients, you might want to include some variables that are only marginally helpful. But if your goal is to actually interpret the coefficient, you will also want to look at whether the coefficients make sense. For example, in the full model, the parameter estimates for thigh and midarm are both negative. This says, the larger your thighs and arms, the smaller your bodyfat, which doesn't make much sense. This happens because skinfold is already in the model, and so the positive influence of thigh on fat has already been included.

Other uses

The **test** statement can be used to perform a significance test for any hypothesis involving a linear combination of the regression coefficients. Examples:

$$\begin{split} \mathrm{H}_{0}: & \beta_{4} = \beta_{5} \; (\texttt{test X4 = X5}) \\ \mathrm{H}_{0}: & \beta_{4} - 3\beta_{5} = 12 \; (\texttt{test X4-3*X5=12}) \end{split}$$

Separate equations by commas; if no = sign appears then = 0 is assumed.

Chapter 7.4: Partial correlations (r) - Coefficients of Partial Determination (r^2)

Partial correlation measures the strength of a linear relation between two variables taking into account other variables.

Recall from SLR that $r^2 = SSM/SST$. In contrast, now we are measuring the marginal increase in SS explained by including a new variable in the model, compared to the SSE. See page 270 for formulas.

This is closely related to extra sums of squares.

Possible procedure to find partial correlation X_i , Y

This is not how you will actually calculate it, but it may help you visualize what is going on

- Predict Y using other X's
- Predict X_i using other X's
- Find correlation between the two sets of residuals

• Plotting the two sets of residuals vs each other on a scatterplot gives a partial correlation plot (later).

KNNL use the term *coefficient of partial determination* for the squared partial correlation r^2 .

Partial correlations in SAS

```
proc reg data=bodyfat;
   model fat=skinfold thigh midarm / pcorr1 pcorr2;
                      Parameter Estimates
                  Squared Partial Squared Partial
Variable
           Pr > |t| Corr Type I Corr Type II
Intercept
           0.2578
skinfold
            0.1699
                         0.71110
                                         0.11435
thigh
            0.2849
                         0.23176
                                         0.07108
midarm
            0.1896
                         0.10501
                                         0.10501
```

These use the Type I and Type II SS for computation:

- Type I Squared Partial Correlation uses Type I SS: $r^2 = \frac{SS1}{SS1+SSE}$
- Type II Squared Partial Correlation uses Type II SS: $r^2 = \frac{SS2}{SS2+SSE}$

Where e.g. for thigh, SS2 = SSM(thigh|skinfold, midarm), and SSE = SSE(skinfold, thigh, midarm). We can similarly define partial correlations as the square root. These give roughly the same information as Extra SS but are scaled

These give roughly the same information as Extra SS but are scaled.

Multicollinearity

Back to body fat example (KNNL page 256)

The *p*-value for ANOVA *F*-test is < .0001.

The p-values for the individual regression coefficients are 0.1699, 0.2849, and 0.1896.

None of these are near our standard significance level of 0.05.

What is the explanation?

Multicollinearity!!!

Numerical analysis problem is that the matrix X'X is close to singular and is therefore difficult to invert accurately.

Statistical problem is that there is too much correlation *among the explanatory variables*, and it is therefore difficult to determine the regression coefficients.

If we solve the statistical problem then the numerical problem will also be solved. Our general goal is to refine a model that has redundancy in the explanatory variables. And we need to do this whether or not X'X can be inverted without difficulty.

Extreme cases can help us understand the problems caused by multicollinearity.

- One extreme: Assume columns in X matrix were uncorrelated (r = 0). In that case Type I and Type II SS will be the same. The contribution of each explanatory variable to the model is the same whether or not the other explanatory variables are in the model. There is no overlap in the variation components explained by each variable.
- Other extreme: Suppose a linear combination of the explanatory variables is a constant (maybe a variable was accidentally included twice in the data set such that $X_1 = X_2$). The Type II SS for the X's involved will be zero because when one is included the other is redundant (it explains NO additional variation over the other variables).

Back to CS example: We did not have a big multicollinearity problem. Let's fake a linear combination to illustrate a point.

```
data cs;
    infile 'H:\System\Desktop\csdata.dat';
    input id gpa hsm hss hse satm satv genderm1;
data cs; set cs;
    sat=satm + satv;
proc reg data=cs;
    model gpa=sat satm satv;
Sum of Mean
```

		hean		
DF	Squares	Square	F Value	Pr > F
2	8.58384	4.29192	7.48	0.0007
221	126.87895	0.57411		
223	135.46279			
	DF 2 221 223	DF Squares 2 8.58384 221 126.87895 223 135.46279	DF Squares Square 2 8.58384 4.29192 221 126.87895 0.57411 223 135.46279	DF Squares Square F Value 2 8.58384 4.29192 7.48 221 126.87895 0.57411 223 135.46279

Something is wrong; $df_M = 2$, but there are 3 X's. What is going on?

0

Error messages from SAS:

0

satv

NOTE: Model is not full rank. Least-squares solutions for the parameters are not unique. Some statistics will be misleading. A reported DF of 0 or B means that the estimate is biased. NOTE: The following parameters have been set to 0, since the variables are a linear combination of other variables as shown. satv = sat - satm Parameter Standard DF Error t Value Pr > |t|Variable Estimate Intercept 1 1.28868 0.37604 3.43 0.0007 В -0.00002456 -0.04 0.9684 0.00061847 sat В 0.00231 0.00110 2.10 0.0365 satm

SAS tells us the design matrix is not of full rank, i.e. its rank is < p. This means one column is a linear combination of some other columns. This in turn means that the determinant of $\mathbf{X'X}$ is 0, and thus $\mathbf{X'X}$ is not invertible.

.

.

Extent of Multicollinearity

This example had one explanatory variable equal to a linear combination of other explanatory variables. This is the most extreme case of multicollinearity and is detected by statistical software because $\mathbf{X}'\mathbf{X}$ does not have an inverse. We are mostly concerned with less extreme cases where $\mathbf{X}'\mathbf{X}$ does have an inverse, though the determinant may be quite small.

Effects of Multicollinearity

What happens when there is multicollinearity in the X's?

- Regression coefficients are not well estimated and may be meaningless, and similarly for standard errors of these estimates. (see chart on page 286)
- Type I SS and Type II SS will differ.
- R^2 and predicted values are usually okay.

Pairwise Correlations

Pairwise correlations can be used to check for "pairwise" collinearity. This is useful but note that they do not show more complicated linear dependence. KNNL p256

$$Corr(skinfold, thigh) = 0.9238$$

 $Corr(skinfold, midarm) = 0.4578$
 $Corr(thigh, midarm) = 0.0847$

But this is not the whole story. For example, if we think of skinfold as a Y variable and model it with both midarm and thigh as explanatory variables, we get a very large $R^2 = 0.9986$, meaning that skinfold is almost exactly predicted from the other two. Simple pairwise correlations do not pick up on this, but the regression results tell us something like this was going on.

Similarly in the example with sat the problem was with a set of three variables: sat, satv, and satm. Any two of these would be okay, but three is impossible. We would not have been aware of the extent of this problem just from the pairwise correlations.

```
proc corr data=cs noprob;
var sat satm satv;
```

	sat	satm	satv
sat	1.00000	0.84450	0.86623
satm	0.84450	1.00000	0.46394
satv	0.86623	0.46394	1.00000

Polynomial Regression

We can fit a quadratic, cubic, etc. relationship by defining squares, cubes, etc., in a data step and using them as additional explanatory variables.

We can also do this with more than one explanatory variable, in which case we also often include an interaction term.

When we do this we generally create a multicollinearity problem, which can often be corrected by standardization.

Warning: You know that with linear regression models it is a bad idea to extrapolate beyond the scope of the data. In polynomial regression it is a VERY bad idea.

KNNL Example, page 300 (nknw302.sas)

Response variable is the life (in cycles) of a power cell. Explanatory variables are

- Charge rate (3 levels)
- Temperature (3 levels)

This is a designed experiment: levels of charge and temperature are planned. The experimenter wants to know if a linear or quadratic function is appropriate, and if an interaction between charge rate and temperature should be included in the model. (An interaction means that the way in which charge rate affects the life of the cell depends on the temperature; a model with no interaction is called *additive*. MUCH more on this in ANOVA.)

Input and check the data.

```
data powercell;
    infile 'H:\System\Desktop\CH07TA09.DAT';
    input cycles chrate temp;
    proc print data=powercell;
```

Obs	cycles	chrate	temp
1	150	0.6	10
2	86	1.0	10
3	49	1.4	10
4	288	0.6	20
5	157	1.0	20
6	131	1.0	20
7	184	1.0	20
8	109	1.4	20
9	279	0.6	30
10	235	1.0	30
11	224	1.4	30

Use a data step to create new variables and run the regression with polynomial and interaction terms. data powercell; set powercell; chrate2=chrate*chrate; temp2=temp*temp; ct=chrate*temp; proc reg data=powercell; model cycles=chrate temp chrate2 temp2 ct / ss1 ss2;

		Sum of	Ν	lean			
	DF	Squares	Squ	lare F	Value	Pr > F	
	5	55366	11	L073	10.57	0.0109	
	5	5240.43860	1048.08	3772			
Total	10	60606					
	Parameter	Standard					
DF	Estimate	Error	t Value	Pr > t	l Ty	pe I SS	Type II SS
1	337.72149	149.96163	2.25	0.074	1	325424	5315.62944
1	-539.51754	268.86033	-2.01	0.101	1	18704	4220.41673
1	8.91711	9.18249	0.97	0.376	1	34202	988.38036
1	171.21711	127.12550	1.35	0.2359	9 164	5.96667	1901.19474
1	-0.10605	0.20340	-0.52	0.6244	4 28	4.92807	284.92807
1	2.87500	4.04677	0.71	0.5092	2 52	9.00000	529.00000
	Total DF 1 1 1 1 1 1	DF 5 5 Total 10 Parameter DF Estimate 1 337.72149 1 -539.51754 1 8.91711 1 171.21711 1 -0.10605 1 2.87500	Sum of DF Squares 5 55366 5 5240.43860 Total 10 60606 Parameter Standard DF Estimate Error 1 337.72149 149.96163 1 -539.51754 268.86033 1 8.91711 9.18249 1 171.21711 127.12550 1 -0.10605 0.20340 1 2.87500 4.04677	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$

Conclusion

Overall F is significant, but no individual t's are significant \rightarrow multicollinearity problem. Also notice the extreme difference between Type I and Type II SS for chrate and temp. Look at the correlations (proc corr) for a clue :

	Pearson	Correlation	Coefficients,	N = 11	
	chrate	temp	chrate2	temp2	ct
chrate	1.00000	0.00000	0.99103	0.00000	0.60532
temp	0.00000	1.00000	0.00000	0.98609	0.75665
chrate2	0.99103	0.00000	1.00000	0.00592	0.59989
temp2	0.00000	0.98609	0.00592	1.00000	0.74613
ct	0.60532	0.75665	0.59989	0.74613	1.00000

There are some very high correlations

r(chrate, chrate2) = 0.99103r(temp, temp2) = 0.98609

Correlation between powers of variables are causing a big multicollinearity problem.

A remedy

We can remove or reduce the correlation between explanatory variables and their powers by *centering*. Centering means that you subtract off the mean from each variable. We often rescale by standardizing (subtract the mean and divide by the standard deviation) but subtracting the mean is the key. Use proc standard to center the explanatory variables. (proc standard will standardize a dataset to any given mean and std dev; we set a mean of 0 to center the variables. To set a given standard deviation, we could also say std=1, for example. The same names are used for the standardized variables.)

Recompute the squares, cubes, etc., using the centered variables, and rerun the regression analysis.

Note the use of "drop" in the data step. (opposite is "keep")

```
data copy; set powercell;
   schrate=chrate; stemp=temp;
   drop chrate2 temp2 ct;
*center the variables and put them in
    dataset std;
proc standard data=copy out=std mean=0;
   var schrate stemp;
* schrate and stemp now have mean 0;
proc print data=std;
```

Obs	cycles	chrate	temp	schrate	stemp
1	150	0.6	10	-0.4	-10
2	86	1.0	10	0.0	-10
3	49	1.4	10	0.4	-10
4	288	0.6	20	-0.4	0
5	157	1.0	20	0.0	0
6	131	1.0	20	0.0	0
7	184	1.0	20	0.0	0
8	109	1.4	20	0.4	0
9	279	0.6	30	-0.4	10
10	235	1.0	30	0.0	10
11	224	1.4	30	0.4	10

Now both schrate and stemp have mean 0.

Recompute squares and interaction term using standardized variables.

```
data std; set std;
schrate2=schrate*schrate;
stemp2=stemp*stemp;
sct=schrate*stemp;
```

Rerun regression...

proc reg data=std; model cycles= chrate temp schrate2 stemp2 sct / ss1 ss2;

Parameter Estimates							
		Parameter	Standard				
Variable	DF	Estimate	Error	t Value	Pr > t	Type I SS	Type II SS
Intercept	1	151.42544	45.45653	3.33	0.0208	325424	11631

chrate	1	-139.58333	33.04176	-4.22	0.0083	18704	18704
temp	1	7.55000	1.32167	5.71	0.0023	34202	34202
schrate2	1	171.21711	127.12550	1.35	0.2359	1645.96667	1901.19474
stemp2	1	-0.10605	0.20340	-0.52	0.6244	284.92807	284.92807
sct	1	2.87500	4.04677	0.71	0.5092	529.00000	529.00000

Now we can see that the linear terms are significant and the quadratic terms are not. See book: general linear test shows that quadratic and interaction terms can be omitted. Things are much more clear now and we can trust our results since the correlations among X's are virtually eliminated.

Type I and Type II SS are almost the same.

proc corr data=std noprob; var chrate temp schrate2 stemp2 sct; Pearson Correlation Coefficients, N = 11schrate2 chrate stemp2 temp sct 1.00000 0.00000 0.00000 0.00000 0.00000 chrate 1.00000 0.00000 0.00000 0.00000 temp 0.00000 schrate2 0.00000 0.00000 1.00000 0.26667 0.00000 stemp2 0.00000 0.00000 0.26667 1.00000 0.00000 0.00000 0.00000 0.00000 0.00000 1.00000 sct

In this particular example, the correlations went to zero after centering. That happened because all possible combinations were equally represented (in ANOVA terms, a balanced design). Correlations will not always go to zero after centering, but they should be reduced. A typical quadratic regression model with two X variables would be

$$Y_{i} = \beta_{0} + \beta_{1}X_{i,1} + \beta_{2}X_{i,2} + \beta_{1,1}(X_{i,1} - \bar{X}_{1})^{2} + \beta_{2,2}(X_{i,2} - \bar{X}_{2})^{2} + \beta_{1,2}(X_{i,1} - \bar{X}_{1})(X_{i,2} - \bar{X}_{2}) + \epsilon_{i}$$

Notice the alternate convention for subscripts sometimes used on the regression coefficients.

Comment on General Linear Test and Extra SS

The test for all second-order terms = 0 uses SSM(schrate2, stemp2, sct|chrate, temp). This is not given directly from our SAS output. We can use the usual test statement in proc reg:

second: test schrate2, stemp2, sct; Test second Results for Dependent Variable cycles Mean DF Source Square F Value Pr > FNumerator 3 819.96491 0.78 0.5527 Denominator 5 1048.08772

But we can also get the necessary SS from SS1 with a little work:

SSM(schrate2, stemp2, sct|schrate, stemp) = SSM(full) - SSM(schrate, stemp)Notice that the Type I SS are

SS1(schrate2) = SSM(schrate2|chrate, temp)= SSM(chrate, temp, schrate2) - SSM(chrate, temp)= SSM(stemp2|chrate, temp, schrate2)SS1(stemp2)= SSM(chrate, temp, schrate2, stemp2) - SSM(chrate, temp, schrate2)SS1(sct) = SSM(sct, chrate, temp, schrate2, stemp2)= SSM(chrate, temp, schrate2, stemp2, sct) - SSM(chrate, temp, schrate2, stemp2)SS1(schrate2) + SS1(stemp2) + SS1(sct)= SSM(chrate, temp, schrate2, stemp2, sct) - SSM(chrate, temp)

= SSM(schrate2, stemp2, sct|chrate, temp)

Thus, we can get the required extra SS from adding certain Type I SS. This only works if the second-order terms (in general, the terms to be tested) are listed last in the model statement.

Interaction Models

With several explanatory variables, we need to consider the possibility that the effect of one variable depends on the value of another variable.

Special cases:

- One binary variable (0/1) and one continuous variable (Section 8.3 doing this out of sequence)
- Two continuous variables

First Special Case: One binary variable and one continuous variable

(From Section 8.3)

 X_1 takes values 0 and 1 corresponding to two different groups or categories.

 X_2 is a continuous variable.

Model: $Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_1 X_2 + \epsilon$

As you will see, this is a convenient way of writing down two separate SLR models for the two categories.

When $X_1 = 0$, the model becomes

$$Y = \beta_0 + \beta_1(0) + \beta_2 X_2 + \beta_3(0) X_2 + \epsilon = \beta_0 + \beta_2 X_2 + \epsilon$$

This is a SLR model for Y as a function of X_2 with intercept β_0 and slope β_2 .

When $X_1 = 1$, the model becomes

$$Y = \beta_0 + \beta_1(1) + \beta_2 X_2 + \beta_3(1) X_2 + \epsilon$$

= $\beta_0 + \beta_1 + \beta_2 X_2 + \beta_3 X_2 + \epsilon$
= $(\beta_0 + \beta_1) + (\beta_2 + \beta_3) X_2 + \epsilon$

This is a SLR model for Y as a function of X_2 with intercept $(\beta_0 + \beta_1)$ and slope $(\beta_2 + \beta_3)$.

These are both SLR models, and we could model them separately (make one data set for each group). But we can more easily model the combination of the two with a multiple regression model using a *binary* (0-1) variable, and this allows us to look at both groups together. It has the added benefit of using all the data together to get the variance estimate, increasing our error degrees of freedom. Some useful tests in this situation include

- $H_0: \beta_1 = \beta_3 = 0$ is the hypothesis that the regression lines are the same.
- $H_0: \beta_1 = 0$ is the hypothesis that the two intercepts are equal.
- $H_0: \beta_3 = 0$ is the hypothesis that the two slopes are equal.

KNNL Example, page 315

(nknw459.sas)

Y is number of months it takes for an insurance company to adopt an innovation.

 X_1 is the type of firm (a qualitative or categorical variable): X_1 takes the value 0 if it is a mutual fund firm and 1 if it is a stock fund firm. X_2 is the size of the firm (a continuous variable).

We ask whether stock firms adopt the innovation slower or faster than mutual firms. We ask the question across all firms, regardless of size.

```
data insurance;
   infile 'H:\System\Desktop\Ch11ta01.dat';
   input months size stock;
proc print data=insurance;
Obs
       months
                          stock
                  size
         17
                            0
  1
                   151
         26
                    92
                            0
  2
. . .
         30
                   124
                            1
 19
 20
         14
                   246
                            1
```

Plot the data with two symbols:

```
symbol1 v=M i=sm70 c=black l=1;
symbol2 v=S i=sm70 c=black l=3;
title1 'Insurance Innovation';
proc sort data=insurance;
by stock size;
title2 'with smoothed lines';
proc gplot data=insurance;
   plot months*size=stock;
```

Note the use of a new syntax in the **plot** statement to plot separate lines for different values of the categorical variable stock.



Interaction Effects

Interaction expresses the idea that the effect of one explanatory variable on the response depends on another explanatory variable.

In this example, this would mean that the slope of the line depends on the type of firm. Are both lines the same?

Use the **test** statement to test whether $\beta_1 = \beta_3 = 0$. Make the X_3 variable which is the product of stock (X_1) and size (X_2) .

Model	3	1504.41904	501.47301	45.49	<.0001
Error	16	176.38096	11.02381		
Corrected Total	19	1680.80000			
Root MSE	3.32021	R-Square	0.8951		

Test	sameline	Results	for Dependent	Variable	months
			Mean		
Source		DF	Square	F Value	Pr > F
Numerato	r	2	158.12584	14.34	0.0003
Denomina	tor	16	11.02381		

The lines are not the same. How are they different?

		Parameter	Standard		
Variable	DF	Estimate	Error	t Value	Pr > t
Intercept	1	33.83837	2.44065	13.86	<.0001
stock	1	8.13125	3.65405	2.23	0.0408
size	1	-0.10153	0.01305	-7.78	<.0001
sizestock	1	-0.00041714	0.01833	-0.02	0.9821

It appears we can ignore β_3 : Two parallel lines.

proc reg data=insurance;

model months = stock size;

		An	alysis of Varia	ance		
			Sum of	Mean		
Source		DF	Squares	Square	F Value	Pr > F
Model		2	1504.41333	752.20667	72.50	<.0001
Error		17	176.38667	10.37569		
Corrected	Total	19	1680.80000			
Root MSE		3.22113	R-Square	0.8951		
Dependent	Mean	19.40000	Adj R-Sq	0.8827		
Coeff Var		16.60377				
		Paramet	er Estimates			
		Parameter	Standard			
Variable	DF	Estimate	Error	t Value	Pr > t	
Intercept	1	33.87407	1.81386	18.68	<.0001	
size	1	-0.10174	0.00889	-11.44	<.0001	
stock	1	8.05547	1.45911	5.52	<.0001	

The two lines have different intercepts, but it is safe to assume they have the same slope. For mutual fund firms, $X_1 = 0$, so the model is $Y = \beta_0 + \beta_2 X_2 + \epsilon$; the fitted equation is $\hat{Y} = 33.784 - 0.102X_2$.

For stock firms, $X_1 = 1$, so the model is $Y = \beta_0 + \beta_1 + \beta_2 X_2 + \epsilon$; the estimated intercept is (33.874 + 8.055) = 41.93, so the fitted equation is $\hat{Y} = 41.93 - 0.102X_2$.

Plot with two regression lines (note this manner of plotting will permit different slopes;

to get the actual lines given by the model we would have to overlay the predicted values on the plot, as we did in Homework 3).

symbol1 v=M i=rl c=black; symbol2 v=S i=rl c=black; proc gplot data=insurance; plot months*size=stock;



Second Special Case: Two Continuous Variables

The model $Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_1 X_2 + \epsilon$ can be rewritten as follows:

$$Y = \beta_0 + (\beta_1 + \beta_3 X_2) X_1 + \beta_2 X_2 + \epsilon$$

$$Y = \beta_0 + \beta_1 X_1 + (\beta_2 + \beta_3 X_1) X_2 + \epsilon$$

The coefficient of one explanatory variable depends on the value of the other explanatory variable.

Variable Selection and Model Building

We usually want to choose a model that includes a subset of the available explanatory variables.

Two separate but related questions:

• How many explanatory variables should we use (i.e., subset size)? Smaller sets are more convenient, but larger sets may explain more of the variation (SS) in the response.

• Given the subset size, which variables should we choose?

Criteria for Model Selection

To determine an appropriate subset of the predictor variables, there are several different criteria available. We will go through them one at a time, noting their benefits and drawbacks. They include R^2 , adjusted R^2 , Mallow's C_p , MSE, PRESS, AIC, SBC. SAS will provide these statistics, so you should pay more attention to what they are good for than how they are computed. To obtain them from SAS, place after the model statement /selection = MAXR ADJRSQ CP. Note that the different criterion may not lead to the same model in every case.

R^2 and Adjusted R^2 (or MSE) Criterion

- The text uses $R_p^2 = R^2 = 1 \frac{SSE}{SSTO}$ (see page 354). Their subscript is just the number of variables in the associated model.
- The goal in model selection is to maximize this criterion. One MAJOR drawback to R^2 is that the addition of any variable to the model (significant or not) will increase R^2 (perhaps not enough to notice depending on the variable). At some point, added variables just get in the way!
- The Adjusted R^2 criterion penalizes the R^2 value based on the number of variables in the model. Hence it eventually starts decreasing as unnecessary variables are added.

$$R_a^2 = 1 - \left(\frac{n-1}{n-p}\right) \frac{SSE}{SSTO} \text{ (we end up subtracting off more as } p \text{ is increased)}$$

• Maximizing the Adjusted R^2 criterion is one way to select a model. As the text points out this is equivalent to minimizing the MSE since

$$R_a^2 = 1 - \left(\frac{n-1}{n-p}\right)\frac{SSE}{SSTO} = 1 - \frac{MSE}{SSTO/(n-1)} = 1 - \frac{MSE}{\text{constant}}$$

Mallow's C_p Criterion

- The basic idea is to compare subset models with the full model.
- The full model is good at prediction, but if there is multicollinearity our interpretations of the parameter estimates may not makes sense. A subset model is good if there is not substantial "bias" in the predicted values (relative to the full model).
- The C_p criterion looks at the ratio of error SS for the model with p variables to the MSE of the full model, then adds a penalty for the number of variables.

$$C_p = \frac{SSE_p}{MSE(Full)} - (n - 2p)$$

- SSE is based on a specific choice of p-1 variables (p is the number of regression coefficients *including the intercept*); while MSE is based on the full set of variables.
- A model is good according to this criterion if $C_p \leq p$. We may choose the smallest model for which $C_p \leq p$, so a benefit of this criterion is that it can achieve for us a "good" model containing as few variables as possible.
- One might also choose to pick the model that minimizes C_p .
- See page 357-359 for details.

PRESS Statistic

Stands for PREdiction Sums of Squares

Obtained by the following algorithm: For each observation i, delete the observation and predict Y for that observation using a model based on the n-1 cases. Then look at SS for the observed minus predicted.

$$PRESS_p = \sum (Y_i - \hat{Y}_{i(i)})^2$$

Models with small PRESS statistic are considered good candidates.

SBC and AIC

Criterion based on -2 * log(likelihood) plus a penalty for more complexity.

$$AIC -- \min n \log\left(\frac{SSE_p}{n}\right) + 2p$$

 $SBC -- \min n \log\left(\frac{SSE_p}{n}\right) + p \log(n)$

Note that different criteria will not give the identical answer.

Model Selection Methods

There are three commonly available. To apply them using SAS, you use the option selection = ***** after the model statement. The methods include

- Forward Selection (FORWARD) starts with the null model and adds variables one at a time.
- *Backward Elimination* (BACKWARD) starts with the full model and deletes variables one at a time.
- Forward Stepwise Regression (STEPWISE) starts with the null model and checks for adds/deletes at each step. This is probably the preferred method (see section on multicollinearity!). It is forward selection, but with a backward glance at each step.

These methods all add/delete variables based on Partial F-tests. (See page 364)

Some additional options in the model statement

INCLUDE=n forces the first n explanatory variables into all models BEST=n limits the output to the best n models of each subset size START=n limits output to models that include at least n explanatory variables

Ordering Models of the Same Subset Size

Use R^2 or SSE.

This approach can lead us to consider several models that give us approximately the same predicted values.

May need to apply knowledge of the subject matter to make a final selection.

If prediction is the key goal, then the choice of variables is not as important as if interpretation is the key.

Surgical Unit Example

- References: KNNL Section 9.2 (p350ff), nknw334.sas
- Y is the survival time
- Potential X's include Blood clotting score (X_1) , Prognostic Index (X_2) , Enzyme Function Test (X_3) , and Liver Function Test (X_4) .
- n = 54 patients were observed.
- Initial diagnostics note curved lines and non-constant variance, suggesting that Y should be transformed with a log. Take a look at the plots in the SAS file and play with some analyses on your own.

```
data surgical;
infile 'H:\System\Desktop\Ch08ta01.dat';
input blood prog enz liver surv;
```

IN	11.2			-	
blood 2.6	-				
	•	98 prog			
			enz 23		
				liver 0.74	
					surv 34

Take the log of survival

```
data surgical;
  set surgical;
  lsurv=log(surv);
proc reg data=surgical;
  model lsurv=blood prog enz liver/
  selection=rsquare cp aic sbc b best=3;
```

Number in Model R-Square C(p) AIC SBC 0.5274 787.9471 -87.3085 -83.33048 1 0.4424 938.6707 -78.3765 -74.39854 1 1 0.3515 1099.691 -70.2286 -66.25061 _____ 2 0.8129 283.6276 -135.3633 -129.39638 2 0.6865 507.8069 -107.4773 -101.51034 2 0.6496 573.2766 -101.4641 -95.49714 _____ 3 0.9723 3.0390 -236.5787 -228.62281 3 0.8829 161.6520 -158.6434 -150.68745 3 -111.4189 -103.46299 0.7192 451.8957 _____ 4 0.9724 5.0000 -234.6217 -224.67680

One model stands out: the first one with 3 variables $(C_p = 3.04 . The full model has <math>C_p = 5 = p$. The parameter estimates indicate that the desired model is the one with blood, prog and enz, but not liver.

Number in						
Model	R-Square	Intercept	blood	prog	enz	liver
1	0.5274	3.90609				0.42771
1	0.4424	3.55863			0.01973	
1	0.3515	3.68138	•	0.02211	•	
2	0.8129	2.08947	·	0.02271	0.02015	
2	0.6865	3.19784			0.01301	0.32010
2	0.6496	3.24325		0.01403		0.34596
3	0.9723	1.11358	0.15940	0.02140	0.02193	
3	0.8829	2.16970		0.01819	0.01612	0.18846
3	0.7192	2.68966	0.09239		0.01604	0.22556
4	0.9724	1.12536	0.15779	0.02131	0.02182	0.00442

In this particular example you would probably come to the same conclusion based on the Type II SS, but not on the individual correlations: *liver* has the *highest* individual correlation with *lsurv* (but also is correlated with the other three).

Below we see that this is the same model chosen by forward stepwise regression:

proc reg data=surgical;

model lsurv=blood prog enz liver / selection=stepwise;

			Summary of	f Stepwise S	Selection			
	Variable	Variable	Number	Partial	Model			
Step	Entered	Removed	Vars In	R-Square	R-Square	C(p)	F Value	Pr > F
1	liver		1	0.5274	0.5274	787.947	58.02	<.0001
2	enz		2	0.1591	0.6865	507.807	25.89	<.0001
3	prog		3	0.1964	0.8829	161.652	83.83	<.0001
4	blood		4	0.0895	0.9724	5.0000	158.65	<.0001
5		liver	3	0.0000	0.9723	3.0390	0.04	0.8442
4 5	blood	liver	4 3	0.0895 0.0000	0.9724 0.9723	5.0000 3.0390	158.65 0.04	< 0